

A Function of SmeDEF, the Major Quinolone Resistance Determinant of *Stenotrophomonas maltophilia*, Is the Colonization of Plant Roots

Guillermo García-León,^a Alvaro Hernández,^a Sara Hernando-Amado,^a Peyman Alavi,^b Gabriele Berg,^b José Luis Martínez^a

Departamento de Biotecnología Microbiana, Centro Nacional de Biotecnología, CSIC, Cantoblanco, Madrid, Spain^a; Graz University of Technology, Institute of Environmental Biotechnology, Graz, Austria^b

Quinolones are synthetic antibiotics, and the main cause of resistance to these antimicrobials is mutation of the genes encoding their targets. However, in contrast to the case for other organisms, such mutations have not been found in quinolone-resistant *Stenotrophomonas maltophilia* isolates, in which overproduction of the SmeDEF efflux pump is a major cause of quinolone resistance. SmeDEF is chromosomally encoded and highly conserved in all studied *S. maltophilia* strains; it is an ancient element that evolved over millions of years in this species. It thus seems unlikely that its main function would be resistance to quinolones, a family of synthetic antibiotics not present in natural environments until the last few decades. Expression of SmeDEF is tightly controlled by the transcriptional repressor SmeT. Our work shows that plant-produced flavonoids can bind to SmeT, releasing it from *smeDEF* and *smeT* operators. Antibiotics extruded by SmeDEF do not impede the binding of SmeT to DNA. The fact that plant-produced flavonoids specifically induce *smeDEF* expression indicates that they are *bona fide* effectors regulating expression of this resistance determinant. Expression of efflux pumps is usually downregulated unless their activity is needed. Since *smeDEF* expression is triggered by plant-produced flavonoids, we reasoned that this efflux pump may have a role in the colonization of plants by *S. maltophilia*. Our results showed that, indeed, deletion of *smeE* impairs *S. maltophilia* colonization of plant roots. Altogether, our results indicate that quinolone resistance is a recent function of SmeDEF and that colonization of plant roots is likely one original function of this efflux pump.

The acquisition of antibiotic resistance by bacterial pathogens is a relevant problem for current medical practice and is also one of the few evolutionary processes amenable to study on a human time scale. Development of resistance can be achieved either by mutation or by acquiring a resistance gene from another microorganism. Since most antibiotics have been isolated from environmental microorganisms (1), it was early suggested that resistance genes originated in natural microbial populations to counteract the inhibitory action of antibiotics present in natural ecosystems (2, 3).

Nevertheless, purely synthetic antimicrobials, such as quinolones, are also in use, and bacteria have developed resistance to these antibacterials. Since there are no quinolone producers in nature, it was supposed that resistance to this family of drugs would be due just to mutations in the genes coding for their targets, the bacterial topoisomerases (4). In the absence of quinolone selective pressure, environmental microorganisms would not require quinolone resistance, and consequently, quinolone resistance genes should not have evolved in nature. Further work showed, however, that this is not true; there are quinolone resistance genes conferring low-level resistance to these drugs via chromosomally encoded multidrug (MDR) efflux pumps (5, 6) or horizontally transferred resistance determinants, such as Qnr (7). However, it is true that despite these findings, mutations in genes encoding bacterial topoisomerases are still the main cause of high-level quinolone resistance in all studied microorganisms. The exception to this rule is *Stenotrophomonas maltophilia*, a free-living and often plant-associated opportunistic pathogen that is considered a prototype of intrinsically resistant bacteria (8, 9), which carries in its genome several operons encoding multidrug efflux pumps (10, 11). Different works have shown that, in contrast to what happens for other bacterial species, clinical quinolone-resistant *S. maltophilia* isolates do not display mutations in genes encoding bacterial topoisomerases (12, 13). One characteristic that

may explain this situation is that, in contrast to the case in other organisms, in which overproduction of efflux pumps usually renders low-level resistance to quinolones, in most cases below clinical breakpoints, the overproduction of the multidrug efflux pump SmeDEF confers high-level quinolone resistance to *S. maltophilia* (14, 15). Furthermore, it has been shown that SmeDEF overexpression is frequent among clinical isolates of *S. maltophilia* and that the level of expression of this efflux pump correlates well with the level of resistance to quinolones (16). Together with the analysis of mutants defective in this efflux pump (17), these results indicate that SmeDEF is an important element in the intrinsic quinolone resistance of *S. maltophilia* (18) and, upon its overexpression, is a major contributor to acquired resistance to that family of antibiotics by this bacterial species (16, 19).

As stated above, quinolones are synthetic antibiotics which were not present in environmental *S. maltophilia* habitats until the last few decades. On the other hand, the SmeDEF genes are intrinsic genes present in the genomes of all known *S. maltophilia* strains, where they have evolved during several million years, rather than being a recent acquisition via horizontal gene transfer. It is difficult to believe that the genes for this element evolved in the genome of *S. maltophilia* to counteract the activity of a family of antibiotics that did not exist in nature until very recently. Fur-

Received 10 April 2014 Accepted 7 May 2014

Published ahead of print 16 May 2014

Editor: M. Kivisaar

Address correspondence to José Luis Martínez, jlmrtnez@cnb.csic.es.

G.G.-L. and A.H. contributed equally to this article.

Copyright © 2014, American Society for Microbiology. All Rights Reserved.

doi:10.1128/AEM.01058-14

thermore, although *S. maltophilia* is a relevant nosocomial pathogen (8), its natural habitats are water and soil, where it can colonize all parts of plants, including the whole endosphere (20). Since efflux pumps of plant-interacting bacteria can be involved in plant colonization/invasion (21–23), we wondered whether SmeDEF may also play this role. Indeed, by using biochemical and functional assays, we found that this efflux pump, which is a major determinant of quinolone resistance in *S. maltophilia*, is involved in the endophytic colonization of plant roots.

MATERIALS AND METHODS

Bacterial strains and growing conditions. The bacterial strains used were the wild-type *S. maltophilia* isolate D457 (15) and its isogenic derivatives MBS411 (D457 Δ *smeE*) and D457R (D457 overexpressing SmeDEF) (14, 24). All strains were grown in LB medium at 37°C, unless indicated otherwise.

In silico prediction of the interactions of SmeT with plant exudates. Docking predictions of the interactions of plant flavonoid exudates (naringenin, phloretin, genistein, apigenin, and quercetin) with SmeT were performed using AutoDock4 software (25), considering the flexibility of the different ligands but not the protein. The docking area was restricted to the protein's binding pocket, and results were analyzed using AutodockTools 1.5.4. Protein-ligand complexes were visualized using PyMol (26).

DNA labeling and EMSA in the presence of plant flavonoid exudates or antibiotics. The probe for the electrophoretic mobility shift assay (EMSA) was a 158-bp double-stranded DNA fragment containing the SmeT operator site, and 5'-end labeled with [γ -³²P]dATP, which was obtained as previously described (27). This probe was incubated with purified SmeT (0.2 μ M) in binding buffer [10 mM Tris-HCl, 50 mM KCl, 10 mM MgCl₂, 1 mM EDTA, pH 7.2, 50 μ g/ml bovine serum albumin, 1 mM dithiothreitol, 5% (vol/vol) glycerol, and 100 μ g/ml poly(dI:dC)], as a nonspecific competitor DNA for 20 min at room temperature. Later, increasing concentrations (0, 0.06, 0.12, 0.25, 0.5, 1, and 2 mM) of commercially available pure flavonoids (naringenin, phloretin, coumestrol, genistein, apigenin, and quercetin) or a 3 mM concentration of antibiotics (chloramphenicol, erythromycin, and ciprofloxacin) was added and incubation continued for 15 min more. Retarded complexes were separated in a 6% nondenaturing polyacrylamide gel (37.5:1 acrylamide:bisacrylamide). Electrophoresis was performed at room temperature with 89 mM Tris-borate and 2 mM EDTA buffer for 90 min at 100 V, and the gel was dried before autoradiography.

RNA preparation and real-time RT-PCR. Fifteen microliters of an *S. maltophilia* D457 overnight culture was used to inoculate 15 ml of LB medium containing subinhibitory concentrations of plant exudates (15 μ g/ml of phloretin and 10 μ g/ml of naringenin, quercetin, apigenin, and genistein) until an optical density at 600 nm (OD₆₀₀) of \approx 1.0 was reached; cells were then spun down at 6,000 \times g for 10 min at 4°C and immediately frozen on dry ice and stored at -80°C. Total RNA extraction, DNA elimination, RNA integrity verification, DNA absence confirmation, cDNA generation, and real-time PCR were performed as described previously (28). Total RNA was extracted from cell pellets by use of an RNeasy mini-kit (Qiagen) according to the manufacturer's instructions. To further eliminate any remaining DNA, Turbo DNA-free (Ambion) was used. RNA integrity was verified on a 1% agarose gel, and the absence of DNA was verified by real-time PCR using primers GyrA-RT.fw and GyrA-RT.rv (Table 1). cDNA was obtained from 1 μ g RNA by use of a High Capacity cDNA reverse transcription (RT) kit (AB Applied Biosystems). RT-PCR was performed according to the method of Morales et al. (29). Briefly, the first denaturation step, 95°C for 10 min, was followed by 40 temperature cycles (95°C for 15 s, 60°C for 1 min) for amplification and quantification. The GyrA-RT.fw and GyrA-RT.rv primers were used to amplify the housekeeping gene *gyrA* as described previously (30) (Table 1). Differences in the relative amounts of mRNA for the different genes were determined according to the 2^{- $\Delta\Delta$ CT} method (31). Real-time RT-PCRs were performed using a Power SYBR green kit (Applied Biosystems) as indi-

TABLE 1 Oligonucleotides used in this work

Oligonucleotide	Sequence	Utilization
GyrA-RT.fw	CCAGGGTAACTTCGGTTCGA	Internal control gene for RT-PCR
GyrA-RT.rv	GCCTCGGTGTATCGCATTG	
SmeD-RT.fw	CGGTACAGCATCCTGATGGA	RT-PCR for <i>smeD</i>
SmeD-RT.rv	TCAACGCTGACTTCGGAGAAGT	
SmeY-RT.fw	AGCTGCTGTTCTCCGGTATCA	RT-PCR for <i>smeY</i>
SmeY-RT.rv	CACCAGGATGCGCAGGAT	
SmeI-RT.fw	ACTGCGATGAACACCGTTACC	RT-PCR for <i>smeI</i>
SmeI-RT.rv	CACGTCACCCTGCTTCACTTC	
SmeA-RT.fw	ACCCGGTCTATCTGGATGTG	RT-PCR for <i>smeA</i>
SmeA-RT.rv	CTCATGGGCATAGGTGCTG	
SmeG-RT.fw	AAGAACGTGAAGACCGATGG	RT-PCR for <i>smeG</i>
SmeG-RT.rv	CCTTCCTGACCTTCTGCAC	
SmeM-RT.fw	ATTTCCAGGTGCCTGAAGTG	RT-PCR for <i>smeM</i>
SmeM-RT.rv	CGCATCAATGGTGCTGAC	
SmeO-RT.fw	CAGGAAAGTCCACTGTCGTTTC	RT-PCR for <i>smeO</i>
SmeO-RT.rv	CACGTCGCCCTTCTTCCAC	
SmeV-RT.fw	CCAGGGTAACTTCGGTTCGA	RT-PCR for <i>smeV</i>
SmeV-RT.rv	GCCTCGGTGTATCGCATTG	

cated by the manufacturer. In all cases, the mean values for relative mRNA expression obtained in three independent triplicate experiments were considered. Student's *t* test was used for statistical analysis of the results.

In vivo plant colonization experiments. The oilseed rape plant (cv. Californium; Kwizda, Austria) was used for *in vivo* assays. Before inoculation of seeds with bacteria, the surfaces of the seeds were sterilized using a modification of the seed infiltration approach described by Müller and Berg (32). In brief, seeds were treated with a 3% sodium hypochlorite (NaClO) solution for 5 min and subsequently washed three times with sterile distilled water. The numbers of CFU ml⁻¹ of *S. maltophilia* strains used in the colonization experiments were determined by diluting and plating 100- μ l aliquots of overnight cultures of each bacterial strain on LB plates. Colonies were counted after incubation of the plates at 37°C for 24 h. Surface-sterilized seeds were placed in a petri dish with 10 ml of 0.85% NaCl solution containing 10⁶ CFU/ml of each bacterial culture on an orbital shaker at 85 rpm and room temperature for 4 h.

The capability of *S. maltophilia* strains D457, MBS411, and D57R to colonize plant rhizospheres was checked by use of seed germination pouches (Mega International, MN). Autoclaved pouches were loaded with 5 inoculated seeds each and moisturized with 20 ml of sterile distilled water. Pouches were placed in sterile, covered polypropylene containers to avoid contamination and dehydration and then incubated in a greenhouse at 23 \pm 2°C under artificial lighting (16-h light period) for 7 days. Bacteria were reisolated from 7-day-old oilseed rape plants. To achieve this purpose, root sections were sampled into sterile Whirl-Pak bags (Carl Roth, Germany), supplemented with 5 ml of 0.85% NaCl solution, and roughly disintegrated using a mortar and pestle. Serial dilutions of the extract were then plated onto LB plates. After incubation for 24 h at 37°C, colonies were counted and the number of CFU/g root was assessed.

FISH. Root sections of oilseed rape plants were fixed with 4% paraformaldehyde-phosphate-buffered saline (PBS) (3:1 [vol/vol]). The fixed samples were then stored in PBS-96% ethanol (1:1) at -20°C. In-tube fluorescent *in situ* hybridization (FISH) was performed as described earlier (33), using commercially available FISH probes purchased from genXpress (Wiener Neudorf, Austria). An equimolar ratio of FISH probes EUC I, EUC II, and EUC III, labeled with the fluorescent dye Cy3, was used to detect the analyzed *S. maltophilia* strains. The hybridization step was carried out using 35% formamide in a 46°C water bath for 90 min. After hybridization, the samples were washed at 48°C for 15 min. Microscopic images of the samples were captured using a Leica TCS SPE confocal microscope (Leica Microsystems, Wetzlar, Germany). A Leica ACS




FIG 3 The presence of plant exudates triggers *smeD* expression. The amounts of *smeD* mRNA in the presence of plant exudates were measured by real-time RT-PCR, and the fold changes estimated with respect to the value determined for the wild-type strain (wt) grown in the absence of exudates are shown. As shown, mRNA levels were increased in the presence of apigenin (Api), genistein (Gen), naringenin (Nar), phloretin (Phl), and quercetin (Que). D457R is a mutant strain in which expression of *smeDEF* is fully derepressed due to a mutation that inactivates SmeT.

To ascertain whether the capability of binding to SmeT correlates with *smeDEF* induction, expression of *smeD* was measured in the presence and absence of plant exudates. As shown in Fig. 3, plant exudates triggered *smeD* expression (in all cases, $P < 0.05$), which supports the hypothesis that these compounds are *bona fide* effectors of SmeT.

To analyze if the observed induction was specific for SmeDEF, the effect of phloretin, the flavonoid triggering the highest induction levels, on the expression of the previously described *S. maltophilia* tripartite efflux pumps (10) was analyzed by measuring the expression of the first gene of each of the operons, namely, *smeA*, *smeD*, *smeV*, *smeY*, *smeG*, *smeM*, *smeO*, and *smeI*. As shown in Fig. 4, phloretin preferentially induced *smeD* ($P = 0.027$), and a minor




FIG 4 Effects of phloretin on the expression of *S. maltophilia* operons encoding multidrug efflux pumps. The mRNA levels of the first genes of the operons encoding MDR determinants in the chromosome of *S. maltophilia* (10), i.e., *smeA*, *smeD*, *smeV*, *smeY*, *smeG*, *smeM*, *smeO*, and *smeI*, were analyzed by real-time RT-PCR with the wild-type D457 strain (black) and the D457R strain (white), a mutant strain in which expression of *smeDEF* is fully derepressed due to a mutation that inactivates SmeT. The expression in the presence of phloretin was measured by real-time RT-PCR with D457 (gray) and D457R (stripes). The fold changes were estimated with respect to the value determined for the wild-type strain grown in the absence of the flavonoid.




FIG 5 Overexpression of SmeDEF reduces the susceptibility of *S. maltophilia* to phloretin. The figure shows the growth of the wild-type strain and mutants either overproducing or lacking *smeE*. (A) Growth in the absence of phloretin. (B) Growth in the presence of phloretin at 300 $\mu\text{g/ml}$. As shown, the susceptibility to phloretin correlates with the SmeDEF expression level.

effect was also observed for *smeO* ($P = 0.048$), and maybe for *smeG* (not statistically significant; $P = 0.16$), while the expression of the other efflux pumps did not change in the presence of the flavonoid. The D457R mutant lacks a functional SmeT repressor, and this causes *smeDEF* overexpression. Since phloretin did not further increase the expression of any of the tested efflux pumps in this mutant (Fig. 4), most likely the induction observed in the wild-type strain is attributable to SmeT, a feature in agreement with the EMSA results displayed in Fig. 2. Although *smeG* and *smeO* are not regulated by SmeT, the regulators of both operons in which these genes are present belong to the TetR family of transcriptional repressors. The induction by phloretin of the expression of *smeO* (and maybe *smeG*) suggests that this flavonoid may bind other TetR-type regulators besides SmeT. In line with this reasoning, previous work showed that phloretin binds TtgR, a TetR regulator homologous to SmeT (44). Since phloretin induces *smeDEF* expression, it might be possible that it is extruded by SmeDEF, which would make the D457R mutant overproducing this efflux pump more resistant and a mutant lacking the efflux pump more susceptible to the presence of high concentrations of the flavonoid. To analyze this possibility, the wild-type strain and mutants lacking *smeE* or presenting constitutive high-level SmeDEF expression were grown in the absence or presence of phloretin (300 $\mu\text{g/ml}$). Consistent with previously published information (40), SmeDEF overexpression conferred a fitness cost to *S. maltophilia* in the absence of phloretin (Fig. 5A). However, in the presence of phloretin, the mutant overexpressing SmeDEF




FIG 6 Colonization of oilseed rape roots by *S. maltophilia*. Colonization was visualized by FISH. (A) Wild-type strain D457. (C) *smeE*-defective mutant strain MBS411. (E) D457R mutant that overexpresses SmeDEF. 3D analysis of the CLSM stacks of D457 (B), MBS411 (D), and D457R (F) was performed with Imaris 7.0 software. (G) Quantification of the numbers of cells colonizing the roots. Results shown are mean values for three independent replicates.

presented normal growth, whereas the wild-type strain was impaired and the mutant lacking *smeE* did not grow. These results strongly suggest that phloretin is a SmeDEF substrate.

As mentioned earlier, MDR efflux pumps are expressed at their highest levels only when needed. Bacteria can determine this need by sensing their environment and identifying the proper effectors. Since the effectors triggering SmeDEF overexpression are plant exudates, not antibiotics, it is conceivable that one original function of this efflux pump was the interaction with plant roots. To study this possibility, we measured the capability for endophytic colonization of an *S. maltophilia* mutant lacking *smeE* in comparison with its wild-type parental counterpart. The D457R mutant, which overexpresses SmeDEF, was also used as a control. As shown in Fig. 6, the mutant lacking *smeE* was impaired in colonizing plant roots ($P = 0.0000000046$), whereas the differences observed between the D457R mutant and the wild-type strain were not significant ($P = 0.082$). This indicates that SmeDEF is required for achieving a normal colonization level. The fact that the D457R mutant does not present enhanced colonization capabilities can be explained by three different (not mutually exclusive) hypotheses. (i) It has been shown that SmeDEF overexpression compromises *in vitro* *S. maltophilia* growth (Fig. 5A) (40); the potential benefits on colonization will be compensated by such impairment. (ii) When *S. maltophilia* grows on plants, the presence of flavonoids induces *smeDEF* in the wild-type strain, to levels similar to those observed for D457R. (iii) The colonization of the plant requires a given threshold of SmeDEF expression, and

increasing the level does not enhance colonization. Altogether, our results support the hypothesis that one original function for which SmeDEF has been selected in *S. maltophilia* is colonization of plants, not resistance against the action of antibiotics such as quinolones.

Our results indicate that antibiotic resistance genes, which are causing problems at hospitals, have not necessarily always evolved in the field to counteract the production of antibiotics by surrounding competitors (45, 46). Rather, enzymes or efflux pumps with substrates similar to an antibiotic can render resistance when the antimicrobial is present. However, this should not downplay their role as significant antibiotic resistance elements but solely points out that, on occasion, they evolved for other purposes in nature. It has been stated that the amounts of antibiotics in natural ecosystems are likely very low for a general role as inhibitors (47–51). However, the wide use of antibiotics by human beings has changed this situation, and there are different ecosystems, including individuals under medical treatment, where antibiotics constitute a highly relevant selective pressure (48).

ACKNOWLEDGMENTS

Work in our laboratory is supported by grant BIO2011-25255 from the Spanish Ministry of Economy and Competitiveness, grant S2010/BMD2414 (PROMPT) from CAM, the Spanish Network for Research on Infectious Diseases (grant REIPI RD12/0015) via the Instituto de Salud Carlos III, and grant HEALTH-F3-2011-282004 (EVOTAR) from the European Union. G.G.-L. has been the recipient of an FPI fellowship.

REFERENCES

- Waksman SA, Woodruff HB. 1940. The soil as a source of microorganisms antagonistic to disease-producing bacteria. *J. Bacteriol.* 40:581–600.
- Benveniste R, Davies J. 1973. Aminoglycoside antibiotic-inactivating enzymes in actinomycetes similar to those present in clinical isolates of antibiotic-resistant bacteria. *Proc. Natl. Acad. Sci. U. S. A.* 70:2276–2280. <http://dx.doi.org/10.1073/pnas.70.8.2276>.
- Davies JE. 1997. Origins, acquisition and dissemination of antibiotic resistance determinants. *Ciba Found. Symp.* 207:15–27.
- Hernandez A, Sanchez MB, Martinez JL. 2011. Quinolone resistance: much more than predicted. *Front. Microbiol.* 2:22. <http://dx.doi.org/10.3389/fmicb.2011.00022>.
- Cohen SP, McMurry LM, Hooper DC, Wolfson JS, Levy SB. 1989. Cross-resistance to fluoroquinolones in multiple-antibiotic-resistant (Mar) *Escherichia coli* selected by tetracycline or chloramphenicol: decreased drug accumulation associated with membrane changes in addition to OmpF reduction. *Antimicrob. Agents Chemother.* 33:1318–1325. <http://dx.doi.org/10.1128/AAC.33.8.1318>.
- Hooper DC. 1999. Mechanisms of fluoroquinolone resistance. *Drug Resist. Updat.* 2:38–55. <http://dx.doi.org/10.1054/drup.1998.0068>.
- Martinez-Martinez L, Pascual A, Jacoby GA. 1998. Quinolone resistance from a transferable plasmid. *Lancet* 351:797–799. [http://dx.doi.org/10.1016/S0140-6736\(97\)07322-4](http://dx.doi.org/10.1016/S0140-6736(97)07322-4).
- Brooke JS. 2012. *Stenotrophomonas maltophilia*: an emerging global opportunistic pathogen. *Clin. Microbiol. Rev.* 25:2–41. <http://dx.doi.org/10.1128/CMR.00019-11>.
- Sanchez MB, Hernandez A, Martinez JL. 2009. *Stenotrophomonas maltophilia* drug resistance. *Future Microbiol.* 4:655–660. <http://dx.doi.org/10.2217/fmb.09.45>.
- Crossman LC, Gould VC, Dow JM, Vernikos GS, Okazaki A, Sebahia M, Saunders D, Arrowsmith C, Carver T, Peters N, Adlem E, Kerhounou A, Lord A, Murphy L, Seeger K, Squares R, Rutter S, Quail MA, Rajandream MA, Harris D, Churcher C, Bentley SD, Parkhill J, Thomson NR, Avison MB. 2008. The complete genome, comparative and functional analysis of *Stenotrophomonas maltophilia* reveals an organism heavily shielded by drug resistance determinants. *Genome Biol.* 9:R74. <http://dx.doi.org/10.1186/gb-2008-9-4-r74>.
- Rocco F, De Gregorio E, Colonna B, Di Nocera PP. 2009. *Stenotrophomonas maltophilia* genomes: a start-up comparison. *Int. J. Med. Microbiol.* 299:535–546. <http://dx.doi.org/10.1016/j.ijmm.2009.05.004>.
- Ribera A, Domenech-Sanchez A, Ruiz J, Benedi VJ, Jimenez de Anta MT, Vila J. 2002. Mutations in *gyrA* and *parC* QRDs are not relevant for quinolone resistance in epidemiological unrelated *Stenotrophomonas maltophilia* clinical isolates. *Microb. Drug Resist.* 8:245–251. <http://dx.doi.org/10.1089/10766290260469499>.
- Valdezate S, Vindel A, Saez-Nieto JA, Baquero F, Canton R. 2005. Preservation of topoisomerase genetic sequences during in vivo and in vitro development of high-level resistance to ciprofloxacin in isogenic *Stenotrophomonas maltophilia* strains. *J. Antimicrob. Chemother.* 56:220–223. <http://dx.doi.org/10.1093/jac/dki182>.
- Alonso A, Martinez JL. 2000. Cloning and characterization of SmeDEF, a novel multidrug efflux pump from *Stenotrophomonas maltophilia*. *Antimicrob. Agents Chemother.* 44:3079–3086. <http://dx.doi.org/10.1128/AAC.44.11.3079-3086.2000>.
- Alonso A, Martinez JL. 1997. Multiple antibiotic resistance in *Stenotrophomonas maltophilia*. *Antimicrob. Agents Chemother.* 41:1140–1142.
- Alonso A, Martinez JL. 2001. Expression of multidrug efflux pump SmeDEF by clinical isolates of *Stenotrophomonas maltophilia*. *Antimicrob. Agents Chemother.* 45:1879–1881. <http://dx.doi.org/10.1128/AAC.45.6.1879-1881.2001>.
- García-León G, Sanchez MB, Martinez JL. 2012. The inactivation of intrinsic antibiotic resistance determinants widens the mutant selection window for quinolones in *Stenotrophomonas maltophilia*. *Antimicrob. Agents Chemother.* 56:6397–6399. <http://dx.doi.org/10.1128/AAC.01558-12>.
- Zhang L, Li XZ, Poole K. 2001. SmeDEF multidrug efflux pump contributes to intrinsic multidrug resistance in *Stenotrophomonas maltophilia*. *Antimicrob. Agents Chemother.* 45:3497–3503. <http://dx.doi.org/10.1128/AAC.45.12.3497-3503.2001>.
- Sanchez P, Alonso A, Martinez JL. 2004. Regulatory regions of *smeDEF* in *Stenotrophomonas maltophilia* strains expressing different amounts of the multidrug efflux pump SmeDEF. *Antimicrob. Agents Chemother.* 48:2274–2276. <http://dx.doi.org/10.1128/AAC.48.6.2274-2276.2004>.
- Zhu B, Liu H, Tian WX, Fan XY, Li B, Zhou XP, Jin GL, Xie GL. 2012. Genome sequence of *Stenotrophomonas maltophilia* RR-10, isolated as an endophyte from rice root. *J. Bacteriol.* 194:1280–1281. <http://dx.doi.org/10.1128/JB.06702-11>.
- Burse A, Weingart H, Ullrich MS. 2004. NorM, an *Erwinia amylovora* multidrug efflux pump involved in *in vitro* competition with other epiphytic bacteria. *Appl. Environ. Microbiol.* 70:693–703. <http://dx.doi.org/10.1128/AEM.70.2.693-703.2004>.
- Martinez JL, Sanchez MB, Martinez-Solano L, Hernandez A, Garmendia L, Fajardo A, Alvarez-Ortega C. 2009. Functional role of bacterial multidrug efflux pumps in microbial natural ecosystems. *FEMS Microbiol. Rev.* 33:430–449. <http://dx.doi.org/10.1111/j.1574-6976.2008.00157.x>.
- Palumbo JD, Kado CI, Phillips DA. 1998. An isoflavonoid-inducible efflux pump in *Agrobacterium tumefaciens* is involved in competitive colonization of roots. *J. Bacteriol.* 180:3107–3113.
- Sanchez P, Alonso A, Martinez JL. 2002. Cloning and characterization of SmeT, a repressor of the *Stenotrophomonas maltophilia* multidrug efflux pump SmeDEF. *Antimicrob. Agents Chemother.* 46:3386–3393. <http://dx.doi.org/10.1128/AAC.46.11.3386-3393.2002>.
- Morris GM, Huey R, Lindstrom W, Sanner MF, Belew RK, Goodsell DS, Olson AJ. 2009. AutoDock4 and AutoDockTools4: automated docking with selective receptor flexibility. *J. Comput. Chem.* 30:2785–2791. <http://dx.doi.org/10.1002/jcc.21256>.
- DeLano WL. 2002. The PyMOL molecular graphics system. <http://www.pymol.org>.
- Hernandez A, Mate MJ, Sanchez-Diaz PC, Romero A, Rojo F, Martinez JL. 2009. Structural and functional analysis of SmeT, the repressor of the *Stenotrophomonas maltophilia* multidrug efflux pump SmeDEF. *J. Biol. Chem.* 284:14428–14438. <http://dx.doi.org/10.1074/jbc.M809221200>.
- Olivares J, Alvarez-Ortega C, Linares JF, Rojo F, Kohler T, Martinez JL. 2012. Overproduction of the multidrug efflux pump MexEF-OprN does not impair *Pseudomonas aeruginosa* fitness in competition tests, but produces specific changes in bacterial regulatory networks. *Environ. Microbiol.* 14:1968–1981. <http://dx.doi.org/10.1111/j.1462-2920.2012.02727.x>.
- Morales G, Ugidos A, Rojo F. 2006. Inactivation of the *Pseudomonas putida* cytochrome o ubiquinol oxidase leads to a significant change in the transcriptome and to increased expression of the CIO and *cbb3-1* terminal oxidases. *Environ. Microbiol.* 8:1764–1774. <http://dx.doi.org/10.1111/j.1462-2920.2006.01061.x>.
- Gould VC, Avison MB. 2006. SmeDEF-mediated antimicrobial drug resistance in *Stenotrophomonas maltophilia* clinical isolates having defined phylogenetic relationships. *J. Antimicrob. Chemother.* 57:1070–1076. <http://dx.doi.org/10.1093/jac/dkl106>.
- Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta Ct}$ method. *Methods* 25:402–408. <http://dx.doi.org/10.1006/meth.2001.1262>.
- Müller H, Berg G. 2008. Impact of formulation procedures on the effect of the biocontrol agent *Serratia plymuthica* HRO-C48 on *Verticillium wilt* in oilseed rape. *BioControl* 53:905–916. <http://dx.doi.org/10.1007/s10526-007-9111-3>.
- Cardinale M, Vieira de Castro J, Jr, Muller H, Berg G, Grube M. 2008. *In situ* analysis of the bacterial community associated with the reindeer lichen *Cladonia arbuscula* reveals predominance of *Alphaproteobacteria*. *FEMS Microbiol. Ecol.* 66:63–71. <http://dx.doi.org/10.1111/j.1574-6941.2008.00546.x>.
- Li XZ, Nikaido H. 2009. Efflux-mediated drug resistance in bacteria: an update. *Drugs* 69:1555–1623. <http://dx.doi.org/10.2165/11317030-000000000-00000>.
- Toone EJ, Nikaido H. 2011. Structure and mechanism of RND-type multidrug efflux pumps. *Adv. Enzymol. Relat. Areas Mol. Biol.* 77:1–60. <http://dx.doi.org/10.1002/9780470920541.ch1>.
- Piddock LJ. 2006. Multidrug-resistance efflux pumps—not just for resistance. *Nat. Rev. Microbiol.* 4:629–636. <http://dx.doi.org/10.1038/nrmicro1464>.
- Yu EW, McDermott G, Zgurskaya HI, Nikaido H, Koshland DE, Jr. 2003. Structural basis of multiple drug-binding capacity of the AcrB multidrug efflux pump. *Science* 300:976–980. <http://dx.doi.org/10.1126/science.1083137>.
- Grkovic S, Brown MH, Skurray RA. 2001. Transcriptional regulation of multidrug efflux pumps in bacteria. *Semin. Cell Dev. Biol.* 12:225–237. <http://dx.doi.org/10.1006/scdb.2000.0248>.
- Grkovic S, Brown MH, Skurray RA. 2002. Regulation of bacterial drug export systems. *Microbiol. Mol. Biol. Rev.* 66:671–701. <http://dx.doi.org/10.1128/MMBR.66.4.671-701.2002>.

40. Alonso A, Morales G, Escalante R, Campanario E, Sastre L, Martinez JL. 2004. Overexpression of the multidrug efflux pump SmeDEF impairs *Stenotrophomonas maltophilia* physiology. *J. Antimicrob. Chemother.* 53:432–434. <http://dx.doi.org/10.1093/jac/dkh074>.
41. Linares JF, Lopez JA, Camafeita E, Albar JP, Rojo F, Martinez JL. 2005. Overexpression of the multidrug efflux pumps MexCD-OprJ and MexEF-OprN is associated with a reduction of type III secretion in *Pseudomonas aeruginosa*. *J. Bacteriol.* 187:1384–1391. <http://dx.doi.org/10.1128/JB.187.4.1384-1391.2005>.
42. Sanchez P, Linares JF, Ruiz-Diez B, Campanario E, Navas A, Baquero F, Martinez JL. 2002. Fitness of in vitro selected *Pseudomonas aeruginosa* *nalB* and *nfxB* multidrug resistant mutants. *J. Antimicrob. Chemother.* 50:657–664. <http://dx.doi.org/10.1093/jac/dkf185>.
43. Hernandez A, Ruiz FM, Romero A, Martinez JL. 2011. The binding of triclosan to SmeT, the repressor of the multidrug efflux pump SmeDEF, induces antibiotic resistance in *Stenotrophomonas maltophilia*. *PLoS Pathog.* 7:e1002103. <http://dx.doi.org/10.1371/journal.ppat.1002103>.
44. Alguel Y, Meng C, Teran W, Krell T, Ramos JL, Gallegos MT, Zhang X. 2007. Crystal structures of multidrug binding protein TtgR in complex with antibiotics and plant antimicrobials. *J. Mol. Biol.* 369:829–840. <http://dx.doi.org/10.1016/j.jmb.2007.03.062>.
45. Martinez JL. 2008. Antibiotics and antibiotic resistance genes in natural environments. *Science* 321:365–367. <http://dx.doi.org/10.1126/science.1159483>.
46. Martinez JL, Fajardo A, Garmendia L, Hernandez A, Linares JF, Martinez-Solano L, Sanchez MB. 2009. A global view of antibiotic resistance. *FEMS Microbiol. Rev.* 33:44–65. <http://dx.doi.org/10.1111/j.1574-6976.2008.00142.x>.
47. Davies J. 2006. Are antibiotics naturally antibiotics? *J. Ind. Microbiol. Biotechnol.* 33:496–499. <http://dx.doi.org/10.1007/s10295-006-0112-5>.
48. Fajardo A, Linares JF, Martinez JL. 2009. Towards an ecological approach to antibiotics and antibiotic resistance genes. *Clin. Microbiol. Infect.* 15(Suppl 1):S14–S16. <http://dx.doi.org/10.1111/j.1469-0691.2008.02688.x>.
49. Linares JF, Gustafsson I, Baquero F, Martinez JL. 2006. Antibiotics as intermicrobial signaling agents instead of weapons. *Proc. Natl. Acad. Sci. U. S. A.* 103:19484–19489. <http://dx.doi.org/10.1073/pnas.0608949103>.
50. Yim G, Wang HH, Davies J. 2007. Antibiotics as signalling molecules. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 362:1195–1200. <http://dx.doi.org/10.1098/rstb.2007.2044>.
51. Yim G, Wang HH, Davies J. 2006. The truth about antibiotics. *Int. J. Med. Microbiol.* 296:163–170. <http://dx.doi.org/10.1016/j.ijmm.2006.01.039>.