

# ETHYLENE TREATED KIWI FRUITS DURING STORAGE. PART I: POSTHARVEST BIOACTIVE, ANTIOXIDANT AND BINDING PROPERTIES

MARTIN POLOVKA<sup>1</sup>, YONG SEO PARK<sup>2,8</sup>, KYUNG-SIK HAM<sup>3</sup>, YANG-KYUN PARK<sup>3</sup>,  
SUCHADA VEARASILP<sup>4</sup>, PAWEŁ PASKO<sup>5</sup>, FERNANDO TOLEDO<sup>6</sup>, PATRICIA ARANCIBIA-AVILA<sup>6</sup> and  
SHELA GORINSTEIN<sup>7,8</sup>

<sup>1</sup>Department of Chemistry and Food Analysis, National Agricultural and Food Centre VUP, Food Research Institute, Bratislava, SK-824 75, Slovak Republic

<sup>2</sup>Department of Horticultural Science, Mokpo National University, Muan, Jeonnam, South Korea

<sup>3</sup>Department of Food Engineering, Mokpo National University, Muan, Jeonnam, South Korea

<sup>4</sup>Postharvest Technology Research Institute/Postharvest Technology Innovation Center, Chiang Mai University, Chiang Mai, 50200, Thailand

<sup>5</sup>Department of Food Chemistry and Nutrition, Medical College, Jagiellonian University, Krakow, 30-688, Poland

<sup>6</sup>Department of Basic Sciences, Universidad Del Bio-Bio, P.O. Box 447, Chillan, 3780000, Chile

<sup>7</sup>The Institute for Drug Research, School of Pharmacy, The Hebrew University, Hadassah Medical School, Jerusalem, 91120, Israel

<sup>8</sup>Corresponding authors.

TEL: 82 61 450 2376;

FAX: 82 61 453 2309;

EMAIL: ypark@mokpo.ac.kr or

TEL: 972 2 6758690;

FAX: 972 2 6757076;

EMAIL: shela.gorin@mail.huji.ac.il

Received for Publication February 19, 2016

Accepted for Publication April 17, 2016

doi:10.1111/jfpp.13084

†This article was written in memory of my dear brother Prof. Simon Trakhtenberg, who died in November 2011, who encouraged me and our research group during all his life.

## ABSTRACT

Kiwi fruits “Hayward” were submitted to ethylene treatment during 24 h, following by storage at 20C for 10 days. Significant differences were found in polyphenols and in the antioxidant capacities in conventional, low chemical and organic kiwi fruits. Ethylene treatment increased the bioactivity of organic, low chemical and conventional kiwi fruit. The antioxidant values for organic fruits were significantly higher than for conventional and nontreated samples. All investigated kiwi fruits showed a high level of correlation between the contents of phenolic compounds, their antioxidant and binding values. The statistical evaluation of bioactivity demonstrated that cultivation system and ethylene treatment following by storage have the potential to enhance the accumulation of health-beneficial food compounds in kiwi fruit and changes of the binding properties.

## PRACTICAL APPLICATIONS

The article describes the impact of three treatments including different agricultural systems, ethylene treatment and storage and biochemical changes in kiwi fruit. Agricultural system produced three different kiwi fruits: organic, semi-organic and conventional. Different cultivation systems affect the polyphenols and antioxidant status of kiwi fruit. During the ethylene treatment and storage the antioxidant status and binding properties have increased. In general these treatments affect the contents of the main bioactive compounds which have high nutritional value. In practice we can select the best treatment and its duration in connection with the best quality of the final product. This could be of economical importance for the kiwi fruit industry in development of new products with long shelf life and high quality of the final product.

## INTRODUCTION

Fruit and vegetable intake is associated with lower risk of chronic diseases as a result of consumption of antioxidant substances. Increased consumption of fruits protects cardiovascular diseases, because of the unique composition

of green kiwi fruit, which has the potential to benefit cardiovascular disease risk (Lee *et al.* 2010; Gammon *et al.* 2012). Organic foods have higher antioxidant capacity, because this form of agricultural management could induce synthesis of secondary compounds such as

polyphenols (Faller and Fialho 2010). Interest in organically produced food is increasing in response to concerns about intensive agricultural practices and their potential effect on human health as well as on the environment (Roghelia and Patel 2013). The effect of cultivation system on phenolic profile and antioxidant capacity in black and red currants, blueberries, tomatoes was evaluated (Wang *et al.* 2008; Wojdylo *et al.* 2013; Pataro *et al.* 2015). Kiwi fruit (*Actinidia deliciosa*) “Hayward,” grown under different conditions, is one of the most known cultivars. The effect of cultivation systems and fruit postharvest management on the antioxidant properties of fruits and vegetables was investigated (Migliori *et al.* 2012). Postharvest performance of organic and conventional “Hayward” kiwi fruits grown on the same farm in Marysville, California, and harvested at the same maturity stage was compared (Amodio *et al.* 2007). “Hayward” treated with ethylene or 1-methylcyclopropene (1-MCP) was compared with nontreated in a number of reports (Park *et al.* 2006; Jhalegar *et al.* 2011; Vieira *et al.* 2010).  $\text{CaCl}_2$  sprays significantly enhanced fruit Ca concentration and improved fruit quality during cold storage and shelf life (Gang *et al.* 2015). However, the main properties of treated and storing kiwi fruit samples were not investigated. The subject of this research was a comparison of the main kiwi fruit properties after ethylene treatment, following storage. To answer these questions, advanced analytical methods were used in combination with statistical calculations.

## MATERIALS AND METHODS

### Fruit Samples

Kiwi fruit cultivar “Hayward” was grown under conventional, low chemical and organic conditions in the orchard Heanam County (longitude 126° 15' and latitude 34° 18'), Jeonnam province, Korea. The harvest date was October 30, 2014 and October 18, 2015. Fruits with defects were discarded and 50 good fruits (80–100 g) were placed in five replicated glass jars. These healthy fruits were divided into two groups, the first group was treated with 100 ppm ethylene for 24 h at 20C, and the second was not treated (control). The fruits were put into an 18 L glass jar and ventilated with humidified flow of air or air mixed with ethylene of 300 mL/mL. Thereafter, the ethylene treated fruits were ripened separately at 20C growth chamber (Percival Scientific, Inc., 505 Research Drive Perry, IA) for 10 days. Control fruits were immediately ripened at 20C in the same growth chamber as above. The samples were treated with liquid nitrogen in order to prevent oxidation of phenolic compounds and then lyophilized as previously described (Gorinstein *et al.* 2011; Park *et al.* 2012).

### Chemicals

Trolox; phenolic standards; Tris, tris (hydroxymethyl) aminomethane; Folin–Ciocalteu reagent;  $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$ ; were purchased from Sigma Chemical Co., St Louis, MO, 4, 6-Tripyridyl-*s*-triazine (TPTZ) was from Fluka Chemie, Buchs, Switzerland.

### Determination of Phenolic Acids and Flavonoids


Chlorogenic, hydroxybenzoic, caffeic and *p*-coumaric acids, catechin and epicatechin (mg/100g FW) were determined in the sample methanol extract which was filtered through Sepak C18 Cartridge with HPLC solvent. 5mM of hexadecyltrimethylammonium bromide and 5 mM  $\text{KH}_2\text{PO}_4$  (pH around 4.6) were added to the sample. The standard and extracted samples were analyzed using HPLC pump system (Waters, Model 510), connected with a UV detector (Waters) set at 365 nm. The Waters Symmetry  $\text{C}_{18}$  column (0.5  $\mu\text{m}$ , 4.6x250 mm) was pre-equilibrated with a mobile phase consisting of methanol: water (5:95, v/v) at a flow rate of 1.5 mL/min on the 25C column temperature (Korsak and Park 2010; Jhalegar *et al.* 2011). Determination of bioactive compounds and total antioxidant capacities: The phenols were extracted with methanol (concentration 20 mg/mL) during 1 h in a cooled ultrasonic bath. Total polyphenols (TP, mg gallic acid equivalents (GAE)/100 g FW) were determined by Folin–Ciocalteu method (Singleton *et al.* 1999) with absorbance measurements at 750 nm using spectrophotometer (Hewlett-Packard, model 8452A, Rockville).

### Antioxidant Assay/Activity

Ferric reducing/antioxidant power (FRAP, mM Fe (II) equivalent)/100 g FW) assay measures the ability of the antioxidants in the investigated samples to reduce ferric-tripyridyltriazine ( $\text{Fe}^{3+}$  TPTZ) to a ferrous form ( $\text{Fe}^{2+}$ ). FRAP reagent (2.5 mL of a 10 mmol ferric-tripyridyltriazine solution in 40 mmol HCl plus 2.5 mL of 20 mmol  $\text{FeCl}_3 \cdot \text{xH}_2\text{O}$  and 25 mL of 0.3 mol/L acetate buffer, pH 3.6) of 900  $\mu\text{L}$  was mixed with 90  $\mu\text{L}$  of distilled water and 30  $\mu\text{L}$  of kiwi fruit extract samples as the appropriate reagent blank. The absorbance was measured at 595 nm (Benzie and Strain 1996).

### Fluorometric Measurements

Two dimensional (2D–FL) fluorescence measurements for kiwi fruit extracts at a concentration of 0.01 mg/mL were recorded on a model FP-6500, Jasco spectrofluorometer, serial N261332, Japan, equipped with 1.0 cm quartz cells




**FIG. 1.** EFFECTS OF ETHYLENE TREATMENT (CONTROL VERSUS TREATED), GROWING CONDITIONS (CONVENTIONAL, LOW-CHEMICAL, ORGANIC) AND STORAGE AT 20°C ON THE CONTENT OF SELECTED PHENOLIC ACIDS (mg/100 g FW): (A) HYDROXYBENZOIC ACID; (B) CHLOROGENIC ACID; (C) CAFFEIC ACID; (D) P-COUMARIC ACID OF "HAYWARD" KIWI FRUIT. Depicted values represent averages of 10 measurements and their standard deviations.

and a thermostat bath. The 2D-FL was taken at emission wavelengths from 300 to 750 nm; and at excitation of 280 nm. The three-dimensional spectra (3D-FL) were collected with subsequent scanning emission spectra from 200 to 795 nm at 1.0 nm increments by varying the excitation wavelength from 200 to 500 nm at 10 nm increments (Gorinstein *et al.* 2011; Park *et al.* 2015).

### Statistical Analysis

The data were processed by statistical analysis in order to assess the effects of production conditions as well as the effects of ethylene treatment and long-term storage on the content of selected phenolic acids, flavonoids and polyphenols on "Hayward" kiwi fruit extracts. Two-way ANOVA




**FIG. 2.** EFFECTS OF ETHYLENE TREATMENT (CONTROL VERSUS TREATED), GROWING CONDITIONS (CONVENTIONAL, LOW-CHEMICAL, ORGANIC) AND STORAGE AT 20C ON THE CONTENT OF SELECTED FLAVONOIDS CATECHINS AND EPICATECHINS (mg/100gFW) IN "HAYWARD" KIWI FRUIT. Numerical values represent the highest averaged value of individual parameters in respective time of storage. The depicted data are the averages of 10 measurements and their standard deviations.

(Friedman test) was used to consider the significance of the differences resulting from the above-mentioned treatments. The differences were recognized as significant at the level of significance of  $P \leq 0.05$ , and the differences in means of individual compared characteristics were recognized as highly significant at  $P < 0.001$ . Multivariate statistics was used to reduce the dimensionality of the space of experimental characteristics and to visualize the effects of treatments on the properties of kiwi fruit samples. From multidimensional pattern recognition techniques, the principal component analysis (PCA) and canonical discriminant analysis (CDA) were used to define, interpret and visualize the significance of individual ways of treatment on kiwi fruit antioxidant characteristics. In order to find possible tendencies in the samples and the discriminant power of individual antioxidant characteristics, principal component factoring was effectively applied. All the statistical calculations and data visualizations were performed using the statistical package Unistat v. 6.0 (Unistat, London, United Kingdom). Origin (Microcalc, Northampton, MA) was also used for data visualization in plots, if not specified otherwise.

## RESULTS AND DISCUSSION

### Phenolic Acids and Flavonoids

The fresh "Hayward" kiwi fruit samples cultivated by conventional, low chemical and organic methods, treated with ethylene for 24 h at 20C and then stored were examined for bioactivity quality parameters: hydroxybenzoic, chlorogenic, caffeic and *p*-coumaric acids, catechin, epicatechin, antioxidant activity (FRAP) and total phenolic content (TPC). Phenolic compounds in kiwi fruit pulp were separated and characterized by reversed-phase HPLC. During ethylene treatment and then storage different changes appeared in the bioactivity of the investigated fruits. Strongly acidic compounds were identified as derivatives of *p*-coumaric and caffeic acids (CAs), including chlorogenic acid, protocatechuic acid, and a derivative of 3, 4-dihydroxybenzoic acid. The hydroxybenzoic acid (mg/100g FW) has changed for the organic treated fruits (from  $10.03 \pm 0.10$  to  $12.66 \pm 0.63$ ) in comparison with the conventional (from  $5.77 \pm 0.29$  to  $7.76 \pm 0.39$ , Fig. 1a). The chlorogenic acid (mg/100g FW) has changed for the



**FIG. 3.** EFFECTS OF ETHYLENE TREATMENT (C-CONTROL VERSUS T-TREATED), GROWING CONDITIONS AND STORAGE AT 20°C ON: (A) FRAP VALUES (mM Fe<sup>2+</sup>/100 g FW) AND (B) TOTAL POLYPHENOLS CONTENT (mg GALLIC ACID EQUIVALENTS (GAE)/100 g FW) OF KIWI FRUIT "HAYWARD"


The depicted data are the averages of 10 measurements.

organic treated fruits (from  $36.22 \pm 2.27$  to  $49.11 \pm 2.46$ ) in comparison with the conventional (from  $25.49 \pm 0.84$  to  $35.48 \pm 2.27$ , Fig. 1b). The CA (mg/100g FW) has changed for the organic treated fruits (from  $37.33 \pm 1.87$  to  $47.52 \pm 2.38$ ) in comparison with the conventional (from  $20.46 \pm 1.02$  to  $27.19 \pm 1.36$ , Fig. 1c). The *p*-coumaric acid

(mg/100g FW) has changed for the organic treated fruits (from  $13.51 \pm 10.68$  to  $17.11 \pm 0.86$ ) in comparison with the conventional (from  $8.89 \pm 0.39$  to  $12.60 \pm 0.54$ , Fig. 1d). As it can be compared from the presented data that all organic kiwi fruit samples significantly differ from conventional associates. This is in line with Doverspike *et al.* (2015) about the importance of some phenolic acids which are involved in fruit metabolism. Chlorogenic acid (CGA) is a secondary plant metabolite with chemotherapeutic and chemopreventative potentials along with other health benefits and the amount of this acid is the highest. CGA and CA, a product of the decomposition of CGA, found in skins and flesh of various fruits, and comparison of concentrations in organically and conventionally grown fruits is important. Our results were in agreement with Doverspike *et al.* (2015), but these phenolic acids were found in tomatoes and peaches/nectarines. CA in organic kiwi fruit was 1.75 times higher than in conventional (Fig. 1c), and chlorogenic acid in 1.38 times higher (Fig. 1b). The low chemical fruits were 1.49 and 1.20 times higher than conventional. Our results were agreed with the fact that phenolic compounds were present at levels of <1-7 mg/L in clarified kiwi fruit juice (Dawes and Keene 1999). In spite of many variables studied in this report the results of *bioactivity* of organic kiwi fruit during two seasons were significantly higher than in conventional fruits (Park *et al.* 2015). Some variables are connected with and harvest dates, post-harvest processing, and storage methods and the production quality of completely different systems. Other reports were dealing with the contents of soluble and total phenolic acids in 29 berries and berry products, 24 fruits and fruit peels, and 12 beverages. Among fruits, the highest contents of phenolic acids (28 mg/100 g) were determined in dark plum, cherry, and one apple variety, and lower amount was found in kiwi fruit as it was determined in this study (Mattila *et al.* 2006; Leontowicz *et al.* 2016). The weakly acidic fraction contained epicatechin, catechin, and procyanidins (B3, B2 or B4 and oligomers). The content of catechin (mg/100g FW) has changed for the organic treated fruits (from  $36.88 \pm 0.164$  to  $50.09 \pm 2.50$ ) in comparison with the conventional (from  $26.19 \pm 1.31$  to  $35.9 \pm 1.80$ , Fig. 2a). Epicatechin (mg/100g FW) has changed for the organic treated fruits (from  $34.86 \pm 1.39$  to  $46.96 \pm 2.35$ ) in comparison with the conventional samples (from  $24.66 \pm 1.39$  to  $33.78 \pm 2.05$ , Fig. 2b). Catechin and epicatechin increased as much as 1.40 times higher in organic fruits than in conventional.

### Antioxidant Activity/Phenolic Acids

The final results of total phenolics and the antioxidant activities after all treatments differed for conventional and organic kiwi fruit and were significantly higher in organic




**FIG. 4.** (A) ABSOLUTE DIFFERENTIATION OF "HAYWARD" KIWI FRUIT PERFORMED BY CANONICAL DISCRIMINANT ANALYSIS ACCORDING TO THE GROWING CONDITIONS - CONVENTIONAL (C), LIGHT-CHEMICAL (LC), AND ORGANIC (O). THE ENTIRE DATA SET OF EIGHT CHARACTERISTICS OF ANTIOXIDANT PROPERTIES WAS USED FOR DISCRIMINATION. (B) DIFFERENTIATION OF THE ETHYLENE-TREATED KIWI FRUIT "HAYWARD" PERFORMED BY CANONICAL DISCRIMINANT ANALYSIS OF THE EXPERIMENTAL CHARACTERISTICS ACCORDING TO THE TIME OF STORAGE (0–10 DAYS). THE ENTIRE DATA SET OF EIGHT CHARACTERISTICS OF ANTIOXIDANT PROPERTIES WAS USED FOR DISCRIMINATION.

kiwi fruit than in conventional samples (Fig. 3). The antioxidant activity by FRAP values ( $\text{mM Fe}^{+2}/100 \text{ g FW}$ ) has changed for the organic treated fruits (from  $4.05 \pm 0.15$  to  $4.51 \pm 0.22$ ) in comparison with conventional (from  $2.25 \pm 0.11$  to  $2.56 \pm 0.32$ , Fig. 3a). The total polyphenols ( $\text{mg GAE}/100 \text{ g FW}$ ) have changed for the organic treated fruits (from  $145.256 \pm 4.225.04$  to  $163.11 \pm 2.96$ , Fig. 3b) in comparison with conventional (from  $137.14 \pm 4.22$  to  $154.18 \pm 7.92$ , Fig. 3b). Previously we showed that not always the quality of organic plants was higher than the normally grown (Park *et al.* 2012). Antioxidant values found in this research were different from the ones shown by Park *et al.* (2006). Our results were in line with Wojdylo *et al.* (2013), where was found that Ribes fruit grown in organic system possess significantly higher total phenolics and antioxidant activity (FRAP) than fruit grown in conventional system. The mean value of total polyphenol content in organically grown currants was similar but statistically higher compared with the conventional cultivation. All currants from organic cultivation possess higher ferric reducing capacity than conventionally grown fruits. The results showed that organic fruits tend to have higher hydrolysable polyphenol contents than conventional ones with values being 11.5% in orange peels, 72.6% in papaya peels, and for organic kiwi fruit about 5.5%. The extraction of phenolic

acids in fruits depends on the temperature and the time of extraction. The temperatures of 41–60°C were suitable for major extraction of bioactive compounds and antioxidant capacity from apple pomace. The phenolic compounds were mostly extracted at 60°C in orange peel, as well as the optimization indicated that flavonoids, and antioxidant capacity (DPPH and FRAP assays) were extracted during 0.5 h at 60°C (Hernández-Carranza *et al.* 2016). Polyphenol content and antioxidant capacity varied among organic and conventional vegetables with no prevalence from either agricultural type. In general, organic agriculture results in food products with similar or slightly higher polyphenol content and antioxidant capacity (Faller and Fialho 2010) showing that blueberry fruit grown from organically culture yielded significantly higher total phenolics and antioxidant activity than fruit from the conventional culture. In organically cultured fruit, the average values for antioxidant activity and polyphenols in comparison with the same indices in conventional were 1.49 and 1.68 times higher (Wang *et al.* 2008) and in this study the values were 1.76 and 1.06 times. These results indicate that significant differences between two cultivation practices were evident (Wang *et al.* 2008). Organic kiwi fruits have high levels of total phenol content, resulting in a high antioxidant activity. Their changes were measured after each storage duration at 0, 35, 72, 90 and 120 days of

**FIG. 5. (A)** PRINCIPAL COMPONENT FACTORING ANALYSIS OF “HAYWARD” KIWI FRUIT SAMPLES PERFORMED IN VARIMAX ROTATION ILLUSTRATING THE RELATIONSHIP BETWEEN THE INDIVIDUAL EXPERIMENTAL CHARACTERISTICS, WITHOUT RESPECT ON TREATMENT, GROWING CONDITIONS OR STORING. **(B)** PRINCIPAL COMPONENT ANALYSIS OF “HAYWARD” KIWI SAMPLES PERFORMED ON THE BASIS OF THEIR ANTIOXIDANT CHARACTERISTICS AND CONCENTRATION OF SELECTED ORGANIC ACIDS. EIGENVECTORS ARE VISUALIZED TAKING INTO CONSIDERATION THE GROWING CONDITIONS (C-CONVENTIONAL, LC-LOW-CHEMICAL, O-ORGANIC) Separation of samples according to the ethylene treatment (control group, treated group) is indicated by the dashed line.




storage at 0C and after 1 week of shelf-life simulation at 20C. In a recent report, as it was mentioned previously, “Hayward” kiwi fruits were treated with 100 ppm ethylene at 20C for 24 h and then immediately ripened at 20C for 10 days (Korsak and Park 2010). Comparing antioxidant profile, organically grown fruits showed significantly higher contents of total phenols, flavonoids (catechin and epicatechin) and total antioxidant activities than conventionally grown fruits. Our results (Park *et al.* 2012) were also in line with Anton *et al.* (2014), where was shown that the content of polyphenols was more dependent on year and cultivar than on cultivation conditions. Generally, the cultivation system had minor impact on polyphenols content, and only a few compounds were influenced by the mode of cultivation in all tested cultivars during 2 years. Therefore it is difficult to measure the differences in all variables even during two collection seasons. Recent investigations of nutritional quality in organic versus conventional produce also indicate that soil nitrogen delivery rates strongly affect nutritional quality. Pal *et al.* (2015) showed a strong correlation between polyphenols and antioxidant activity. Positive relationship between total phenol-flavonoids, total phenol-total antioxidant capacity and flavanoids-total antioxidant capacity among organically grown fruits is presented in this

study. Hence, the study concludes that organically grown fruits offer higher amount of nutritional as well as antioxidant properties with inclusion of organically grown fruits in daily diet which could lead to a positive health effects (Roghelia and Patel 2013).

### Processing of Data by Multivariate Statistics

**Canonical Discriminant Analysis.** When the data are processed by CDA separately for the control group of samples and samples treated by ethylene, in both cases absolute differentiation of organically, low-chemically and conventionally grown samples was reached. While for the control group, the dominant role in discrimination had the polyphenols content and FRAP in the first discriminant function and CA and catechin concentrations in the second one. In the ethylene treated group, the dominant role in discrimination played the concentration of hydroxybenzoic acid and FRAP values for the first and hydroxybenzoic and *p*-coumaric acids concentrations for the second discriminant function construction. As expected on the basis of statistical tests performed with respect on ethylene treatment, the discrimination of kiwi fruit samples remain absolute also when this factor has been ignored, as is clearly depicted on Fig. 4a.



**FIG. 6.** SPECTRAL STUDIES OF HUMAN SERUM ALBUMIN (HSA) WITH KIWI FRUIT METHANOL EXTRACTS (A) Emission spectra of HSA in the absence and presence of kiwi fruit methanol polyphenol extracts at  $\lambda_{ex}$  280 nm and  $\lambda_{em}$  300 nm: lines 1–7, HSA in buffer; HSA in methanol; ConvC + HSA; OrgC + HSA; ConvTr + HSA; LChemTr + HSA; OrgTr + HSA. (B, C, D) 3D-spectrum of OrgC + HSA; ConvTr + HSA; ConvC + HSA, Bb, Cc, Dd, cross spectrum of OrgC + HSA; ConvTr + HSA; ConvC + HSA.

Values of the standardized discriminant coefficients, the dominant role in discrimination exhibited the FRAP values/TPC content and the concentration of *p*-coumaric/hydroxybenzoic acids, as regards the first and the second discrimination function construction. It is also apparent from the Fig.

4a, that the properties of conventionally grown samples are closer to that low-chemically grown than those, grown under the organic conditions. This visual tendency is also supported by the calculated distances between the centroids (C-LC = 14.97; C-O = 47.74; LC-O = 34.05). Looking on

**TABLE 1.** FLUORESCENCE DATA OF KIWI FRUIT METHANOL EXTRACTS, HUMAN SERUM ALBUMIN (HSA) AND STANDARD: A, 2D- FLUORESCENCE; B, 3-D FLUORESCENCE

A								
Line number		Samples	$\lambda$ Em	FI	Binding (2D/3D)			
1		HSA*	355	953.22	0			
2		HSA**	353	878.14	7.88			
3		ConvC + HSA	353	825.38	6.01/7.45			
4		OrgC + HSA	352	800.44	8.85/10.81			
5		ConvTr + HSA	353	683.07	22.10/29.15			
6		LChemTr + HSA	347	592.58	32.52/37.08			
7		OrgTr + HSA	355	306.12	65.14/59.76			


B								
Peaks	Indices	Samples						
		OrgC	ConvTr	ConvC	LChemTr	OrgTr	CaffAc	HSA
A	$\lambda_{ex}/\lambda_{em}$	228/347	228/352	228/347	228/350	230/352	228/352	228/347
	FI (A.U.)	436.52	346.08	496.68	301.50	180.21	310.63	502.77
B	$\lambda_{ex}/\lambda_{em}$	278/354	278/352	278/352	280/352	281/355	278/335	280/354
	FI (A.U.)	702.88	559.64	663.01	505.78	342.85	824.01	767.85
C	$\lambda_{ex}/\lambda_{em}$	375/469	375/451	370/444	373/469	376/448	278/650	370/443
	FI (A.U.)	201.18	142.77	62.81	275.47	1000	96.12	68.33
D	$\lambda_{ex}/\lambda_{em}$	275/659	278/657	276/652	278/650	–	–	278/654
	FI(A.U.)	90.04	80.36	90.31	75.18	–	–	102.04
E	$\lambda_{ex}/\lambda_{em}$	229/644	–	–	229/627	–	–	229/640
	FI(A.U.)	63.65	–	–	62.16	–	–	76.24

Abbreviations: line numbers shown in Fig. 1A, Emission mode,  $\lambda_{ex} = 280; \lambda_{em} = 300$ ; \*HSA in buffer; \*\*HSA in methanol; All kiwi fruit methanol extracts and standard were used after interaction with HSA; ConvC, conventional control; OrgC, organic control; ConvTr, conventional treated; LChemTr, low chemical treated; OrgTr, organic treated; CaffAc, caffeic acid;  $\lambda_{ex}/\lambda_{em}$ , (nm/nm); FI, fluorescence intensity, A. U., arbitrary units; binding 2D/3D is determined in %. In 2D the binding properties were found by the decrease of the fluorescence of the main peak (Fig. 6A, Table 1). In 3D, the binding properties were determined by the decrease of fluorescence intensities in peaks **a** and **b** Figs. 6 and 7, Table 1).

these data it can be also concluded, that the kiwi fruit (in terms of their acids content and antioxidant properties) are significantly influenced by the growing conditions. The classification of kiwi fruit samples according to the time of storage brings only partially successful discrimination and classification, reaching thus the average classification score of 50%. It should be mentioned, that all the other factors (growing conditions, ethylene treatment) were ignored. When the variability of the experimental characteristics is decreased (discrimination is performed separately for the control group of samples nontreated by ethylene, and for the ethylene treated samples), then the results of discrimination are much more promising. In both cases, samples were classified into separate groups according to the time of storage with 100% correctness, as is illustrated on Fig. 4a. However, as the signs of the discrimination coefficients in individual functions are opposite, when discriminated all-together, the discrimination score decreases as indicated above. From the above-presented results, it is apparent that ethylene treatment influences the properties of kiwi fruit samples, as well.

**Principal Component Factoring.** All the factors lay in the same sector of the graph practically in one cluster, indicating the potential correlation of individual characteristics. Only slight vector differences for FRAP and polyphenols content are obvious (Fig. 5). However, such result can be satisfactory explained by the fact, that all these characteristics can be considered as antioxidants or revealing the antioxidant properties. Thus, they can be partially used for characterization of antioxidant properties of the complex system. Results of factor analysis fully supported also the values of Pearson's correlation coefficients. For majority of cases, very high correlations ( $> 0.9$ ) were calculated; with exception for just FRAP and polyphenols content. Last two mentioned are characterized by still high but slightly lower correlation coefficients ( $\approx 0.8$ ).

**Principal Component Analysis.** PCA explains the variability of the data set of experimental characteristics and is used to reduce the number of potentially correlating variables via their transformation to noncorrelating group of principal components. In case of PCA of eight experimental



**FIG. 7.** SPECTRAL STUDIES OF HUMAN SERUM ALBUMIN (HSA) WITH KIWI FRUIT METHANOL EXTRACTS

**(E, F, G, H)** 3D-spectrum of LChemTr + HSA; OrgTr + HSA; Caffeic acid + HSA; HSA in methanol. **Ee, Ff, Gg, Hh**, cross spectrum of LChemTr + HSA; OrgTr + HSA; Caffeic acid + HSA; HSA in methanol. Abbreviations for Figs. 6 and 7: All kiwi fruit methanol extracts and standard were used after interaction with HSA; ConvC, conventional control; OrgC, organic control; ConvTr, conventional treated; LChemTr, low chemical treated; OrgTr, organic treated; CaffAc, caffeic acid;  $\lambda_{ex}/\lambda_{em}$ , (nm/nm); FI, fluorescence intensity, A. U., arbitrary units. at  $\lambda_{ex}$  200–500 nm and  $\lambda_{em}$  200–795 nm. The reaction was 1 h at 25C; Concentration of ConvC = 5.6  $\mu\text{g}/\text{mL}$ ; OrgC = 5.7  $\mu\text{g}/\text{mL}$ ; ConvTr = 9.35  $\mu\text{g}/\text{mL}$ ; LChemTr = 10.26  $\mu\text{g}/\text{mL}$ ; OrgTr = 17.16  $\mu\text{g}/\text{mL}$ ; Caffeic acid = 0.05 mM; HSA =  $2 \text{ M} \times 10^{-5}$

characteristics related to antioxidant properties of kiwi fruit, it can be concluded that the PCA explains the variability very effectively – even the 1st principal component is capable of explanation of more than 92% of the whole system variability, while the 2nd component itself explains 5.5% of the remaining variability, reaching thus very promising almost 98% explana-

tion of the whole system variability. 1st PC is constructed equally from all the experimental characteristics, while in the 2nd PC, the dominant role of FRAP value was identified on the basis of eigenvalues comparison. Plot of principal components clearly differentiated the eigenvectors according to the samples growing conditions (conventional - low-chemical -

organic) but also the differentiation according to the ethylene treatment (control versus treated group) can be performed, as indicated by the dashed line in the Fig. 5b. All eight characteristics were utilized for PCs construction. Scores (in %) indicate the contribution of individual components to the explanation of the properties (variability) of the data set of experimental characteristics and the dominant parameter in PC.

### Fluorescence Measurements

The changes in the kiwis' bioactivities during different treatments were interesting to investigate by fluorescence. The 2D fluorescence spectra showed the results described below. At emission of the wavelengths from 347 to 355 nm the recording of the peaks in the fluorescence spectra for conventional, low chemical and organic ethylene treated kiwi fruit after interaction with HSA were with fluorescence intensities (FIs) 683.07, 592.58 and 306.12 arbitrary units (Fig. 6A, lines 5–7 from the top) in comparison with nontreated of 815.38 and 814.44 (Table 1, Fig. 6A, lines 3, 4 from the top). The changes in the fluorescence intensity of treated and untreated samples corresponded with the changes of polyphenol compounds (Table 1). The decrease in the intensity of the main peak during binding of polyphenols to HSA was the smallest for organic treated kiwi fruit (Table 1, Fig. 7F, 7Ff) in comparison with conventional control (Table 1, Fig. 6D, 6Dd). Conventionally, semi-organically (low chemical) and organically grown kiwi fruit was stored at 20C for 10 days. Fluorescence measurements showed a difference between the investigated samples, especially in their antioxidant status. The quenching properties of these samples are correlated with their antioxidant activities and the amounts of polyphenols. The decrease in fluorescence intensity (Park *et al.* 2015) was proportional to the binding properties of extracted polyphenols.

### CONCLUSIONS

From all the statistical analyses, it can be stated here that the antioxidant properties and selected phenolic acids contents of kiwi fruit significantly influenced by both ethylene treatment and the growing conditions. The effect of long term storage is apparent, as well, however, it can only be evaluated in dependence on ethylene treatment. The binding properties of organically grown kiwi fruit were higher than in conventional.

### ACKNOWLEDGMENTS

This research was partly supported by the Rural Development Administration (RDA), Korea. The authors are thankful to Dr. Elena Katrich (School of Pharmacy, Hebrew

University of Jerusalem) for her technical assistance in determination of antioxidant activity and 3 D fluorescence.

### REFERENCES

- AMODIO, M.L., COLELLI, G., HASEY, J.K. and KADER, A.A. 2007. A comparative study of composition and postharvest performance of organically and conventionally grown kiwifruits. *J. Sci. Food Agric.* 87, 1228–1236.
- ANTON, D., MATT, D., PEDASTSAAR, P., BENDER, I., KAZIMIERCZAK, R., ROASTO, M., KAART, T., LUIK, A. and PUSSA, T. 2014. Comparative study of polyphenol contents and antioxidant capacities in fruits of tomato (*Lycopersicon esculentum* Mill.) cultivars grown under organic and conventional conditions. *J. Agric. Food Chem.* 62, 5173–5180.
- BENZIE, I.F.F. and STRAIN, J.J. 1996. The ferric reducing ability of plasma (FRAP) as a measure of antioxidant power: The FRAP assay. *Anal. Biochem.* 239, 70–76.
- DAWES, H.M. and KEENE, J.B. 1999. Phenolic composition of kiwifruit juice. *J. Agric. Food Chem.* 47, 2398–2403.
- DOVERSPIKE, J.C., RADFORD, C.A. and OWENS, J.E. 2015. Determination of chlorogenic acid and caffeic acid in fruits with evaluation of pesticide concentrations. Abstracts of Papers, 249th ACS National Meeting & Exposition, Denver, CO, United States, March 22–26, 2015, CHED-248.
- FALLER, A.L.K. and FIALHO, E. 2010. Polyphenol content and antioxidant capacity in organic and conventional plant foods. *J. Food Compos. Anal.* 23, 561–568.
- GAMMON, C.S., KRUGER, R., MINIHAINE, A.M., CONLON, C.A., VON HURST, P.R. and STONEHOUSE, W. 2012. Kiwifruit consumption favourably affects plasma lipids in a randomised controlled trial in hypercholesterolaemic men. *Br. J. Nutr.* 14, 1–11.
- GANG, C., LI, J., CHEN, Y., WANG, Y., LI, H., PAN, B. and ODEH, I. 2015. Synergistic effect of chemical treatments on storage quality and chilling injury of honey peaches. *J. Food Process. Preserv.* 39, 1108–1117.
- GORINSTEIN, S., POOVARODOM, S., LEONTOWICZ, H., LEONTOWICZ, M., NAMIESNIK, J., VEARASILP, S., HARUENKIT, R., RUAMSUKE, P., KATRICH, E. and TASHMA, Z. 2011. Antioxidant properties and bioactive constituents of some rare exotic Thai fruits and comparison with conventional fruits: In vitro and in vivo studies. *Food Res. Intern.* 44, 2222–2232.
- HERNÁNDEZ-CARRANZA, P., ÁVILA-SOSA, R., GUERRERO-BELTRÁN, J.A., NAVARRO-CRUZ, A.R., CORONA-JIMÉNEZ, E. and OCHOA-VELASCO, C.E. 2016. Optimization of antioxidant compounds extraction from fruit by-products: Apple pomace, orange and banana peel. *J. Food Process. Preserv.* 40, 103–115.
- JHALEGAR, M.J., SHARMA, R.R., PAL, R.K., ARORA, A. and DAHUJA, A. 2011. Analysis of physiological and biochemical changes in kiwifruit (*Actinidia deliciosa* cv. Allison) after the postharvest treatment with 1-methylcyclopropene. *J. Plant Biochem. Biotechnol.* 20, 205–210.

- KORSACK, T. and PARK, Y.S. 2010. Ethylene metabolism and bioactive compounds in ethylene-treated 'Hayward' kiwifruit during ripening. *Hort. Environ. Biotechnol.* *51*, 89–94.
- LEE, D.E., SHIN, B.J., HUR, H.J., KIM, J.H., KIM, J., KANG, N.J., KIM, D.O., LEE, C.Y., LEE, K.W. and LEE, H.J. 2010. Quercetin, the active phenolic component in kiwifruit, prevents hydrogen peroxide-induced inhibition of gap-junction intercellular communication. *Br. J. Nutr.* *104*, 164–170.
- LEONTOWICZ, H., LEONTOWICZ, M., LATOCHA, P., JESION, I., PARK, Y.S., KATRICH, E., BARASCH, D., NEMIROVSKI, A. and GORINSTEIN, S. 2016. Bioactivity and nutritional properties of hardy kiwi fruit *Actinidia arguta* in comparison with *Actinidia deliciosa* 'Hayward' and *Actinidia eriantha* 'Bidan'. *Food Chem.* *196*, 281–291.
- MATTILA, P., HELLSTRÖM, J. and TÖRRÖNEN, R. 2006. Phenolic acids in berries, fruits, and beverages. *J. Agric. Food Chem.* *54*, 7193–7199.
- MIGLIORI, C., DI CESARE, L.F., LO SCALZO, R., CAMPANELLI, G. and FERRARI, V. 2012. Effects of organic farming and genotype on alimentary and nutraceutical parameters in tomato fruits. *J. Sci. Food Agric.* *92*, 2833–2839.
- PAL, R.S., KUMAR, V.A., ARORA, S., SHARMA, A.K., KUMAR, V. and AGRAWAL, S. 2015. Physicochemical and antioxidant properties of kiwifruit as a function of cultivar and fruit harvested month. *Braz. Arch. Biol. Technol.* *58*, 262–271.
- PATARO, G., SINIK, M., CAPITOLI, M.M., DONSI, G. and FERRARI, G. 2015. The influence of post-harvest UV-C and pulsed light treatments on quality and antioxidant properties of tomato fruits during storage. *Innov. Food Sci. Emerg. Technol.* *30*, 103–111.
- PARK, Y.S., JUNG, S.T., KANG, S.G., DELGADO-LICON, E., ARANCIBIA AVILA, P., KATRICH, E., TASHMA, Z., TRAKHTENBERG, S. and GORINSTEIN, S. 2006. Effect of ethylene treatment on kiwifruit bioactivity. *Plant Foods Hum. Nutr.* *61*, 151–156.
- PARK, Y.S., HAM, K.S., KANG, S.G., PARK, Y.K., NAMIESNIK, J., LEONTOWICZ, H., LEONTOWICZ, M., EZRA, A., TRAKHTENBERG, S. and GORINSTEIN, S. 2012. Organic and conventional kiwifruit, myths versus reality: Antioxidant, antiproliferative, and health effects. *J. Agric. Food Chem.* *60*, 6984–6993.
- PARK, Y.S., POLOVKA, M., SUHAJ, M., HAM, K.-S., KANG, S.-G., PARK, Y.-K., ARANCIBIA-AVILA, P., TOLEDO, F., ROBLES SANCHEZ, M. and GORINSTEIN, S., 2015. The postharvest performance of kiwi fruit after long cold storage. *Eur. Food Res. Technol.* *241*, 601–613.
- ROGHELIA, V. and PATEL, V.H. 2013. Nutritional profile and antioxidant potential of selected organically and conventionally grown fruits. *Res. Rev. J. Food Sci. Technol.* *2*, 33–41.
- SINGLETON, V.L., ORTHOFER, R. and LAMUELA-RAVENTOS, R.M. 1999. Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. *Methods Enzymol.* *299*, 152–158.
- VIEIRA, M.J., ARGENTA, L.C., DO AMARANTE, C.V.T., STEFFENS, C.A. and VIEIRA, A.M.F.D. 2010. Postharvest quality preservation of 'Bruno' kiwifruit by ethylene control. *Rev. Bras. Fruticult.* *32*, 397–406.
- WANG, S.Y., CHEN, C.-T., SCIARAPPA, W., WANG, C.Y. and CAMP, M.J. 2008. Fruit quality, antioxidant capacity, and flavonoid content of organically and conventionally grown blueberries. *J. Agric. Food Chem.* *56*, 5788–5794.
- WOJDYLO, A., OSZMIANSKI, J., MILCZAREK, M. and WIETRZ, J. 2013. Phenolic profile, antioxidant and antiproliferative activity of black and red currants (*Ribes* spp.) from organic and conventional cultivation. *Intern. J. Food Sci. Technol.* *48*, 715–726.