

Effects of high temperature on nodulation and nitrogen fixation by *Phaseolus vulgaris* L.

MARIANGELA HUNGRIA and AVÍLIO ANTÔNIO FRANCO
EMBRAPA-CNPSo, Caixa Postal 1061, CEP 86001-970, Londrina, PR, Brazil and
EMBRAPA-CNPBS, Caixa Postal 74505, CEP 23851, Seropédica, RJ, Brazil

Received 2 July 1991. Accepted in revised form 23 November 1992

Key words: common bean, nitrogenase, nodule senescence, *Rhizobium*

Abstract

Screening of *Rhizobium leguminosarum* bv. *phaseoli* strains showed some that were able to nodulate common beans (*Phaseolus vulgaris* L.) at high temperatures (35 and 38°C/8 h/day). The nodulation ability was not related to the capability to grow or produce melanin-like pigment in culture media at high temperatures. However, nodules formed at high temperatures were ineffective and plants did not accumulate N in shoots. Two thermal shocks of 40°C/8 h/day at flowering time drastically decreased nitrogenase activity and nodule relative efficiency of plants otherwise grown at 28°C. Recovery of nitrogenase activity began only after seven days, when new nodules formed; total incorporation of N in tops did not recover for 2 weeks. Non-inoculated beans receiving mineral N were not affected by the thermal shock, and when growing continuously at 35 or 38°C had total N accumulated in shoots reduced by only 18%.

Introduction

Symbiotic N₂ fixation by legumes is strongly affected by temperature conditions and nodulated legumes grow optimally within narrower temperature ranges than those receiving mineral N (Lie, 1981).

High temperature is a major factor limiting N₂ fixation in tropical, and even some subtropical regions. It can depress root hair formation, reducing the sites for nodulation (Frings, 1976; Jones and Tisdale, 1921) and affect adherence of bacteria to hairs (Frings, 1976). The infection process seems to be particularly sensitive (Barrios et al., 1963; Frings, 1976; Lie, 1981). High temperature also affects nodule development (Dart and Mercer, 1965; Jones and Tisdale, 1921; Piha and Munns, 1987) and functioning (Hernandez-Armenta et al., 1989; Meyer and Anderson, 1959; Munévar and Wollum, 1981a;

Piha and Munns, 1987) and may accelerate nodule senescence (Pankhurst and Gibson, 1973; Sutton, 1983).

Legume species vary in their temperature tolerance, and common bean (*Phaseolus vulgaris* L.) is thought to be more sensitive to high temperature than cowpea and soybean (Piha and Munns, 1987). Although the upper limits for N₂ fixation with bean were considered to be between 30°C (Graham and Halliday, 1977; Pankhurst and Sprent, 1976) and 33°C (Barrios et al., 1963), declines in N₂ fixation have been observed even above 28°C (Piha and Munns, 1987). Soil temperatures in tropics often exceed 40°C (Dennet, 1984).

The development of a methodology to allow the selection of temperature-tolerant *Rhizobium* strains under laboratory conditions could contribute greatly to the success of a nitrogen fixing symbiosis in high temperature conditions. Based

on the reports showing that the information for nodulation and for a melanin-like pigment (Beynon et al., 1980) were harbored in the same plasmid of *R. leguminosarum* bv. *phaseoli* (Borthakur et al., 1987; Lamb et al., 1982), Oliveira et al. (1983) isolated eighteen strains that were able to produce dark pigment in culture medium under fluorescent light. In an attempt to use pigment production as an indicator in their selection program, Oliveira et al. (1983) submitted these strains to high temperature, which caused loss of pigment production or capacity to nodulate, but not simultaneously, suggesting that either the genes for those characters are not on the same plasmid, or that they are on the same plasmid but are subject to different temperature control.

In the experiments reported in this paper, we describe the attempts to detect N_2 fixation in beans with some thermal-tolerant strains (both melanin producers and unpigmented) identified by Oliveira et al. (1983). We also examine the effects of a thermal shock on the function of nodules containing a relatively temperature-tolerant strain.

Materials and methods

Two experiments were performed under greenhouse conditions at EMBRAPA-CNPBS, km 47, Rio de Janeiro, Brazil.

In the first, seven strains of *R. leguminosarum* bv. *phaseoli* that were able to grow in plates exposed continuously to high temperature (38°C) were tested for nodulation with common bean (*Phaseolus vulgaris* L. cv. Negro Argel) at 28, 35 and 38°C/8 h/day. Seed N content was of 7.5 mg N/seed. Four of the strains produced melanin-like pigment when grown in TY medium (Beringer, 1974) under the conditions established by Beynon et al. (1980).

For plant inoculation, strains were grown in YM medium (Vincent, 1970) for five days at 28°C. Bean seeds were surface sterilized (Vincent, 1970) and soaked for 1 h in 1-mL of inoculum (about 10^8 cells mL⁻¹) for each fifteen seeds. Five seeds were planted per pot, and covered with a layer of sterile sand. Pots were made of PVC tubes filled with two thick plastic

bags, and received 2.5 kg of sterile sand and vermiculite (1/2, v/v). Four days after emergence (DAE) plants were thinned to two per pot. At the time of sowing, pots were put into water tanks so that the water level was just above that of the substrate inside the pot. Thermostats controlling the water temperature at 28, 35 or 38°C were turned on from 7:00 a.m. to 5:00 p.m., taking about 2 h to reach the desired maxima. A pump circulated the water in the tank during the hot period. After the heating was turned off, the temperature of the pots quickly dropped to 23°C. Temperatures around the shoots averaged 29/23°C, day/night.

Pots were watered through a capillary 'candle filter' system in each pot to maintain the pots at the moisture of about 75% of maximum water holding capacity (Boddey et al., 1987).

Nutrient solution without mineral N was provided weekly and contained: 0.725 mM NaH₂PO₄; 0.725 mM KH₂PO₄; 3.1 mM MgSO₄ 7H₂O; 2.9 mM CaSO₄ 2H₂O; 1.5 mM K₂SO₄; 1 μM MnSO₄ 4H₂O; 0.1 μM CuSO₄ 5H₂O; 0.1 μM ZnSO₄ 7H₂O; 5.0 μM H₃BO₃; 10.0 μM NaCl; 0.5 μM NaMoO₄ 2H₂O; 0.02 μM CoSO₄ 6H₂O and 5.0 μM Fe-EDTA; pH 6.0 to 6.2. The treatment without inoculation received a total of 8 mg of N as KNO₃/plant/day from 5 to 26 DAE.

The experiment was laid out in a split-plot design, each temperature replicated in two tanks with two pots in each tank. Plants were harvested at 28 DAE (early flowering) and the following parameters were evaluated: shoot dry weight and total N (Boddey et al., 1987), nodule number and dry weight.

In the second experiment, of similar experimental design, cv. Negro Argel was inoculated with strain SEMIA 487 (IPAGRO, RS, Brazil), or received mineral N (10 mg N as KNO₃/plant/day, starting at 5 DAE). Inoculation and planting were as described for the first experiment. Plants were grown in the water tanks at 28°C during the day and 23°C at night. At 30 DAE (early flowering), a thermal shock of 40°C for 8 h was applied to half of the pots, and the shock was repeated on day 31. Plants were harvested at 30, 31, 32, 33, 36, 38 and 47 DAE.

Acetylene reduction activity was evaluated on excised roots using a continuous flow system with

12% of acetylene in air at a rate of 100 mL/min, as described before (Boddey et al., 1987). Previously (Hungria and Neves, 1987a), a comparative experiment was performed to evaluate whether the nitrogenase was affected by the presence of acetylene (Minchin et al., 1983). Plants were transferred to a continuous flow system, detopped and flushed with an air stream containing 12% of C₂H₂. Samples were taken after 1, 3, 4, 5, 6, 7, 10, 15 and 30 minutes. Percent decreases in nitrogenase activity for thirty symbiotic combinations (five bean cultivars and six *Rhizobium* strains) were just 4.5% after 30 minutes. In the experiments reported in this paper, we detopped the plants, immediately incubated with acetylene and samples were taken after 4 minutes with acetylene. Although we have not investigated the effect of detopping, it is unlikely that any significant effect would take place in this short interval.

Hydrogen evolution by the nodules was measured on excised root systems that were incubated in air in 280-mL jars under normal atmosphere for 10 minutes, and hydrogen was analyzed by gas chromatography as described previously (Boddey et al. 1987). Relative efficiency of nitrogenase was determined according to Schubert and Evans (1976).

Nitrate reductase activity of leaves, stems and

roots was evaluated according to Hungria and Neves (1987b). Nodules were examined for texture (hard/soft) and internal color (pink/brown) and classified as senesced (soft and/or brown) or non-senesced (hard and pink). Number and dry weight of nodules were also evaluated. Plant dry weight and total N (shoots + roots) were analyzed according to Boddey et al. (1987).

Results

All seven *R. leguminosarum* bv. *phaseoli* strains were able to nodulate beans when root systems were exposed at 35 and 38°C/8 h/day. This ability was not related to their growth or melanin-like pigment production in plates exposed continuously to 38°C (Table 1). However, nitrogenase activity was not detected and plants had a very low N content. On the other hand, plants receiving mineral N were not greatly affected, and total N accumulated in shoots decreased by only 18% at 35 or 38°C (Table 1).

A thermal shock of 40°C/8 h/day at 30 DAE, of plants previously grown at 28°C, resulted in a 70% loss of nitrogenase activity 24 h after, and with a second thermal shock at 31 DAE, there was a further loss in activity of 15% (Fig. 1). The proportion of electrons diverted to hydrogen

Table 1. Nodulation and N accumulation in shoots of common bean, cv. Negro Argel, inoculated with seven *Rhizobium* strains or receiving mineral N (8 mg N/plant/day from 5 to 26 DAE) and grown at 28, 35 or 38°C/8 h/day and 23°C at night. Plants were harvest at early flowering (28 DAE) and data represent the average of four replicates

<i>Rhizobium</i> strains	Production of dark pigment in TY medium at 28°C	Growth in YM medium		Nodulation						Shoot N (mg N/plant)		
		35	38	Number			Dry weight			28	35	38
				28	35	38	28	35	38			
SEMIA 487 ^a	+	+	+	95	63	39	50	40	23	52	18	18
SEMIA 4021 ^a	+	+	+	79	77	13	50	23	10	50	20	24
F 413 ^b	+	+	+	84	70	32	40	43	33	54	24	20
F 413 Mn3 ^c	+	+	–	82	61	13	40	27	17	40	18	20
C05 ^d	–	–	–	59	33	13	47	27	20	46	26	20
CIAT 57 ^e	–	–	–	115	79	15	73	37	17	54	18	18
CIAT 255 ^e	–	–	–	96	87	13	84	30	13	58	18	20
Mineral N				0	0	0	0	0	0	92	76	74
Analyses of variance												
Temperature ($p < 0.05$)							17		12			9
Inoculation ($p < 0.05$)							20		16			8

^aIPAGRO, RS, Brazil; ^b Nitragin Co., USA; ^cF 413 isolated from soils with high levels of manganese; ^dCENA, SP, Brazil; ^eCIAT, Cali, Colombia.




Fig. 1. Effects of two thermal shocks (40°C/8 h/day) at 30 and 31 DAE on the nitrogenase activity of bean, cv. Negro Argel, inoculated with *Rhizobium* strain SEMIA 487 and grown at room temperature (28°C). Means of four replicates and vertical bars denote LSD for each harvest ($p = 0.05$).




Fig. 2. Effects of two thermal shocks (40°C/8 h/day) at 30 and 31 DAE on the relative efficiency of nitrogenase of bean cv. Negro Argel, inoculated with SEMIA 487 and grown at room temperature (28°C). Means of four replicates, and vertical bars denote LSD for each harvest ($p = 0.05$).

evolution was also changed, and the relative efficiency decreased with the increase in the temperature (Fig. 2). By 33 DAE, both nitrogenase (Fig. 1) and relative efficiency (Fig. 2) had begun to recover, but 7 days after the second thermal shock, the nitrogenase activity was still less than half of the control level. However, total nitrate reductase (nitrate reductase of leaves + stems + roots) was not affected by the thermal shocks, declining only at 36 DAE for both treatments (Fig. 3).

Five days after the thermal shock, the number of senesced nodules increased by 33%, but 7 days after the plants had been submitted to 40°C new nodules could be seen, increasing nodule number per plant and decreasing significantly the percentage of senesced nodules. So, at 47 DAE, only 17 days after the first thermal shock, the number of nodules in stressed plants was 54% higher than that of the control (Fig. 4). New nodules led to recovery of nitrogenase activity and relative efficiency, with values significantly higher in stressed plants at 47 DAE (Figs. 1 and 2).

There was no effect of high temperature shock

on N accumulation by the plants that received mineral N (Fig. 5). Also at the final harvest there was no difference between plants given mineral N and those inoculated with strain SEMIA 487 and growing at optimal temperatures.

Discussion

Other workers have also shown that the ability of rhizobia and bradyrhizobia to grow in culture medium at high temperatures (up to 47°C) does not necessarily relate to the infectivity and N₂ fixation under temperature-stress conditions (Karanja and Wood, 1988; La Favre and Eaglesham, 1986). Also the attempts to use a genetic marker, such as the production of melanin-like pigment, to identify temperature-tolerant rhizobia have not been successful (Oliveira et al., 1983). The role of melanin-like pigment production by *R. leguminosarum* strains is still unclear. We found no relation between the production of this pigment in plates and the nodulation ability of *R. leguminosarum* by.




Fig. 3. Effects of two thermal shocks (40°C/8 h/day) at 30 and 31 DAE on nitrate reductase activity in leaves of beans, cv. Negro Argel, receiving 10 mg N/plant/day and grown at room temperature (28°C). Means of 4 replicates and vertical bars denote LSD for the effect of harvesting time ($p = 0.05$).




Fig. 4. Effects of two thermal shocks (40°C/8 h/day) at 30 and 31 DAE on nodulation of beans cv. Negro Argel inoculated with SEMIA 487 and grown at 28°C. Mean of 4 replicates and vertical bars denote LSD for each harvest ($p = 0.05$).

phaseoli strains. Also in *R. leguminosarum* bv. *viceae*, Hynes et al. (1988) showed that the loss of dark-pigment production had no effect on the




Fig. 5. Nitrogen accumulation by bean cv. Negro Argel, inoculated with strain SEMIA 487 or receiving 10 mg N/pl/day. Plants were grown at room temperature (28°C) and received or not thermal shocks of 40°C/8 h/day at 30 and 31 DAE. Means of 4 replicates and vertical bars denote LSD for each harvest ($p = 0.05$)

symbiotic performance or competitive ability of this strain.

The effect of a thermal shock is unlikely to reflect simply the loss of Sym plasmid, since Soberón-Chávez et al. (1986) showed that strains of *R. leguminosarum* bv. *phaseoli* lost symbiotic properties at a frequency of 62% after growth at 37°C for five days, but maintained the same electrophoretic pattern and the same plasmid molecular weight. In that work, the loss of Fix phenotype resulted from genomic rearrangements that could only be shown by hybridizing techniques (Soberón-Chávez et al., 1986).

Although in the first experiment seven *Rhizobium* strains were capable of nodulating beans when root day temperatures were 35 and 38°C, data on the acetylene reduction activity, total N accumulated by plants (Table 1) and number of infected cells (data not shown) showed that these nodules were not functional.

Similar results were also obtained for beans by Gitonga et al. (1989). There was also no relation between the origin of the strains (from tropical or temperate regions) and the resistance to high temperatures, as also observed by Munévar and Wollum (1981a).

Differences among strains in tolerance to high temperature were observed for *Bradyrhizobium japonicum* (Dart et al., 1976; La Favre and Eaglesham, 1986; Munévar and Wollum, 1981a,b) and *R. leguminosarum* bv. *phaseoli* strains (Karanja and Wood, 1988; Piha and Munns, 1987), but as in our experiments, there was almost no N₂ fixation with temperatures above 33°C. Since plants receiving mineral N in experiment 1 had total N decreased by only 18%, the limitation appears to be in the symbiosis rather than in the growth potential of the bean plant.

In the second experiment, plants grown under optimal temperature conditions and exposed to a thermal shock at flowering time decreased nitrogenase activity by 70% after 24 h (Fig. 1). Similar results were obtained by Hernandez-Armenta et al. (1989) confirming that the symbiosis of *Phaseolus vulgaris*-*R. leguminosarum* bv. *phaseoli* is extremely sensitive to high temperatures. The proportion of electrons allocated to the proton reduction was increased after 38°C (Fig. 2), as it was also observed before for nodulated cowpea (Rainbird et al., 1983). In our experiment, plants started to recover after one week, due to the development and/or formation of new nodules (Fig. 4).

The data presented here demonstrate that high temperature (35°C or above) can affect the symbiosis of *P. vulgaris*-*R. leguminosarum* bv. *phaseoli* and none of the strains tested was able to fix N₂ under these stressed conditions. However, in tropical and even some subtropical regions, soil temperatures between 40–60°C are often observed, and prolonged periods at 38°C in the surface 4 cm of soil were reported in North Carolina, USA (Munévar and Wollum, 1981a). Since we have shown that the plant growth per se was not greatly affected by heat (35 or 38°C/8 h/day during growth or a thermal shock of 40°C/8 h/day at flowering), there would still seem to be potential for improvement of bean yields from inoculation if the appropriate strain

could be found. To this end we have investigated other sources of *Rhizobium* for N₂ fixation with beans at high temperatures, and we describe this selection work in the following paper (Hungria et al., 1993).

Acknowledgements

We thank Dr. Johanna Döbereiner for suggestions during the experiment and Dr. Allan R J Eaglesham for criticism of the paper and manuscript review. And we would like to thank specially the valuable contribution and suggestions of Dr. Janet I Sprent. The work was funded by the Financiadora de Estudos e Projetos-FINEP.

References

- Barrios S, Raggio N and Raggio M 1963 Effect of temperature on infection of isolated bean roots by rhizobia. *Plant Physiol.* 38, 178–174.
- Beringer J E 1974 R factor transfer in *Rhizobium leguminosarum*. *J. Gen. Microbiol.* 84, 188–198.
- Beynon J L, Beringer J E and Johnston A W B 1980 Plasmids and host-range in *Rhizobium leguminosarum* and *Rhizobium phaseoli*. *J. Gen. Microbiol.* 120, 421–429.
- Boddey R M, Pereira J A R, Hungria M, Thomas R J and Neves M C P 1987 Methods for the study of nitrogen assimilation and transport in grain legumes. *MIRCEN J. Appl. Environm. Biotechnol.* 3, 3–22.
- Borthakur D, Lamb J W and Johnston A W B 1987 Identification of two classes of *Rhizobium phaseoli* genes required for melanin synthesis, one of which is required for nitrogen fixation and activates the transcription of the other. *Mol. Gen. Genet.* 207, 155–160.
- Dart P J, Day J, Islam R and Döbereiner J 1976 Symbiosis in tropical grain legumes: some effects of temperature and the composition of the rooting medium. *In Symbiotic Nitrogen Fixation in Plants*. Ed. P S Nutman. pp 361–383. Cambridge University Press, Cambridge.
- Dart P J and Mercer F V 1965 The effect of growth temperature, level of ammonium nitrate, and light intensity on the growth and nodulation of cowpea (*Vigna sinensis* Endl. ex. Hassk.). *Aust. J. Agric. Res.* 16, 321–345.
- Dennet M D 1984 The tropical environment. *In The Physiology of Tropical Field Crops*. Eds. R P Goldsworthy and N M Fisher. pp 1–38. Wiley, New York.
- Frings J F J 1976 The *Rhizobium*-Pea Symbiosis as Affected by High Temperatures. Thesis. Wageningen Agricultural University, Wageningen. 76 p.
- Gitonga N M, Widdowson D and Keya S O 1989 Interaction of *Phaseolus vulgaris* with thermotolerant isolates of

- Rhizobium leguminosarum* biovar *phaseoli* from Kenyan soils. MIRCEN J. 5, 493–504.
- Graham P H and Halliday J 1977 Inoculation and nitrogen fixation in the genus *Phaseolus*. In *Exploiting the Legume Symbiosis in Tropical Agriculture*. Eds. J M Vincent, A S Whitney and J Bose. pp 313–333. College of Tropical Agriculture Publication, 145. University of Hawaii, Hawaii, HI.
- Hernandez-Armenta R, Wien H C and Eaglesham A R J 1989 Carbohydrate partitioning and nodule functioning in common bean after heat stress. *Crop Sci.* 29, 1292–1297.
- Hungria M, Franco A A and Sprent J I 1993 New sources of high-temperature tolerant rhizobia for *Phaseolus vulgaris*. *Plant and Soil* 149, 103–109.
- Hungria M and Neves M C P 1987a Cultivar and *Rhizobium* strain effects on nitrogen fixation and transport in *Phaseolus vulgaris* L. *Plant and Soil* 103, 111–121.
- Hungria M and Neves M C P 1987b Partitioning of nitrogen from biological fixation and fertilizer in *Phaseolus vulgaris* L. *Physiol. Plant.* 69, 55–63.
- Hynes M F, Brucksch K and Priefer U 1988 Melanin production encoded by a cryptic plasmid in a *Rhizobium leguminosarum* strain. *Arch. Microb.* 150, 326–332.
- Jones F R and Tisdale W B 1921 Effect of soil temperature upon the development of nodules on the roots of certain legumes. *J. Agric. Res.* 22, 17–37.
- Karanja N K and Wood M 1988 Selecting *Rhizobium phaseoli* strains for use with beans (*Phaseolus vulgaris* L.) in Kenya: Tolerance of high temperature and antibiotic resistance. *Plant and Soil* 112, 15–22.
- La Favre A K and Eaglesham A R J 1986 The effects of high temperatures on soybean nodulation and growth with different strains of bradyrhizobia. *Can. J. Microbiol.* 32, 22–27.
- Lamb J W, Hombrecher G and Johnston A W B 1982 Plasmid-determined nodulation and nitrogen-fixation abilities in *Rhizobium phaseoli*. *Mol. Gen. Genet.* 186, 449–452.
- Lie T A 1981 Environmental physiology of the legume-*Rhizobium* symbiosis. In *Nitrogen Fixation*. Vol I. Ed. W J Broughton. pp 104–134. Clarendon Press, Oxford.
- Meyer D R and Anderson A J 1959 Temperature and symbiotic nitrogen fixation. *Nature* 183, 161.
- Minchin F R, Witty J F, Sheehy J E and Muller M 1983 A major error in the acetylene reduction assay: Decreases in nodular nitrogenase activity under assay conditions. *J. Exp. Bot.* 34, 641–649.
- Munévar F and Wollum II A G 1981a Growth of *Rhizobium japonicum* strains at temperatures above 27°C. *Appl. Environm. Microb.* 42, 272–276.
- Munévar F and Wollum II A G 1981b Effect of high root temperature and *Rhizobium* strain on nodulation, nitrogen fixation and growth of soybeans. *Soil Sci. Soc. Am. J.* 45, 1113–1120.
- Oliveira L C B de, Franco A A and Döbereiner J 1983 Estabilidade da efetividade e eficiência de estirpes de *Rhizobium phaseoli* submetidas a altas temperaturas. *An. Cong. Bras. Ci. Solo*, 19. Curitiba, SBCS, p. 3.
- Pankhurst C E and Gibson A H 1973 *Rhizobium* strain influence on disruption of clover nodule development at high root temperature. *J. Gen. Microb.* 74, 219–231.
- Pankhurst C E and Sprent J I 1976 Effects of temperature and oxygen tension on the nitrogenase and respiratory activities of turgid and water-stressed soybeans and French bean root nodules. *J. Exp. Bot.* 27, 1–9.
- Piha M I and Munns D N 1987 Sensitivity of the common bean (*Phaseolus vulgaris*) symbiosis to high soil temperature. *Plant and Soil* 98, 183–194.
- Rainbird R M, Atkins C A and Pate J S 1983 Effect of temperature on nitrogenase functioning in cowpea nodules. *Plant Physiol.* 73, 392–394.
- Schubert K R and Evans H J 1976 Hydrogen evolution: A major factor affecting the efficiency of nitrogen fixation in nodulated symbionts. *Proc. Natl. Acad. Sci. USA* 73, 1207–1211.
- Soberón-Chaves G, Nájera R, Oliveira H and Segovia L 1986 Genetic rearrangements of a *Rhizobium phaseoli* symbiotic plasmid. *J. Bacteriol.* 167, 487–491.
- Sutton W D 1983 Nodule development and senescence. In *Nitrogen Fixation*. Vol. 3. Ed. W J Broughton, pp 144–212. Clarendon Press, Oxford.
- Vincent J M 1970 *Manual for the Practical Study of Root Nodule Bacteria*. IBP Handbook nr. 15. Blackwell, Oxford. 164 p.

Section editor: R O D Dixon