

## Diversity of Endophytic Enterobacteria Associated with Different Host Plants

Adalgisa Ribeiro Torres<sup>1</sup>, Wellington Luiz Araújo<sup>1,2\*</sup>, Luciana Cursino<sup>1</sup>, Mariangela Hungria<sup>3</sup>,  
Fábio Plotegher<sup>3</sup>, Fábio Luís Mostasso<sup>3</sup>, and João Lúcio Azevedo<sup>1,2</sup>

<sup>1</sup>Genetics Department, "Luiz de Queiroz" College of Agriculture, University of São Paulo (ESALQ/USP), Piracicaba, SP, Av. Pádua Dias, 11, P. O. Box 83, 13400-970, Brazil

<sup>2</sup>Laboratory of Microbial Genetics, NIB, University of Mogi das Cruzes, SP, Brazil

<sup>3</sup>Brazilian Agricultural Research Corporation – Embrapa Soybean, Londrina, PR, Brazil

(Received September 18, 2007 / Accepted June 24, 2008)

**Fifty-three endophytic enterobacteria isolates from citrus, cocoa, eucalyptus, soybean, and sugar cane were evaluated for susceptibility to the antibiotics ampicillin and kanamycin, and cellulase production. Susceptibility was found on both tested antibiotics. However, in the case of ampicillin susceptibility changed according to the host plant, while all isolates were susceptible to kanamycin. Cellulase production also changed according to host plants. The diversity of these isolates was estimated by employing BOX-PCR genomic fingerprints and 16S rDNA sequencing. In total, twenty-three distinct operational taxonomic units (OTUs) were identified by employing a criterion of 60% fingerprint similarity as a surrogate for an OTU. The 23 OTUs belong to the *Pantoea* and *Enterobacter* genera, while their high diversity could be an indication of paraphyletic classification. Isolates representing nine different OTUs belong to *Pantoea agglomerans*, *P. ananatis*, *P. stewartii*, *Enterobacter* sp., and *E. homaechei*. The results of this study suggest that plant species may select endophytic bacterial genotypes. It has also become apparent that a review of the *Pantoea/Enterobacter* genera may be necessary.**

**Keywords:** endophytes, BOX-PCR, diversity, cellulase production, antibiotic susceptibility

The study of endophytic microorganisms' genetic diversity is not only important for understanding their ecological role in natural environments, but can also their biotechnological application (Wise *et al.*, 1996). The fate and distribution of a species in a natural environment may in part be ruled by genetic diversity within the species. As such, it is necessary to estimate this diversity. Several high-resolution molecular fingerprinting techniques have been used to assess species and subspecies diversity (Vanechoutee *et al.*, 1998; Rademaker *et al.*, 2000; Schloter *et al.*, 2000). Ribotyping, repetitive DNA sequences and PCR (Dombek *et al.*, 2000; Carson *et al.*, 2003) techniques have been successfully applied to cluster bacteria from the *Enterobacteriaceae*, *Rhizobiaceae*, and *Actinobacteridae* families (Terasawa *et al.*, 2003; Davelos *et al.*, 2004; Yang *et al.*, 2004). BOX-PCR is a technique based on primers targeting the highly conserved repetitive DNA sequences of the BOXA subunit in bacterial genomes (Rademaker *et al.*, 2000; Lanoot *et al.*, 2004). Moreover, BOX-PCR is based on the observation that outwardly facing oligonucleotide primers, complementary to interspersed repeated sequences, enable the amplification of different-sized DNA fragments, consisting of sequences lying in between these elements. It has been applied in a number of taxonomic studies on plant-associated bacteria (Rademaker *et al.*, 2000; Berg *et al.*, 2002), while only used in two studies on endophytic bacteria (Verma *et al.*, 2001; Krechel *et al.*,

2002). Interestingly, no studies have analyzed endophytic bacteria from the *Enterobacteriaceae* family.

The aim of the present study was evaluate the genetic and physiological aspects of endophytic enterobacteria from five commercially important Brazilian host plants (citrus, cocoa, eucalyptus, soybean, and sugar cane) based on production of cellulase (endoglucanase), antibiotic susceptibility and molecular methods, such as BOX-PCR and 16S rRNA gene sequencing.

### Materials and Methods

#### Bacterial isolates and culture conditions

The bacterial isolates used in this study were obtained from a collection belonging to the Laboratory of Microbial Genetics, Department of Genetics, ESALQ/USP. The isolates were grown in liquid LB media (Sambrook *et al.*, 1989) at 28°C under agitation (200 rpm).

#### Susceptibility analysis

The susceptibility of 53 isolates to antibiotics was determined in the following manner: bacteria were grown in LB broth until  $5 \times 10^5$  CFU/ml according to the guidelines of the National Committee for Clinical Laboratory Standards (NCCLS, 2004). The culture was then spotted with a 16-pin handheld replicator onto the LB agar supplemented with antibiotics. Selective concentrations used were 50 and 100 µg/ml for ampicillin and kanamycin, respectively.

\* To whom correspondence should be addressed.  
(Tel) 55-19-3429-4251; (Fax) 55-19-3433-6706  
(E-mail) wlaraujo@esalq.usp.br

### Plate cellulase enzyme assays

Plate assays were performed according to the methods of Teather and Wood (1982). Isolated colonies of 53 bacterial cultures were picked and inoculated using a toothpick on LB agar supplemented with 1.5% (w/v) carboxy-methyl-cellulose (CMC). The inoculated plates were incubated for 24 h at 28°C. After incubation, the plates were flooded with Congo red (1 mg/ml) solution, incubated at room temperature for 15 min and washed several times with sterile 1 M NaCl solution for 10–15 min to remove unbounded excess dye. A clearing zone surrounded by a red background was indicative of CMC hydrolysis. The cellulase activity index was achieved by the ratio of CMC degradation clear zone diameter and that of bacterial colony.

### Genomic DNA extraction

Genomic DNA was extracted from endophytic bacteria using the following protocol: a 1.5-ml sample of an overnight bacterial culture was centrifuged for 2 min at 12,000×g and re-suspended in 1 ml of TE buffer (10 mM Tris-HCl, 1 mM EDTA, pH 8.0), re-centrifuged, decanted, and re-suspended in 500 µl of TE buffer plus 0.5 g of 0.1-mm diameter glass beads and 30 µl of 10% sodium dodecyl sulfate. The cells were homogenized for 30 sec in a bead beater (Braun cell homogenizer; B. Braun, Germany). A 500 µl vol of Tris-buffered phenol was then added, the solution was well mixed and centrifuged for 10 min at 12,000×g. The aqueous phase was extracted once with phenol-chloroform (1:1) and once with chloroform, and the DNA was precipitated with isopropanol (5 min at room temperature) and collected by centrifugation (10 min at 12,000×g). The pellet was washed with 70% ethanol, air dried, and re-suspended in 50 µl of TE buffer.

### BOX-PCR analysis

BOX-PCR reactions were performed in a 25-µl volume containing 5–10 ng of isolated DNA, 7.5 mM MgCl<sub>2</sub>, 1×10<sup>6</sup> µM of BOX A1R primer; 5'-CTACGGCAAGGCGACGCTGACG-3' (Koeuth *et al.*, 1995), 1 U of the *Taq* DNA polymerase (Invitrogen, USA), and 0.6 mM each dNTP. DNA amplification was carried out in a thermal cycler (PTC-200, MJ Research, USA) under the following conditions: 95°C for 10 min followed by 30 cycles of 1 min at 95°C, 1 min 30 sec at 72°C. The final extension step was performed at 72°C for 10 min. The PCR products were analyzed in 1.5% agarose gels in Tris-borate-EDTA (TBE) buffer at 100 V. DNA bands were prepared for visualization by staining with ethidium bromide (0.5 µg/ml) for 15 min and washing in water. Subsequently, they were analyzed under UV light and photographed using the Gel Doc 2000 documentation system (Bio-Rad, USA). A 1 kb molecular size ladder (Fermentas, Inc., USA) was included in each agarose run.

### Computer-assisted analysis of BOX-PCR DNA fingerprints

PCR reactions were electrophoresed on 1.5% agarose gel for 2 h at 100 V. Gels were stained with ethidium bromide (15 min). Images were captured and saved directly as TIFF files and processed by BioNumerics 3.0 (Applied Maths, Belgium). Similarity matrices of densitometric curves were

calculated by a Jaccard correlation coefficient. Cluster analyses of similarity matrices were performed by an unweighted pair group method with an arithmetic mean (UPGMA) algorithm. The correlation was expressed as percent similarity. Minimum similarity at a level of 60% between isolates was randomly used as the criterion to define an operational taxonomic unit (OTU) (Yang *et al.*, 2004).

### Bacterial identification by 16S rRNA gene sequencing


The molecular identification of isolates representative of some OTUs was done based on 16S rRNA gene sequencing. For this, the 16S rDNA was amplified using the primers 27F; 5'-AGAGTTTGTATCMTGGCTCAG-3' and 1378R; 5'-CGG TGTGTACAAGGCCCGGGAACG-3' under the following PCR conditions: 4 min at 94°C, 30 cycles of 30 sec at 94°C, 1 min at 62.5°C, and 1 min at 72°C, followed by a final extension step of 1 min at 72°C. The PCR master mix contained 3.75 mM MgCl<sub>2</sub>, 0.2 mM dNTP, 0.2 µM primer, 2.5 U *Taq* DNA polymerase, 1× PCR buffer and 1 µl DNA template (0.5–10 ng). The PCR products were purified using a GFX PCR DNA and gel band purification kit (Amersham Biosciences) and sequenced using the 1378R primer. Analyses of sequences were performed with the basic sequence alignment program, BLAST, which ran against the database [National Center for Biotechnology Information website (<http://www.ncbi.nlm.nih.gov/BLAST>)]. The determined sequence were aligned using CLUSTAL W and the distance matrices and phylogenetic trees were calculated by Jukes and Cantor (1969) and neighbour-joining (Saitou and Nei, 1987) algorithms, respectively, using MEGA4.0 software. The nucleotide sequences obtained in this study have been submitted to the GenBank and assigned accession numbers AY914088, AY914093, AY914097, AY914789, AY914792, AY914795, AY920254, AY920255, AY920524.

### Diversity index estimation

The Shannon diversity index ( $H = -\sum(x_i/x_0)\ln(x_i/x_0)$ ) where  $x_i$ =number of isolates in an OTU and  $x_0$ =total number of isolates) was calculated for each host plant in the BOX-PCR experiments (Margurran, 1988).

### Statistical analysis

Data analysis was performed using the SAS software package version 6.0 (SAS, 1987). For BOX-PCR experiments, a



**Fig. 1.** Cellulase activity index (AI) of bacterial isolates from five host plants. AI for each host plant was averaged.

**Table 1.** Bacterial isolates used, with their relevant characteristics

Isolates	Species	Host plant	OTUs	Ampicillin phenotype	Cellulase production	Reference or source <sup>a</sup>	
ARB18	<i>Pantoea agglomerans</i>	Citrus	10	S	+	Araújo <i>et al.</i> (2002)	
ARB18/10		Citrus	10	R	+	Araújo <i>et al.</i> (2002)	
CL12		Citrus	9	R	-	Araújo <i>et al.</i> (2001)	
CTO2		Citrus	9	R	-	Araújo <i>et al.</i> (2001)	
CTO5		Citrus	9	R	-	Araújo <i>et al.</i> (2001)	
LC32		Citrus	9	R	-	Araújo <i>et al.</i> (2001)	
LCO6		Citrus	9	R	-	Araújo <i>et al.</i> (2001)	
LT37		Citrus	9	R	-	Araújo <i>et al.</i> (2001)	
LV23		Citrus	9	R	-	Araújo <i>et al.</i> (2001)	
LV24		Citrus	9	R	-	Araújo <i>et al.</i> (2001)	
PR1.5/2		Citrus	5	S	+	Araújo <i>et al.</i> (2002)	
PR1.5/4		Citrus	5	R	+	Araújo <i>et al.</i> (2002)	
PR1.5/9		Citrus	3	R	-	Araújo <i>et al.</i> (2002)	
PR1/4		Citrus	15	R	-	Araújo <i>et al.</i> (2002)	
PR2/7	<i>Enterobacter cloacae</i>	Citrus	12	S	-	Araújo <i>et al.</i> (2002)	
TC41		Citrus	9	R	-	Araújo <i>et al.</i> (2001)	
TO45		Citrus	9	R	-	Araújo <i>et al.</i> (2001)	
TO47		Citrus	9	R	-	Araújo <i>et al.</i> (2001)	
TS49		Citrus	9	R	-	Araújo <i>et al.</i> (2001)	
TS51		Citrus	9	R	-	Araújo <i>et al.</i> (2001)	
C32.2		Cocoa	12	R	-	ESALQ/USP <sup>b</sup>	
C33.2		Cocoa	12	R	+	ESALQ/USP	
C35.1		Cocoa	12	R	-	ESALQ/USP	
C39.1		Cocoa	1	R	-	ESALQ/USP	
C40.1		Cocoa	1	R	-	ESALQ/USP	
C41.1		Cocoa	12	R	+	ESALQ/USP	
C45.1		Cocoa	12	R	-	ESALQ/USP	
C49.1		Cocoa	12	R	-	ESALQ/USP	
C55.1	Cocoa	1	R	-	ESALQ/USP		
C56.1	Cocoa	1	R	-	ESALQ/USP		
ECE5	<i>Pantoea agglomerans</i>	Eucalyptus	8	R	-	Procópio (2004)	
ECE6		Eucalyptus	4	R	+	Procópio (2004)	
ECE7		Eucalyptus	6	R	+	Procópio (2004)	
EGE1		Eucalyptus	10	S	+	Procópio (2004)	
EGE6		Eucalyptus	13	S	+	Procópio (2004)	
EGE8		Eucalyptus	10	S	+	Procópio (2004)	
EGS2		Eucalyptus	17	S	+	Procópio (2004)	
EIE3		Eucalyptus	14	S	+	Procópio (2004)	
EIE6		Eucalyptus	22	S	+	Procópio (2004)	
ETE7		<i>Hafnia alvei</i>	Eucalyptus	6	R	-	Procópio (2004)
ETE8			Eucalyptus	4	R	+	Procópio (2004)
EUE6		<i>Pantoea ananatis</i>	Eucalyptus	18	S	+	Procópio (2004)
EUE7			Eucalyptus	7	R	+	Procópio (2004)
EN80		Soybean	Soybean	2	S	+	Kuklinsky-Sobral <i>et al.</i> (2004)
EN103	Soybean		21	R	+	Kuklinsky-Sobral <i>et al.</i> (2004)	
EN179	Soybean		16	S	+	Kuklinsky-Sobral <i>et al.</i> (2004)	
EN192	Soybean		23	S	+	Kuklinsky-Sobral <i>et al.</i> (2004)	
EN306	Soybean		11	S	+	Kuklinsky-Sobral <i>et al.</i> (2004)	
ENB73	Soybean		8	R	-	Kuklinsky-Sobral <i>et al.</i> (2004)	
2B9	Sugar cane		6	R	+	ESALQ/USP	
64 R1	Sugar cane		20	S	+	ESALQ/USP	
65 R1	Sugar cane	19	S	+	ESALQ/USP		
66 R1	Sugar cane	19	S	+	ESALQ/USP		

<sup>a</sup> All strains are deposited in Bacterial Collection of Laboratory of Microbial Genetics, Department of Genetics, ESALQ/USP, Piracicaba, Brasil, FAX 55 19 3433 6706

<sup>b</sup> ESALQ/USP: isolates from the endophytic bacteria collection of Laboratório de Genética de Microrganismos, ESALQ/USP

- S and R are the susceptibility and resistance to ampicillin, respectively. + and - are the phenotypes of production or not of cellulase, respectively.


series of T-tests were performed according to the methods proposed by Magurran (1988) in order to detect statistical difference in diversity indices.

### Results


#### Cellulase production and antibiotic susceptibility

We characterized 53 enterobacteria endophytes for the production of cellulase and antibiotic susceptibility. The lowest

cellulase production was found for cocoa plants in which 2 out of 10 (20%) isolates were able to produce this enzyme *in vitro*. A similar production percentage was observed for citrus in which 4 out of 20 (20%) isolates produced cellulase. The highest average of cellulase production was observed for soybean (5 out of 6), eucalyptus (11 out of 13), and sugar-cane (4 out of 4), showing 83, 84.6, and 100%, respectively. Similarly, the cellulase activity index (AI) was lower for cocoa followed by citrus. Eucalyptus, soybean, and sugar cane, on



**Fig. 2.** Dendrogram by UPGMA clustering of BOX-PCR fingerprints of 53 bacterial isolates from citrus, cocoa, eucalyptus, soybean, and sugar cane. Clustering using Jaccard product-moment coefficient was expressed as percentage similarity.



**Fig. 3.** Phylogenetic relationships of *Enterobacteriaceae* based on partial 16S rDNA sequences obtained from the endophytic bacteria and closely related sequences, based on a distance analysis (neighbour-joining algorithm with Jukes-Cantor model; 1,000 bootstrap replicates performed).

the other hand, presented higher AI values (Fig. 1). For antibiotic susceptibility, all tested isolates were susceptible to kanamycin. Regarding ampicillin, susceptibility was heterogeneous according to the host plant (Table 1). The susceptibility to this antibiotic was observed as follows: citrus (15%), eucalyptus (54%), soybean (67%), and sugar cane (75%) (data not shown). Cocoa isolates did not exhibit susceptibility to ampicillin (Table 1).

### BOX-PCR DNA fingerprint of endophytic enterobacteria

BOX-PCR was performed on 53 endophytic bacteria belonging to the *Enterobacteriaceae* family isolated from cocoa, citrus, eucalypti, soybean, and sugar cane host plants. Curve-based product-moment correlation coefficients were used for pairwise fingerprint comparison, and UPGMA was used to perform cluster analysis on all isolates (Fig. 2). Through cluster analysis of the BOX-PCR profiles at 55% as arbitrary similarity level, 23 different OTUs (comprising 9 clusters and 14 unique profiles) were identified. We observed that 52.7% of the isolates were included in only 4 OTUs, with OTU9 and OTU12 being most frequent with 24.5 and 13.2% of the isolates.

Representative isolates from the most frequent OTUs were identified by 16S rRNA gene sequencing, which showed that the population evaluated is composed of at least 6 spe-

cies, including *Pantoea agglomerans*, *P. ananatis*, *P. stewartii*, *Enterobacter cancerogenus*, *E. hormaechei*, and *Tatumella pyseos* (Fig. 3). OTU9 comprised of 13 isolates obtained from citrus plants and identified according to biochemical profile as *P. agglomerans* (Araújo *et al.*, 2002). However, 16S rRNA gene sequencing of the isolate TO45 showed that it is close to *Enterobacter cancerogenus* LMG2693<sup>T</sup> (Fig. 3). OTU12 included 6 cocoa and 1 citrus isolates, and the isolate C33.2, which seems to be a divergent genotype in *Enterobacter* genera (Fig. 3). In OTU2, OTU10, OTU19, and OTU23, the isolates EN80, EGE1, 65R1, and EN192, respectively, showed similarity with *P. agglomerans* ATCC 27155<sup>T</sup>, while the isolates included in OTU5 and OTU7 are closed related to *P. ananatis* ATCC 33244<sup>T</sup> and *P. stewartii* subsp. *stewartii* LMG 2715<sup>T</sup> (Fig. 3).

OTU6 had three isolates (one from sugar cane and two from eucalyptus), and since the sequence of isolate ETE7 matched (98%) with unidentified *Enterobacteriaceae* (EU 029106), this group was assumed to be a divergent genotype. However, the phylogenetic tree suggests that this group is related to *Tatumella* genera (Fig. 3).

### Interaction among cellulase, antibiotic phenotypes, and BOX-PCR DNA fingerprint

We analyzed the interaction between phenotypes (cellulase and ampicillin resistance) and genotypes (BOX-PCR DNA

fingerprint) of the evaluated bacteria. Isolates from OTUs 9 and 1 (citrus and cocoa plants, respectively) were resistant to ampicillin and negative for cellulase production (Table 1). However, this result was not observed for OTUs 10 and 12, which showed variable phenotypes. In contrast, except for isolate PR2/7 (OTU 12), ampicillin susceptible isolates showed cellulase production (16 out of 17 isolates).

## Discussion

Understanding microbial biodiversity in association with recombinant DNA technique may provide the basis for utilization of endophytes in agriculture. Detailed molecular diversity information may help us to understand the evolution of microbial functionality in their particular environment, creating new genetic variants below the species level. The selection of newly acquired functions may help us to understand the functional diversity of microbial communities and ultimately ecosystem function as a whole (Schloter *et al.*, 2000).

Our research goals were to characterize endophytic enterobacteria isolates from five commercially important crops for susceptibility to antibiotics and cellulase (endoglucanase) production. We also evaluated the genetic diversity of these isolates using the powerful BOX-PCR technique followed by 16S rRNA gene sequencing.

Initially, we analyzed all the isolates for antibiotic susceptibility and cellulase production. We found susceptibility to both tested antibiotics, which in ampicillin changed according to the host plant. Cellulase production also changed according to the host plant. As such, cellulase production is a characteristic possibly required for endophytic colonization (Germaine *et al.*, 2004), since endophytic colonization is an active process and not simply the passive diffusion of bacteria into the intercellular spaces of plants (Dong *et al.*, 2003). Furthermore, cell wall degradation by endophytes may trigger systemic resistance induction (Hallmann *et al.*, 1997).

According to Germaine *et al.* (2004), endophytic bacteria may lose antibiotic resistance after its introduction in plants. This finding is corroborated by our results, since we observed that in ampicillin sensible population, 94.1% were able to produce cellulase, while in antibiotic resistant population this decreased to 27.8%. Therefore, we suggest that the ability to establish inside the host plant, by cell wall degradation, may be related to the loss of some phenotypes, such as antibiotic resistance.

After this, BOX-PCR was selected as the molecular typing technique for the enterobacteria endophytic isolates. This is because BOX-PCR is reproducible, rapid, easy to perform, and highly discriminatory at the subspecies level (Olive and Bean, 1999), yielding results that correlate well with pairwise DNA-DNA analyses (Rademaker *et al.*, 2000). The BOX-PCR genomic fingerprint patterns were analyzed by a curve-based protocol, which retains more information than merely the number and position of fingerprint fragments (Hane *et al.*, 1993; Yang *et al.*, 2004). In the present study, 23 OTUs were observed in 53 isolates analyzed by BOX-PCR fingerprint. OTU9 and OTU1 were observed only in citrus and cocoa bacterial population, respectively.

The isolates belonging to these OTUs didn't produced cellulase and were ampicillin resistant. Also, all cocoa isolates belonged to OTU1 and OTU12, and showed ampicillin resistance. These results suggest that correlation between host genotype, and genetic and phenotypic characteristic of the bacteria, could drive the endophytic-plant interaction.

In our work, the genetic diversity of 53 isolates analyzed by BOX-PCR fingerprint grouped in 23 different OTUs mostly showed a non-strictly OTU-host relationship. This had implications in the diversity indices of these hosts, which was higher for eucalyptus and soybean isolates. According to Yang *et al.* (2004), the presence of 23 OTUs could indicate that the 53 isolates are comprised of up to 23 different species, since when microbial diversity is inferred from BOX-PCR, individual OTUs must be defined as species surrogates. However, no consistent definition of OTUs is currently employed, leading to a non-comparable estimate. Here, isolates representing nine different OTUs was identified by 16S rRNA gene sequencing, belonged to *P. agglomerans*, *P. ananatis*, *P. stewartii*, *Enterobacter cancerogenus*, *E. hormaechei*, and *Tatumella pyseos*. This suggests that the OTUs observed in our work could be species closely related to the *Pantoea* genus or subspecies of *P. agglomerans* and *P. ananatis*. This hypothesis is supported by Yang *et al.* (2004) who state that BOX-PCR is a higher genomic resolution technique for diversity estimation within species, which means that it is powerful for subspecies diversity examination.

In this study, we observed high variability in endophytic *Enterobacteriaceae*, including OTUs that share less than 20% of the BOX-PCR bands. Furthermore, the *P. agglomerans* strains were not in the same group, being the type strain JCM1236 closely related to *E. hormaechei*. These results suggest that some species, such as *P. agglomerans* and *P. stewartii*, could be polyphyletic.

In summary, the BOX-PCR technology was highly successful in discriminating closely related bacteria from the *Enterobacteriaceae* family. Finally, it is hoped that this line of research will be continued and expanded in order to provide further interesting insights into the understanding of the endophytic enterobacteria community from other commercially important crops.

## Acknowledgements

Funding for this research was provided by the FAPESP project no. 02/14143-3. We thank M.C. Quecine (ESALQ/USP) for assistance and production of statistical analysis. We also gratefully acknowledge CAPES and CNPQ for providing the scholarship to A.R. Torres and L. Cursino, and FAPESP for providing the fellowship to W.L. Araújo.

## References

- Araújo, W.L., W. Maccheroni, C.I. Aguilar-Vildoso, P.A.V. Barroso, H.O. Saridakis, and J.L. Azevedo. 2001. Variability and interactions between endophytic bacteria and fungi isolated from leaf tissues of citrus rootstocks. *Can. J. Microbiol.* 47, 229-236.
- Araújo, W.L., J. Marcon, W. Maccheroni, J.D. Van Elsas, J.W.L. Van Vuurde, and J.L. Azevedo. 2002. Diversity of endophytic bacterial populations and their interaction with *Xylella fastidiosa* in citrus plants. *Appl. Environ. Microbiol.* 68, 4906-4914.

- Berg, G., B. Roskot, A. Steidle, L. Eberl, A. Zock, and K. Smalla. 2002. Plant-dependent genotypic and phenotypic diversity of antagonistic rhizobacteria isolated from different *Verticillium* host plants. *Appl. Environ. Microbiol.* 68, 3328-3338.
- Carson, C.A., B.L. Shear, M.R. Ellersieck, and J.D. Schnell. 2003. Comparison of ribotyping and repetitive extragenic palindromic-PCR for identification of fecal *Escherichia coli* from humans and animals. *Appl. Environ. Microbiol.* 69, 1836-1839.
- Davelos, A.L., L.L. Kinkel, and D.A. Samac. 2004. Spatial variation in the frequency and intensity of antibiotic interactions among Streptomycetes from prairie soil. *Appl. Environ. Microbiol.* 70, 1051-1058.
- Dombek, P.E., L.K. Johnson, S.T. Zimmerley, and M.J. Sadowsky. 2000. Use of repetitive DNA sequences and the PCR to differentiate *Escherichia coli* isolates from human and animal sources. *Appl. Environ. Microbiol.* 66, 2572-2577.
- Dong, Y., A.L. Iniguez, and E.W. Triplett. 2003. Quantitative assessments of the host range and strain specificity of endophytic colonization by *Klebsiella pneumoniae* 342. *Plant Soil* 257, 49-59.
- Germaine, K., E. Keogh, G. Garcia-Cabellos, B. Borremans, D. Van Der Lelie, T. Barac, L. Oeyen, J. Vangronsveld, F.P. Moore, E.R.B. Moore, C.D. Campbell, D. Ryan, and D.N. Dowling. 2004. Colonisation of poplar trees by *gfp* expressing bacterial endophytes. *FEMS Microbiol. Ecol.* 48, 109-118.
- Hallmann, J., A. Quadt-Hallmann, W.F. Mahaffee, and J.W. Kloepper. 1997. Bacterial endophytes in agricultural crops. *Can. J. Microbiol.* 43, 895-914.
- Hane, B.G., K. Jager, and H.G. Drexler. 1993. The Pearson product moment correlation coefficient is better suited for identification of DNA fingerprints profiles than band matching algorithms. *Electrophoresis* 14, 967-972.
- Jukes, T.H. and C.R. Cantor. 1969. Evolution of protein molecules, p. 21-132. In H.N. Munro (ed.), *Mammalian protein metabolism*. Academic Press, New York, N.Y., USA.
- Koeuth, T., J. Versalovic, and J.R. Lupski. 1995. Differential subsequence conservation of interspersed repetitive *Streptococcus pneumoniae* BOX elements in diverse bacteria. *Genome Res.* 5, 408.
- Krechel, A., A. Faupel, J. Hallmann, A. Ulrich, and G. Berg. 2002. Potato-associated bacteria and their antagonistic potential towards plant-pathogenic fungi and the plant-parasitic nematode *Meloidogyne incognita* (Kofoid & White) Chitwood. *Can. J. Microbiol.* 48, 772-785.
- Kuklinsky-Sobral, J., W.L. Araújo, R. Mendes, I.O. Geraldi, A.P. Pizzirani-Kleiner, and J.L. Azevedo. 2004. Isolation and characterization of soybean-associated bacteria and their potential for plant growth promotion. *Environ. Microbiol.* 6, 1244-1251.
- Lanoot, B., M. Vancanneyt, P. Dawyndt, M. Cnockaert, J. Zhang, Y. Huang, Z. Liu, and J. Swings. 2004. BOX-PCR fingerprinting as a powerful tool to reveal synonymous names in the genus streptomycetes. Emended descriptions are proposed for the species *Streptomyces cinereorectus*, *S. fradiae*, *S. tricolor*, *S. colombiensis*, *S. filamentosus*, *S. vinaceus*, and *S. phaeoauripureus*. *Syst. Appl. Microbiol.* 27, 84-92.
- Margurran, A.E. 1998. *Ecological diversity and its measurements*. Princeton University Press, Princeton, New Jersey, USA.
- NCCLS - National Committee for Clinical Laboratory Standards. Performance Standards for Antimicrobial Susceptibility Testing. 15<sup>th</sup> ed. NCCLS document M. 100 - S15: National Committee for Clinical Laboratory Standards, 2004 Villanova, PA, USA.
- Olive, D.M. and P. Bean. 1999. Principles and applications of methods for DNA-based typing of microbial organisms. *J. Clin. Microbiol.* 37, 1661-1669.
- Procópio, R.E.L. 2004. Diversidade bacteriana endofítica de *Eucahyptus* spp. e avaliação do seu potencial biotecnológico. Ph. D. thesis. Universidade de São Paulo, São Paulo, Brazil.
- Rademaker, J.L.W., B. Hoste, F.J. Louws, K. Kertters, J. Swings, L. Vauterin, P. Vauterin, and F.J. De Bruijn. 2000. Comparison of AFLP and rep-PCR genomic fingerprints with DNA-DNA homology studies: *Xanthomonas* as a model system. *Int. J. Syst. Evol. Microbiol.* 50, 665-677.
- Saitou, N. and M. Nei. 1987. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* 4, 406-425.
- SAS Institute, Inc. 1987. *SAS/STAT guide for personal computers*, version 6. SAS Institute, Inc., Cary, N.C., USA.
- Schlöter, M., M. Leubhn, T. Heulin, and A. Hartmann. 2000. Ecology and evolution of bacterial microdiversity. *FEMS Microbiol. Rev.* 24, 647-660.
- Teather, R.M. and P.J. Wood. 1982. Use of Congo red-polysaccharide interactions in enumeration and characterization of cellulolytic bacteria from bovine rumen. *Appl. Environ. Microbiol.* 43, 777-780.
- Terasawa, G.L.V., B.C. Glienke, and M. Hungria. 2003. Diversity of a soybean rhizobial population adapted to a cerrados soil. *World J. Microbiol. Biotechnol.* 19, 933-939.
- Vanechoutte, M., P. Boerlin, H.V. Tichy, E. Bannerman, B. Jager, and J. Bille. 1998. Comparison of PCR-based DNA fingerprinting techniques for the identification of *Listeria* species and their use for atypical *Listeria* isolates. *Int. J. Syst. Bacteriol.* 48, 127-139.
- Verma, S.C., J.K. Ladha, and A.K. Tripathi. 2001. Evaluation of plant growth promoting and colonization ability of endophytic diazotrophs from deep water rice. *J. Biotechnol.* 91, 127-141.
- Wise, M.G., J. Vaun McArthur, C. Wheat, and L.J. Shimkets. 1996. Temporal variation in genetic diversity and structure of a lotic population of *Burkholderia (Pseudomonas) cepacia*. *Appl. Environ. Microbiol.* 62, 1558-1562.
- Yang, H.H., R.T. Vinopal, D. Grasso, and B.F. Smets. 2004. High diversity among environmental *Escherichia coli* isolates from a bovine feedlot. *Appl. Environ. Microbiol.* 70, 1528-1536.