

Rhizobium freirei sp. nov., a symbiont of *Phaseolus vulgaris* that is very effective at fixing nitrogen

Rebeca Fuzinato Dall'Agnol,^{1,2†} Renan Augusto Ribeiro,^{1,3†}
Ernesto Ormeño-Orrillo,^{4†} Marco Antonio Rogel,⁴
Jakeline Renata Marçon Delamuta,^{1,3} Diva Souza Andrade,⁵
Esperanza Martínez-Romero⁴ and Mariangela Hungria^{1,2,3}

Correspondence

Mariangela Hungria
mariangela.hungria@embrapa.br
or hungria@pq.cnpq.br

¹Embrapa Soja, C.P. 231, 86001-970 Londrina, Paraná, Brazil

²Universidade Estadual de Londrina, Dept of Biochemistry and Biotechnology, C.P. 6001, 86051-990 Londrina, Paraná, Brazil

³Universidade Estadual de Londrina, Dept of Microbiology, C.P. 60001, 86051-990 Londrina, Paraná, Brazil

⁴Centro de Ciencias Genómicas, Universidad Nacional Autónoma de México, Cuernavaca, Morelos, Mexico

⁵IAPAR, C.P. 481, 86001-970 Londrina, Paraná, Brazil

Common bean (*Phaseolus vulgaris* L.) can establish symbiotic associations with several *Rhizobium* species; however, the effectiveness of most strains at fixing nitrogen under field conditions is very low. PRF 81^T is a very effective strain, usually referred to as *Rhizobium tropici* and used successfully in thousands of doses of commercial inoculants for the common bean crop in Brazil; it has shown high rates of nitrogen fixation in all areas representative of the crop in the country. Here, we present results that indicate that PRF 81^T, although it belongs to the 'R. tropici group', which includes 10 *Rhizobium* species, *R. tropici*, *R. leucaenae*, *R. lusitanum*, *R. multihospitium*, *R. miluonense*, *R. hainanense*, *R. calliandrae*, *R. mayense*, *R. jaguaris* and *R. rhizogenes*, represents a novel species. Several morpho-physiological traits differentiated PRF 81^T from related species. Differences were also confirmed in the analysis of rep-PCR (sharing less than 45 % similarity with the other species), MLSA with *recA*, *atpD* and *rpoB* genes, and DNA–DNA hybridization. The novel species, for which we propose the name *Rhizobium freirei* sp. nov., is able to establish effective root nodule symbioses with *Phaseolus vulgaris*, *Leucaena leucocephala*, *Leucaena esculenta*, *Crotalaria juncea* and *Macroptilium atropurpureum*. The type strain is PRF 81^T (=CNPSO 122^T=SEMIA 4080^T=IPR-Pv81^T=WDCM 440^T).

Common bean (*Phaseolus vulgaris* L.) represents a major source of protein for poor populations of South and Central America, Africa and India. An important characteristic of common bean is its capacity to establish symbiotic partnerships with a variety of *Rhizobium* species

(Martínez-Romero, 2003). These symbioses include both Fix⁺ (*Rhizobium leguminosarum* sv. phaseoli, *R. phaseoli*, *R. tropici*, *R. etli*, *R. leucaenae*, *R. giardinii* sv. phaseoli, *R. gallicum*, *R. lusitanum* and *R. pisi*) and Fix⁻ (*R. giardinii* sv. giardinii and *R. miluonense*) species. This promiscuous nature (Michiels *et al.*, 1998) is probably responsible for the commonly reported low contribution of biological N₂ fixation to common bean nutrition, as the legume often nodulates with very competitive but inefficient indigenous rhizobia (Graham, 1981; Hardarson, 1993).

Strains belonging to the broad-host-range species *R. tropici* have been used as inoculants in the tropics because of their high tolerance to environmentally stressful conditions and high genetic stability (Hungria *et al.*, 2000, 2003). The most competitive and effective strain used in commercial inoculants for common bean in Brazil is PRF 81^T, which

†These authors contributed equally to this work.

Abbreviations: ANI, average nucleotide identity; DDH, DNA–DNA hybridization; IGS, intergenic spacer; MLSA, multilocus sequence analysis.

The GenBank/EMBL/DDBJ accession numbers for the 16S rRNA and *recA* gene sequences of strain PRF 81^T are EU488742 and EU488827. The accession number of the whole genome shotgun sequencing project for strain PRF 81^T is AQHN00000000.

Four supplementary tables and two supplementary figures are available with the online version of this paper.

is closely related to *R. tropici* on the basis of 16S rRNA gene sequence analysis; however, it may be distinguished from *R. tropici* by phenotypic and genetic characteristics (Hungria *et al.*, 2000; Ribeiro *et al.*, 2009). A recent comparison of the genome of PRF 81^T to that of *R. tropici* CIAT 899^T highlighted the high similarity of their symbiotic plasmids, but considerable differences between the rest of their genomes, with low average nucleotide identity (ANI) and *in silico* DNA–DNA hybridization (DDH) (Ormeño-Orrillo *et al.*, 2012). High divergence of PRF 81^T from other rhizobial species has also been revealed in the reference proteomic map of this strain (Gomes *et al.*, 2012) and by multilocus sequence analysis (MLSA) (Ribeiro *et al.*, 2009).

Strain PRF 81^T was isolated in Irati county, State of Paraná, Brazil (25° 28' 01" S 50° 37' 03" W; Köppen's classification of climate *Cfb*; altitude 812 m), by one of us (D. S. A.) on 28 May 1992. The DNA G+C content of PRF 81^T was determined previously as 59.9 mol% (Ormeño-Orrillo *et al.*, 2012), which is within the range reported for *Rhizobium* (Jordan, 1984). In extensive common bean and *Leucaena* spp. rhizobial isolations performed in our laboratory, we have never found other strains showing high similarity with PRF 81^T in rep-PCR profiles; however, we have obtained isolates showing identical or nearly identical profiles to PRF 81^T (>90% in rep-PCR) in several soils from Brazil (states of Pernambuco, Goiás, Santa Catarina, Paraná, Mato Grosso do Sul, Federal District). Examples of unique rep-PCR profiles obtained in our studies are given elsewhere (Pinto *et al.*, 2007; Stocco *et al.*, 2008). These isolates possessed sequences identical to PRF 81^T in their 16S rRNA and housekeeping genes (data not shown). This seemed to indicate that PRF 81^T is a member of a broadly distributed *Rhizobium* clonal group that nodulates common bean as well as leucaenas. Due to its relevant economic importance, we sought to establish the taxonomic position of PRF 81^T as a representative of this group, using a polyphasic approach with data published by us and new evidence obtained in this study.

16S rRNA gene sequences from all *Rhizobium* type strains were retrieved from the GenBank database and aligned with MUSCLE (Edgar, 2004). Neighbour-joining (Saitou & Nei, 1987) and maximum-likelihood (Felsenstein, 1981) phylogenies were reconstructed with MEGA (Tamura *et al.*, 2011) using the Tamura–Nei model (Tamura & Nei, 1993). The 16S rRNA gene sequence of PRF 81^T was part of a clade with bootstrap support of 98% (Figs 1 and S1, available in IJSEM Online) that included all species of the '*R. tropici* group' (Ribeiro *et al.*, 2012). The 16S rRNA gene sequence identity between PRF 81^T and *R. multihospitium* CCBAU 83401^T was 100%, and identity ranged from 98.8 to 99.8% with the remaining species of the clade (Table S1). It was surprising to find such level of identity with *R. multihospitium* CCBAU 83401^T, as the similarity between the BOX-PCR profiles of PRF 81^T and *R. multihospitium* CCBAU 83401^T was lower than 45% (Fig. S2). It is known that phylogenetic analysis based exclusively on the 16S

rRNA gene often does not reflect the correct taxonomic position of a species (Coenye *et al.*, 2001; Martens *et al.*, 2007; Menna *et al.*, 2009; Wang & Martínez-Romero, 2000). We tested whether the similarity between PRF 81^T and *R. multihospitium* also included other parts of the ribosomal (*rrn*) operon by comparing the sequences of their 16S–23S intergenic spacer (IGS) regions. The available IGS sequences of four *R. multihospitium* strains were identical to each other (Han *et al.*, 2008), as were the sequences of the three *rrn* operons in the PRF 81^T genome (Ormeño-Orrillo *et al.*, 2012). Interestingly, the PRF 81^T and *R. multihospitium* IGS sequences were only 76.4% identical (Table S1), indicating that they may have recombined at their 16S rRNA gene loci at some point in their evolutionary history. PRF 81^T showed the highest IGS sequence identity to *R. hainanense* (77.5% to the type strain), while *R. multihospitium* showed the highest identity to *R. tropici* (84.1% to the type strain) (Table S1).

An MLSA approach with the *recA*, *atpD* and *rpoB* genes was used to explore the relationships of PRF 81^T and related species further. Sequences were retrieved from GenBank or extracted from the PRF 81^T genome, and phylogenies were constructed as described above for the




Fig. 1. Part of a neighbour-joining phylogeny of 16S rRNA gene sequences (1308 nt) of *R. freirei* sp. nov. PRF 81^T and other *Rhizobium* species (the complete phylogram is presented in Fig. S1). GenBank accession numbers are given in parentheses. Bootstrap support values based on 500 resamplings are shown as percentages at nodes only when $\geq 60\%$. Nodes with adjacent filled circles also had $\geq 60\%$ support in a separate maximum-likelihood phylogeny. Bar, 0.005 substitutions per nucleotide position.




Fig. 2. Neighbour-joining phylogeny based on a concatenated alignment of *recA* (442 nt), *atpD* (437 nt) and *rpoB* (650 nt) gene sequences of *R. freirei* sp. nov. PRF 81^T and closely related *Rhizobium* species. GenBank accession numbers are given in parentheses (*recA*, *atpD*, *rpoB*). Bootstrap support values based on 500 resamplings are shown as percentages at nodes only when $\geq 60\%$. Nodes with adjacent filled circles also had $\geq 60\%$ support in a separate maximum-likelihood phylogeny. Bar, 0.02 substitutions per nucleotide position.

Table 1. Strains used in this study for phenotypic characterization

Strain	Other strain nomenclature	Host species	Geographical origin	Reference or source
<i>R. freirei</i> sp. nov. PRF 81 ^T	CNPSo 122 ^T =SEMIA 4080 ^T =IPR-Pv81 ^T =WDCM 440 ^T	<i>Phaseolus vulgaris</i>	Paraná, Brazil	Hungria <i>et al.</i> (2000)
<i>R. leucaenae</i> CFN 299 ^T	USDA 9039 ^T =LMG 9517 ^T =UMR1026 ^T =CENA 183 ^T =SEMIA 4083 ^T =CNPSo 141 ^T	<i>P. vulgaris</i>	Brazil	Ribeiro <i>et al.</i> (2012)
<i>R. tropici</i> CIAT 899 ^T	USDA 9030 ^T =ATCC 49672 ^T =UMR1899 ^T =TAL 1797 ^T =HAMBI 1163 ^T =CM0 ^T =SEMIA 4077 ^T =DSM 11418 ^T =BR 322 ^T =CNPSo 142 ^T	<i>P. vulgaris</i>	Colombia	Martínez-Romero <i>et al.</i> (1991)
<i>R. rhizogenes</i> ATCC 11325 ^T	DSM 30148 ^T =LMG 150 ^T =IFO 13257 ^T =IAM 13570 ^T =CNPSo 1976 ^T	Apple	NK	Velázquez <i>et al.</i> (2010)
<i>R. hainanense</i> I66 ^T	BCRC 15793 ^T =CCRC 15793 ^T =CECT 4658 ^T =DSM 11917 ^T =ICMP 13690 ^T =LMG 18074 ^T =USDA 3588 ^T =CNPSo 2450 ^T	<i>Desmodium sinuatum</i>	Hainan, China	Chen <i>et al.</i> (1997)
<i>R. lusitanum</i> P1-7 ^T	CECT 7016 ^T =CIP 109524 ^T =LMG 22705 ^T =CNPSo 2055 ^T	<i>P. vulgaris</i>	Arcos de Valdevez, Portugal	Valverde <i>et al.</i> (2006)
<i>R. multihospitium</i> CCBAU 83401 ^T	HAMBI 2975 ^T =LMG 23946 ^T =LMG 24298 ^T =CNPSo 2054 ^T	<i>Halimodendron halodendron</i>	Xinjiang, China	Han <i>et al.</i> (2008)
<i>R. miluonense</i> CCBAU 41251 ^T	HAMBI 2971 ^T =LMG 24208 ^T =CNPSo 2056 ^T	<i>Lespedeza chinensis</i>	Hunan, China	Gu <i>et al.</i> (2008)
<i>R. calliandrae</i> CCGE 524 ^T	ATCC BAA-2435 ^T =CIP110456 ^T =LBP2-1 ^T =CNPSo 2466 ^T	<i>Calliandra grandiflora</i>	Chiapas, Mexico	Rincón-Rosales <i>et al.</i> (2013)
<i>R. mayense</i> CCG E526 ^T	CIP 110454 ^T =NSJP1-1 ^T =CNPSo 2464 ^T	<i>C. grandiflora</i>	Chiapas, Mexico	Rincón-Rosales <i>et al.</i> (2013)
<i>R. jaguaris</i> CCG E525 ^T	ATCC BAA-2445 ^T =CIP 110453 ^T =SJP1-2 ^T =CNPSo 2465 ^T	<i>C. grandiflora</i>	Chiapas, Mexico	Rincón-Rosales <i>et al.</i> (2013)

NK, Not known.

Table 2. Distinctive phenotypic features of *Rhizobium freirei* sp. nov. PRF 81^T and type strains of phylogenetically related species of the '*R. tropici* group'

Strains: 1, *R. freirei* PRF 81^T; 2, *R. leucaenae* CFN 299^T; 3, *R. tropici* CIAT 899^T; 4, *R. lusitanum* P1-7^T; 5, *R. multihospitium* CCBAU 83401^T; 6, *R. miluonense* CCBAU 41251^T; 7, *R. rhizogenes* ATCC 11325^T; 8, *R. hainanense* I66^T; 9, *R. calliandrae* CCGE 524^T; 10, *R. mayense* CCGE 526^T; 11, *R. jaguaris* CCGE 525^T. +, Growth, -, no growth; w, weak growth; ND, not determined. Data were obtained in this study. All strains are sensitive to tetracycline (30 µg).

Characteristic	1	2	3	4	5	6	7	8	9	10	11
Carbon source											
Glycerol	+	+	+	+	+	+	w	+	+	w	+
Erythritol	+	-	+	+	+	+	+	+	+	+	+
D-Arabinose	w	+	+	w	+	+	+	+	w	w	w
D-Xylose	w	+	+	+	+	+	w	+	w	w	w
L-Xylose	+	+	-	+	+	+	+	+	w	w	w
D-Adonitol	+	+	+	+	+	w	+	+	w	w	w
Methyl β-D-xylopyranoside	w	+	+	+	+	+	+	+	w	+	w
D-Glucose	+	+	+	+	+	+	w	+	+	w	w
D-Fructose	w	+	+	+	+	+	w	+	+	w	w
D-Mannose	+	+	+	+	+	+	w	+	+	w	w
L-Sorbose	w	-	-	w	w	+	w	w	-	-	-
L-Rhamnose	w	+	+	+	+	+	w	+	w	+	+
Dulcitol	-	w	-	-	w	w	w	w	w	-	-
Inositol	w	+	+	+	+	+	w	w	w	w	w
D-Mannitol	w	+	+	w	+	+	w	+	+	+	w
D-Sorbitol	w	-	-	w	w	+	w	+	-	w	w
Methyl α-D-mannopyranoside	-	-	-	w	w	-	-	-	-	-	-
Methyl α-D-glucopyranoside	w	-	+	w	+	+	+	+	w	w	w
N-Acetylglucosamine	+	-	-	-	+	+	+	w	-	w	-
Amygdalin	-	-	-	w	w	w	+	+	-	w	-
Arbutin	+	-	w	+	+	+	+	+	w	w	w
Salicin	+	-	w	+	+	+	+	w	w	w	w
Cellobiose	w	w	+	+	+	+	+	+	+	+	w
Maltose	w	w	+	+	+	+	w	+	w	w	w
Lactose	w	w	w	+	+	+	w	w	w	w	w
Melibiose	w	w	w	+	+	+	w	w	w	w	w
Sucrose	w	w	+	+	+	+	+	w	w	w	w
Trehalose	w	w	+	+	+	+	+	+	w	w	w
Melezitose	-	-	-	-	-	-	-	w	-	-	-
Raffinose	w	w	w	+	+	+	w	w	w	w	w
Glycogen	+	-	-	+	+	+	+	+	+	+	+
Xylitol	w	-	-	w	w	+	w	+	-	w	-
Gentiobiose	w	w	w	+	+	+	+	+	w	w	w
D-Turanose	w	w	w	w	+	+	w	+	w	w	w
D-Lyxose	w	+	+	+	+	+	+	+	+	+	+
D-Tagatose	w	w	-	w	w	+	+	w	w	-	-
D-Arabitol	w	+	+	w	+	+	w	+	w	w	w
L-Arabitol	+	-	w	w	w	w	+	+	-	-	-
Resistance to:											
Nalidixic acid (30 µg)	+	+	+	+	+	+	-	w	+	+	+
Ampicillin (10 µg)	w	-	+	-	+	+	-	-	-	-	-
Chloramphenicol (30 µg)	-	-	+	-	+	-	w	+	w	-	-
Cephuroxyn (30 µg)	+	-	+	+	+	+	-	-	-	-	-
Erythromycin (15 µg)	+	+	+	+	+	+	+	+	+	+	-
Streptomycin (10 µg)	-	-	+	-	w	w	w	+	-	-	-
Neomycin (30 µg)	w	-	-	w	-	w	-	w	w	+	w
Penicillin (10 U)	+	-	+	+	+	+	+	+	+	w	w
Growth in/at:											
PY without Ca medium	w	-	+	+	+	+	+	+	-	-	-

Table 2. cont.

Characteristic	1	2	3	4	5	6	7	8	9	10	11
LB medium	w	–	+	+	+	–	–	+	–	–	–
37 °C	+	+	+	+	+	+	–	+	–	–	–
40 °C	+	w	+	–	–	–	–	+	–	–	–
1 % NaCl	+	–	+	w	+	–	–	+	–	–	–
pH 4	+	w	+	–	+	–	ND	–	–	–	–

16S rRNA gene. In the concatenated-alignment phylogeny, PRF 81^T was grouped with *R. tropici*, *R. multihospitium*, *R. hainanense* and *R. miluonense* in a subclade with bootstrap support of 99% (Fig. 2) that was part of a clade that included all species of the '*R. tropici* group'. The range of nucleotide identity between described species of this clade, calculated from the concatenated genes, was 85.4–96% (Table S1). Konstantinidis *et al.* (2006) found that 96% ANI in concatenated housekeeping genes is equivalent to 70% DDH. PRF 81^T showed the highest identity with *R. hainanense* (95.5% to the type strain), while it showed 95.2% identity or less to all other species, indicating that this strain is a member of a species different from all others of the '*R. tropici* group'. In order to ascertain that PRF 81^T represents a distinct species, DDH experiments were performed using a filter hybridization methodology (Martínez-Romero *et al.*, 1991) to estimate the overall DNA relatedness with type strains of related species. DDH values obtained were lower than 70% with all type strains tested, including those of *R. multihospitium* and *R. hainanense* (Table S2), indicating that PRF 81^T represents a novel species.

Comparative phenotypic characterization of PRF 81^T and strains of related species (Table 1) was performed by using previously described methodologies (Hungria *et al.*, 2001; Ribeiro *et al.*, 2012). All tests were performed in triplicate. Differential phenotypic characteristics are displayed in Table 2. Additionally, Table S3 shows the differences found between PRF 81^T and well-known symbionts of common bean. It is interesting to observe that PRF 81^T could be easily distinguished from *R. tropici*, the species in which it is usually classified, from *R. multihospitium*, which has an identical 16S rRNA gene sequence, and from *R. hainanense*, which shares 95.5% ANI. PRF 81^T differs from *R. tropici* CIAT 899^T in mucus production, colour and morphology of colonies (Table S3); likewise, it grows very poorly in PY minus Ca and LB medium, while *R. tropici* CIAT 899^T grows abundantly in those media. The two latter characteristics also distinguished PRF 81^T from *R. multihospitium* CCBAU 83401^T and *R. hainanense* 166^T, as well as its ability to grow at 40 °C, in contrast to *R. multihospitium* CCBAU 83401^T. Additionally, differential utilization of 24, 23 and 21 out of the 49 carbon sources tested with the API 50CH kit can be used to distinguish PRF 81^T from *R. tropici* CIAT 899^T, *R. multihospitium* CCBAU 83401^T and *R. hainanense* 166^T, respectively, as well as different tolerance to four of five antibiotics.

The fatty acid profiles of PRF 81^T, *R. tropici* CIAT 899^T, *R. multihospitium* CCBAU 83401^T and *R. hainanense* 166^T were obtained with the MIDI system using FAME library TSBA6 of a culture grown for 5 days at 28 °C on YM agar plates. The results shown in Table S4 indicate that all strains had summed feature 8 (C_{18:1}ω7c/C_{18:1}ω6c) as a major fatty acid, like other *Rhizobium* strains (Tighe *et al.*, 2000); however, differences in the relative amounts of some fatty acids seem to be useful to distinguish the strains. For example, PRF 81^T had significant less C_{19:0} cyclo ω8c than *R. tropici* CIAT 899^T and *R. multihospitium* CCBAU 83401^T, and about half the content of summed feature 2 (C_{14:0} 3OH/16:1 iso I) compared with *R. hainanense* 166^T (Table S4).

In view of the genotypic and phenotypic data presented here, which clearly indicate the distinctiveness of PRF 81^T in relation to all described species of *Rhizobium*, we propose to create a separate species, named *Rhizobium freirei* sp. nov., to accommodate this strain.

Description of *Rhizobium freirei* sp. nov.

Rhizobium freirei (frei're.i. N.L. masc. gen. n. *freirei* of Freire, named after Professor João Ruy Jardim Freire, a distinguished Brazilian rhizobiologist).

Cells are Gram-negative, aerobic, non-spore-forming rods. Colonies on YMA medium are circular, convex, rose, opaque and gummy, with moderate production of mucus, and usually 2–4 mm in diameter within 2–3 days of incubation at 28 °C. It acidifies YMA medium after 3 days. It can tolerate 40 °C and grow at pH 4; optimum growth occurs at pH 5–7 and at 25–28 °C. Grows weakly in LB medium and PY minus Ca medium and is sensitive to chloramphenicol (30 µg), streptomycin (10 µg) and tetracycline (30 µg) and resistant to nalidixic acid (30 µg), cephalosporin (30 µg), erythromycin (15 µg) and penicillin (10 U). As sources of carbon, uses glycerol, erythritol, L-arabinose, D-ribose, L-xylose, D-adonitol, D-galactose, D-glucose, D-mannose, N-acetylglucosamine, arbutin, aesculin, salicin, starch, glycogen, D- and L-fucose and L-arabitol; weakly utilizes D-arabinose, D-xylose, methyl β-D-xylopyranoside, D-fructose, L-sorbose, L-rhamnose, inositol, D-mannitol, D-sorbitol, methyl α-D-glucopyranoside, cellobiose, maltose, lactose, melibiose, sucrose, trehalose, raffinose, xylitol, gentiobiose, turanose, D-lyxose, D-tagatose and D-arabitol. Induces the formation of root nodules and fixes N₂ with *Phaseolus vulgaris*, *Leucaena leucocephala*,

Leucaena esculenta, *Crotalaria juncea* and *Macroptilium atropurpureum*.

The type strain is PRF 81^T (=CNPSO 122^T=SEMIA 4080^T=IPR-Pv81^T=WDCM 440^T), isolated from an effective nodule of *Phaseolus vulgaris* in Paraná State, Brazil. The DNA G+C content of the type strain is 59.9 mol%.

Acknowledgements

The study was partially supported by CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico, Brazil), Project Repensa (562008/2010-1). The authors acknowledge Ligia Maria O. Chueire for help in several steps of this work, Dr Allan R. J. Eaglesham for suggestions on the manuscript, Dr Itamar Soares de Melo for help in the fatty acid analysis and Professor Jean P. Euzéby for help in the creation of the specific epithet. R. F. D. and J. R. M. D. acknowledge PhD fellowships from CAPES. M. H. is also a research fellow of CNPq (300547/2010-2). Embrapa Soja and Centro de Ciências Genômicas had a bilateral collaboration project CNPq/CONACYT (490048/2009-9).

References

- Chen, W.-X., Tan, Z.-Y., Gao, J.-L., Li, Y. & Wang, E.-T. (1997). *Rhizobium hainanense* sp. nov., isolated from tropical legumes. *Int J Syst Bacteriol* **47**, 870–873.
- Coenye, T., Vandamme, P., Govan, J. R. & LiPuma, J. J. (2001). Taxonomy and identification of the *Burkholderia cepacia* complex. *J Clin Microbiol* **39**, 3427–3436.
- Edgar, R. C. (2004). MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* **32**, 1792–1797.
- Felsenstein, J. (1981). Evolutionary trees from DNA sequences: a maximum likelihood approach. *J Mol Evol* **17**, 368–376.
- Gomes, D. F., Batista, J. S. D., Torres, A. R., de Souza Andrade, D., Galli-Terasawa, L. V. & Hungria, M. (2012). Two-dimensional proteome reference map of *Rhizobium tropici* PRF 81 reveals several symbiotic determinants and strong resemblance with agrobacteria. *Proteomics* **12**, 859–863.
- Graham, P. H. (1981). Some problems of nodulation and symbiotic nitrogen-fixation in *Phaseolus vulgaris* L. – a review. *Field Crops Res* **4**, 93–112.
- Gu, C. T., Wang, E. T., Tian, C. F., Han, T. X., Chen, W. F., Sui, X. H. & Chen, W. X. (2008). *Rhizobium miluonense* sp. nov., a symbiotic bacterium isolated from *Lespedeza* root nodules. *Int J Syst Evol Microbiol* **58**, 1364–1368.
- Han, T. X., Wang, E. T., Wu, L. J., Chen, W. F., Gu, J. G., Gu, C. T., Tian, C. F. & Chen, W. X. (2008). *Rhizobium multihospitium* sp. nov., isolated from multiple legume species native of Xinjiang, China. *Int J Syst Evol Microbiol* **58**, 1693–1699.
- Hardarson, G. (1993). Methods for enhancing symbiotic nitrogen-fixation. *Plant Soil* **152**, 1–17.
- Hungria, M., Andrade, D. S., Chueire, L. M. O., Probanza, A., Guttierrez-Mañero, F. J. & Megias, M. (2000). Isolation and characterization of new efficient and competitive bean (*Phaseolus vulgaris* L.) rhizobia from Brazil. *Soil Biol Biochem* **32**, 1515–1528.
- Hungria, M., Chueire, L. M. O., Coca, R. G. & Megias, M. (2001). Preliminary characterization of fast growing rhizobial strains isolated from soyabean nodules in Brazil. *Soil Biol Biochem* **33**, 1349–1361.
- Hungria, M., Campo, R. J. & Mendes, I. C. (2003). Benefits of inoculation of the common bean (*Phaseolus vulgaris*) crop with efficient and competitive *Rhizobium tropici* strains. *Biol Fertil Soils* **39**, 88–93.
- Jordan, D. C. (1984). Family III. *Rhizobiaceae* Conn 1938. In *Bergey's Manual of Systematic Bacteriology*, vol. 1, pp. 234–235. Edited by N. R. Krieg & J. G. Holt. Baltimore: Williams & Wilkins.
- Konstantinidis, K. T., Ramette, A. & Tiedje, J. M. (2006). Toward a more robust assessment of intraspecies diversity, using fewer genetic markers. *Appl Environ Microbiol* **72**, 7286–7293.
- Martens, M., Delaere, M., Coopman, R., De Vos, P., Gillis, M. & Willems, A. (2007). Multilocus sequence analysis of *Ensifer* and related taxa. *Int J Syst Evol Microbiol* **57**, 489–503.
- Martínez-Romero, E. (2003). Diversity of *Rhizobium-Phaseolus vulgaris* symbiosis: overview and perspectives. *Plant Soil* **252**, 11–23.
- Martínez-Romero, E., Segovia, L., Mercante, F. M., Franco, A. A., Graham, P. & Pardo, M. A. (1991). *Rhizobium tropici*, a novel species nodulating *Phaseolus vulgaris* L. beans and *Leucaena* sp. trees. *Int J Syst Bacteriol* **41**, 417–426.
- Menna, P., Barcellos, F. G. & Hungria, M. (2009). Phylogeny and taxonomy of a diverse collection of *Bradyrhizobium* strains based on multilocus sequence analysis of the 16S rRNA gene, ITS region and *glnII*, *recA*, *atpD* and *dnaK* genes. *Int J Syst Evol Microbiol* **59**, 2934–2950.
- Michiels, J., Dombrecht, B., Vermeiren, N., Xi, C., Luyten, E. & Vanderleyden, J. (1998). *Phaseolus vulgaris* is a non-selective host for nodulation. *FEMS Microbiol Ecol* **26**, 193–205.
- Ormeño-Orrillo, E., Menna, P., Almeida, L. G., Ollero, F. J., Nicolás, M. F., Pains Rodrigues, E., Shigeyoshi Nakatani, A., Silva Batista, J. S., Oliveira Chueire, L. M. & other authors (2012). Genomic basis of broad host range and environmental adaptability of *Rhizobium tropici* CIAT 899 and *Rhizobium* sp. PRF 81 which are used in inoculants for common bean (*Phaseolus vulgaris* L.). *BMC Genomics* **13**, 735.
- Pinto, F. G. S., Hungria, M. & Mercante, F. M. (2007). Polyphasic characterization of Brazilian *Rhizobium tropici* strains effective in fixing N₂ with common bean (*Phaseolus vulgaris* L.). *Soil Biol Biochem* **39**, 1851–1864.
- Ribeiro, R. A., Barcellos, F. G., Thompson, F. L. & Hungria, M. (2009). Multilocus sequence analysis of Brazilian *Rhizobium* microsymbionts of common bean (*Phaseolus vulgaris* L.) reveals unexpected taxonomic diversity. *Res Microbiol* **160**, 297–306.
- Ribeiro, R. A., Rogel, M. A., López-López, A., Ormeño-Orrillo, E., Barcellos, F. G., Martínez, J., Thompson, F. L., Martínez-Romero, E. & Hungria, M. (2012). Reclassification of *Rhizobium tropici* type A strains as *Rhizobium leucaenae* sp. nov. *Int J Syst Evol Microbiol* **62**, 1179–1184.
- Rincón-Rosales, R., Villalobos-Escobedo, J. M., Rogel, M. A., Martínez, J., Ormeño-Orrillo, E. & Martínez-Romero, E. (2013). *Rhizobium calliandrae* sp. nov., *Rhizobium mayense* sp. nov. and *Rhizobium jaguaris* sp. nov., rhizobial species nodulating the medicinal legume *Calliandra grandiflora*. *Int J Syst Evol Microbiol* **63**, 3423–3429.
- Saitou, N. & Nei, M. (1987). The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* **4**, 406–425.
- Stocco, P., Santos, J. C. P., Vargas, V. P. & Hungria, M. (2008). Avaliação da biodiversidade de rizóbios simbiotes do feijoeiro (*Phaseolus vulgaris* L.) em Santa Catarina. *Rev Bras Cienc Solo* **32**, 1107–1120 (in Portuguese).
- Tamura, K. & Nei, M. (1993). Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Mol Biol Evol* **10**, 512–526.
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M. & Kumar, S. (2011). MEGA5: molecular evolutionary genetics analysis using

maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol* **28**, 2731–2739.

Tighe, S. W., de Lajudie, P., Dipietro, K., Lindström, K., Nick, G. & Jarvis, B. D. W. (2000). Analysis of cellular fatty acids and phenotypic relationships of *Agrobacterium*, *Bradyrhizobium*, *Mesorhizobium*, *Rhizobium* and *Sinorhizobium* species using the Sherlock Microbial Identification System. *Int J Syst Evol Microbiol* **50**, 787–801.

Valverde, A., Igual, J. M., Peix, A., Cervantes, E. & Velázquez, E. (2006). *Rhizobium lusitanum* sp. nov. a bacterium that nodulates *Phaseolus vulgaris*. *Int J Syst Evol Microbiol* **56**, 2631–2637.

Velázquez, E., Palomo, J. L., Rivas, R., Guerra, H., Peix, A., Trujillo, M. E., García-Benavides, P., Mateos, P. F., Wabiko, H. & Martínez-Molina, E. (2010). Analysis of core genes supports the reclassification of strains *Agrobacterium radiobacter* K84 and *Agrobacterium tumefaciens* AKE10 into the species *Rhizobium rhizogenes*. *Syst Appl Microbiol* **33**, 247–251.

Wang, E. T. & Martínez-Romero, E. (2000). Phylogeny of root- and stem-nodule bacteria associated with legumes. In *Prokaryotic Nitrogen Fixation: A Model System for the Analysis of a Biological Process*, pp. 177–186. Edited by E. W. Triplett. Wymondham, UK: Horizon Scientific Press.

Table S1. Percentage nucleotide identity between species of the “*R. tropici* group” and between these species and *R. freirei* sp. nov. at five loci.

Species	Loci					
	16S rRNA	IGS	<i>recA</i>	<i>atpD</i>	<i>rpoB</i>	concatenat e [‡]
	98.3-	-	88.9-	88.7-97.7	90.3-96.4	85.4-96
Between species range [†]	99.9		95.2			
Between <i>R. freirei</i> and						
<i>R. multihospitium</i>	1	76.4	92.3	96.7	96.1	95.2
<i>R. miluonense</i>	99.8	-	93.8	96.5	95.3	95.2
<i>R. hainanense</i>	99.8	77.5	95.2	96.5	95	95.5
<i>R. tropici</i>	99.5	74.9	92.7	96.7	95.2	94.9
<i>R. leucaenae</i>	99.5	76	91.4	94.7	93.2	93.1
<i>R. lusitanum</i>	99.4	64.3	92.3	92.6	94.1	93.1
<i>R. rhizogenes</i>	99.3	64.2	90.4	93.1	92.1	91.9
<i>R. calliandrae</i>	99.1	-	89.8	89	91.5	90.3
<i>R. mayense</i>	99	-	89.1	91.9	91.5	90.9
<i>R. jaguaris</i>	98.8	-	88.6	89.2	92.6	90.5

* Sequence alignments length were 1308, 442, 437, 650, 450 and 582 nt for 16S rRNA, *recA*, *atpD*, *rpoB*, *glnII* and *gyrB*, respectively.

[†] Without *R. freirei*.

[‡] concatenate of *recA*, *atpD* and *rpoB* gene sequences.

Table S2. DNA-DNA hybridization* between *R. freirei* PRF 81^T and related species.

Strain	DDH (%)
<i>R. freirei</i> PRF 81 ^T	100
<i>R. tropici</i> CIAT 899 ^T	27
<i>R. leucaenae</i> CFN 299 ^T	28
<i>R. miluonense</i> CCBAU 41251 ^T	50
<i>R. lusitanum</i> P1-7 ^T	31
<i>R. hainanense</i> I66 ^T	8
<i>R. rhizogenes</i> IAM13570 ^T	36
<i>R. mayense</i> CCGE526 ^T	14
<i>R. multihospitium</i> CCBAU83401 ^T	23
<i>R. jaguaris</i> CCGE525 ^T	23
<i>R. calliandrae</i> CCGE524 ^T	11
<i>R. etli</i> CFN 42 ^T	10

* The standard error of the mean between replicate experiments was 4% for the DDH involving *R. miluonense* and in the range of 2-13% with the remaining strains.

Table S3. Distinctive phenotypic features of *Rhizobium freirei* sp. nov. and the predominant symbionts of common bean.

	<i>R. freirei</i> PRF 81 ^T	<i>R. leucaenae</i> CFN 299 ^T	<i>R. tropici</i> CIAT 899 ^T	<i>R. etli</i> CFN 42 ^T	<i>R. leguminosarum</i> USDA 2671	<i>R. gallicum</i> R602 ^T	<i>R. giardinii</i> H152 ^T
Growth characteristics							
Colony morphology in YM medium	opaque, gummy	opaque, dry	translucent, gummy	translucent, gummy	translucent, gummy	semi-translucent, gummy	translucent, wet
Color of colony in YMA medium	rose	rose	white	white	white	white	white
Colony elevation	convex	flat	convex	convex	convex	convex	convex
LB ^b	w	–	+	–	–	–	–
PY - Ca ^b	w	–	+	–	–	–	–
37°C ^b	+	+	+	–	–	NA	NA
40°C ^b	+	w	+	–	–	–	–
pH 4.0 ^b	+	w	+	–	–	–	–
Synthesis of melanin ^b	w	–	+	–	–	NA	NA
Carbohydrate source							
Glycerol	+	+	+	+	+	–	–
D-Arabinose	w	+	w	+	+	–	–
D-Xylose	w	–	–	–	w	–	–
D-Adonitol	+	w	–	+	–	–	–
D-Galactose	+	+	+	+	+	–	–
D-Glucose	+	+	w	w	+	w	w
D-fructose	w	+	–	+	+	–	–
D-Mannose	+	+	–	w	+	–	–
L-Rhamnose	w	w	–	–	+	w	–
D-Mannitol	w	w	w	w	+	w	w
D-Sorbitol	w	w	+	–	+	–	–
Metil- α -D-Glucopyranoside	w	w	–	–	–	+	–
Salicin	+	w	w	–	–	+	–
D-Celiobiose	w	+	–	–	w	–	–
D-Lactose	w	–	–	w	w	–	+
D-Rafinose	w	w	w	–	–	+	+
Xylitol	w	+	+	w	–	–	–
D-Tagatose	w	w	+	+	–	+	–
L-fucose	+	+	–	–	w	w	–
D-Arabitol	w	+	–	w	–	–	–
L-Arabitol	+	+	–	–	+	–	–

^b From Hungria *et al.* (2000) and Pinto *et al.* (2007); ^T Type strains ; NA, Not avalied

Table S4. Fatty acid profile of *Rhizobium freirei* PRF 81^T and type strains of three related species grown for 5 d at 28°C on yeast extract-mannitol agar plates.

Data obtained with the MIDI system using FAME library TSBA6.

Fatty acid	<i>R. freirei</i> PRF 81 ^T	<i>R. tropici</i> CIAT 899 ^T	<i>R. multihospitum</i> CCBAU 83401 ^T	<i>R. hainanense</i> I66 ^T
iso-C15:0 3-OH	3.07	4.03	3.86	
C16:0	5.26	5.37	4.49	15.99
C16:0 3-OH	4.13	4.39	5.39	
iso-C17:0		0.61	0.68	
C17:0 3-OH			0.55	
C18:0	3.92	3.67	1.58	
C18:0 3-OH	2.26	2.33	2.39	
C18:1 2-OH	3.95	1.15	2.81	
C18:1 ω7c11-methyl		0.98		
C19:0 cyclo ω8c	9.30	30.07	23.18	13.12
C20:2 ω6,9c		0.72		
Summed Features				
2 [*]	3.59	3.13	3.99	7.77
3 [†]				
8 [‡]	64.52	43.11	50.64	63.13

* Summed Feature 2=C12:0 aldehyde, unknown 10.928, iso-C16:1 I/C14:0 3-OH, C14:0 3-OH/iso-C16:1 I.

† Summed Feature 3 = C_{16:1} ω7c/_{16:1} ω6c.

‡ Summed Feature 8=C_{18:1}ω7c/ω9t/ω12t, C_{18:1}ω7c/ω9c/ω12t.




Fig. S1. Neighbour joining phylogeny of 16S rRNA gene sequences of *R. freirei* sp. nov. and other *Rhizobium* species. GenBank accession numbers are given within parenthesis. Bootstrap support values based on 500 resamplings are shown at nodes only when they were $\geq 60\%$. Nodes marked with the ● symbol also had support values $\geq 60\%$ in a separate maximum likelihood phylogeny. Bar, 0.005 substitutions per nucleotide position.




Fig. S2. Dendrogram based on cluster analysis of BOX-A1R-PCR products using the unweighted pair-group method with arithmetic mean (UPGMA) algorithm and the Jaccard coefficient of *R. freirei* sp. nov. and strains belonging to the “*R. tropici* group”. Clusters were obtained with Bionumerics (Applied Mathematics, Kortrijk, Bélgica, v. 4.6).