

# Dynamic communication between plants and rhizobacteria via volatile signals

Choong-Min Ryu<sup>1</sup>, Hwe-Su Yi<sup>1</sup>, Yeo-Rim Ahn<sup>2</sup>, Won-il Kim<sup>3</sup>, Huiming Zhang<sup>4</sup>, Seung-Hwan Park<sup>1</sup>, Chang Seuk Park<sup>3</sup>, Mohamed A. Farag<sup>4</sup>, Paul W. Paré<sup>4</sup>, and Joseph W. Kloepper<sup>5</sup>

<sup>1</sup>Systems Microbiology Research Center, KRIBB, Daejeon, S. Korea; <sup>2</sup>Department of Biological Science, KAIST, Daejeon, S. Korea; <sup>3</sup>Department of Agricultural Biology and Environmental Science, Gyeongsang National University, Jinju, S. Korea; <sup>4</sup>Department of Chemistry and Biochemistry, Texas Tech University, Lubbock, TX USA; <sup>5</sup>Department of Entomology and Plant Pathology, Auburn University, Auburn, AL USA

cmryu@kribb.re.kr

## Abstract

Certain plant growth-promoting rhizobacteria (PGPR), in the absence of physical contact with plants, stimulate plant growth and elicit induced systemic resistance (ISR) via volatile organic compound (VOC) emissions. Gas chromatographic analysis of VOCs collected from the PGPR strains *Bacillus subtilis* strain GB03 and *B. amyloliquefaciens* strain IN937a reveals consistent patterns in VOC emissions in comparison with non-growth promoting strains such as *E. coli* DH5 $\alpha$ . The two most abundant compounds, 2,3-butanediol and 3-hydroxy-2-butanone, are consistently emitted from GB03 and IN937a while these metabolites are not released from DH5 $\alpha$ . Transcriptional approaches have been employed to prove how *Arabidopsis* respond to biologically active bacterial VOCs. To assess potential utilization of PGPR VOCs for crop plants, volatile blends from GB03, IN937a, and DH5 $\alpha$  have been applied separately to pepper, tomato, and cucumber roots. Bacterial survival capacity of 2,3-butanediol null mutants was significantly reduced in proximity with plant tissue. These reduced bacterial survival rates suggest that in addition to bacterial VOCs triggering plant growth and induced systemic resistance in plants, such chemicals provide protection for

PGPR via chemical signaling within the host plant.

### **Rhizobacteria help plant growth and defense against multiple pathogens**

Plant growth-promoting rhizobacteria (PGPR) include a wide range of root-colonizing bacteria with the capacity to enhance plant growth by increasing seed emergence, plant weight, and crop yields (Kloepper, 1980). Seed or seedling treatments with PGPR have been used to enhance growth of several crops (Glick, 1995) as well as to suppress the growth of plant pathogens and deleterious rhizosphere microorganisms. Proposed mechanisms for plant growth promotion by PGPR include bacterial synthesis of the plant hormones indole-3-acetic acid (IAA), cytokinin, and gibberellin; breakdown of plant-produced ethylene by bacterial production of 1-aminocyclopropane-1-carboxylate (ACC) deaminase; and increased mineral and nitrogen availability (Glick et al., 1995; 1999).

In addition to promoting plant growth, application of certain PGPR strains to seeds or seedlings leads to a state of induced systemic resistance (ISR) in the treated plant (Kloepper et al., 1992). PGPR can influence pathogens by direct or indirect effects. Indirectly, root-colonization by certain rhizobacteria induces systemic resistance that is effective against viral, bacterial, and fungal pathogens, as well as nematodes (Kloepper et al., 1999). ISR occurs in many plants including carnation, cucumber, tobacco, tomato, bean, radish, and *Arabidopsis* (Van Loon et al., 1998; Ryu et al., 2003b; 2004b). In contrast to signaling pathways of necrotic pathogens or chemical-elicited systemic acquired resistance, ISR does not require a salicylic acid-dependent pathway (Van Loon et al., 1998; Ryu et al., 2003b; 2004b). Several microbial determinants have been associated with elicitation of ISR: 2,4-diacetylphloroglucinol (Iavicoli et al., 2003), the O-antigen from lipopolysaccharide (Van Peer and Schippers, 1992), and salicylic acid (Maurhofer et al., 1998).

### **Bacterial volatiles are as aromatic stimulants for plant growth and defense**

By physically separating plant growth promotion rhizobacteria from their host plant airborne chemicals from certain soil bacteria have been identified as effective signals for triggering plant growth and ISR. (Ryu *et al.*, 2003a; Ryu *et al.*, 2004a). Of several PGPR tested, two strains, *Bacillus subtilis* GB03 and *B. amyloliquefaciens* IN937a, elicited plant growth promotion and ISR

by volatile organic compounds (VOCs) emissions. In the case of growth promotion by VOCs from strain GB03, the cytokinin receptor-deficient (*cre1*) and cytokinin- and ethylene-insensitive (*ein2*) mutants were insensitive to growth promotion effects triggered by GB03 volatiles, while ethylene-insensitive (*etr1*), auxin-transporter-deficient and ethylene-insensitive (*eir1*), gibberellic acid-insensitive (*gai2*), and brassinosteroid-insensitive (*cbb1*) mutants were all sensitive to GB03 VOCs as evidenced by measurement of growth promotion.

Of several *Arabidopsis* mutant lines tested for regulatory control of ISR against *Erwinia carotovora* subsp. *carotovora*, only the ethylene-insensitive line (*ein2*) did not exhibit an amelioration of disease symptoms when *Arabidopsis* plants were pre-treated with GB03 volatiles. To examine if ISR is mediated at the level of transcription, three transgenic  $\beta$ -glucuronidase (GUS) fusion lines involved in plant-defense signaling were tested. Of those lines assayed, the jasmonic acid and ethylene-responsive PDF 1.2 gene GUS fusion plant alone exhibited elevated GUS activity (>1000 fold) over untreated control plants (Ryu *et al.*, 2004).

Besides study of strain GB03 VOCs, characterization of the VOCs produced by IN937a, coupled with bioassays of fractions of VOCs elicited plant growth promotion allowed for an initial characterization of potentially active bacterial volatiles. Exposure of *Arabidopsis* to VOCs from strain IN937a resulted in a significant reduction in disease severity caused by *E. carotovora* subsp. *carotovora*. Elicitation of ISR occurred with exposure to bacterial VOCs for as little as 4 days. Various mutant lines of *Arabidopsis* were exposed to whole VOCs. Mutant lines included a jasmonic acid-insensitive line (*coi1*), an ethylene-insensitive line (*ein2*), a salicylic acid-degrading line (NahG), and a line that lacks a regulatory gene (*npr1*). VOCs from strain IN937a elicited ISR on all of these lines. Hence, elicitation of ISR by VOCs of IN937a is independent of jasmonic acid, ethylene, salicylic acid, and *npr1*. Such a pattern of signal pathway has not been reported with ISR elicited by bacteria, and therefore, it is likely that VOCs of IN937a elicit a distinct and uncharacterized pathway in *Arabidopsis*. The two most abundant compounds, 2,3-butanediol and 3-hydroxy-2-butanone (also referred to as acetoin), were consistently released from strains GB03 and IN937a while these metabolites were not released from DH5 $\alpha$  or water-treated MS media. Other components of the complex bacterial bouquet that did not elicit ISR included dodecane, 2-undecanone, 2-tridecanone, 2-tridecanol, and tetramethyl pyrazine.

In *Bacillus* sp., 2,3-butanediol and acetoin are produced with low atmospheric O<sub>2</sub> partial pressure to provide an alternative electron sink for the

regeneration of  $\text{NAD}^+$  when aerobic respiration is limited possible. This additional metabolic pathway functions analogously to alcohol fermentation activated in yeast under anaerobic conditions. The biological activity of 2,3-butanediol in triggering ISR was surmised in *Arabidopsis* when pre-exposure of plants to low doses (pg to ng range) of 2,3-butanediol activated ISR. The priming activity of 2,3-butanediol to reduce a plant's susceptibility to disease was confirmed when Bacilli strains genetically blocked in the production of 2,3-butanediol exhibited no disease protection. More interestingly, direct application of bacterial volatiles into plant roots increased plant growth significantly and elicited ISR in the greenhouse (unpublished data).

### **Uncharacterized bacterial odors**

The involvement of known signaling pathways in *Arabidopsis* were screened by exposing defined mutants and transgenic plant lines to bacterial emissions containing 2,3-butanediol. ISR triggered by strain GB03 VOC emissions was independent of salicylic acid, NPR1, and jasmonic acid signaling pathways, but did appear to be mediated by ethylene. Interestingly plant growth promotion activation by strain IN937a was independent of all the signaling pathways that were tested suggesting that additional VOCs utilize alternative pathways to trigger plant growth promotion (Ryu et al., 2003a). In fact, solid phase microextraction (SPME) analysis of bacterial odors turned up substantial differences in VOC profiles between two active strains GB03 and IN937a for growth promotion. Major branched-chain alcohols 3-methyl-1-butanol, 2-methyl-1-butanol and their conjugates were detected in IN937a, albeit not in GB03. Differences in volatile profiles might account for the dissimilarity observed among both strains in triggering growth promotion via volatile chemical signals, and suggests the existence of diverse VOC metabolism existing in two closely related Bacilli species (Farag et al., 2006).

### **Plant and bacteria responses to rhizobacterial volatiles**

To understand plant-signaling pathways induced by bacterial VOCs during plant growth promotion and ISR, we have begun to characterize global changes in the *Arabidopsis* transcriptome using available microarray technology. Microarray results revealed physiological changes associated with growth, photosynthesis, and stress tolerance. In addition to bacterial synthesis of plant growth regulators such as auxin, bacterial VOCs induce auxin synthesis and transport *in planta*. Transcriptional analysis also

identified changes in cell wall loosening providing a possible mechanism for rapid cell expansion and leaf growth promotion that is observed in GB03 VOC exposure (Zhang et al 2007). 2,3-Butanediol also appears to participate in bacterial colonization with its host. In case of *Vibrio cholerae* the mutant defective in 2,3-butanediol synthesis exhibited reduced capacity to colonize the gut of its host the mouse and did not develop its usual biofilm. Preliminary observations with the 2,3-butanediol non-producing *Bacillus subtilis* strain also indicates poor colonization of its host, in this case, pepper roots and overall poor bacterial fitness *in situ*. It appears that 2,3-butanediol produced by *B. subtilis* may serve in a dual functions eliciting the production of plant antimicrobial compounds and as a protecting agent for bacterial cells (Yi et al., unpublished data). Additional investigations are underway to better characterize how PGPR utilize volatile signaling to transmit information between plants and rhizobacteria.

## Acknowledgements

This work was supported by Biogreen21 (20070401034005), the Welch Foundation (Grant D1478) and Technology Development Program for Agriculture and Forestry, Ministry of Agriculture and Forestry, S. Korea.

## Literature cited

- Cleyet-Marcel, J.C., Larcher, M., Bertrand, H., Rapior, S., and Pinochet, X. 2001. Plant growth enhancement by rhizobacteria. In: Nitrogen assimilation by plants, physiological, biochemical and molecular aspects. ed. Morot-Gaudry, J-F. Science Publishers, Inc. Enfield, NH, USA.
- Iavicoli, A., Boutet, E., Buchala, A., and Metraux, J.P. 2003. Induced systemic resistance in *Arabidopsis thaliana* in response to root inoculation with *Pseudomonas fluorescens* CHA0. *Mol. Plant-Microbe Interact.* 16:851-858.
- Farag, M.A., Ryu, C.M., Sumner, L.W., and Paré P.W. 2006. Profiling of rhizobacterial emissions reveals prospective inducers of growth promotion and induced systemic resistance in plants. *Phytochemistry* 67:2262-2268.
- Glick, B.R. 1995. The enhancement of plant growth by free-living bacteria. *Can. J. Microbiol.* 41:109-117.
- Glick, B.R. 1999. Biochemical and Genetic Mechanisms Used by Plant Growth Promoting Bacteria, (eds). Glick BR, Patten CN, Holguin G, Penrose DM, Imperial College Press, London, UK.
- Kloepper, J.W., Leong, J., Teintze, M., and Schroth, M.N. 1980. Enhanced plant growth by siderophores produced by plant growth-promoting rhizobacteria. *Nature* 286:885-886.
- Kloepper, J.W., Tuzun, S., and Kuc, J. 1992. Proposed definitions related to induced disease resistance. *Biocontrol Sci. Technol.* 2:349-351.

- Kloepper, J.W., Ryu, C.M., and Zhang, S. 2004. Induced systemic resistance and promotion of plant growth by *Bacillus* spp. *Phytopathology* 94:1259-1266.
- Kloepper, J.W., and Ryu, C.M. 2006. Bacterial endophytes as elicitors of induced systemic resistance. In: *Soil Biology, Vol.9 Microbial Root Endophytes*. B. Schulz, C. Boyle, and T.N. Sieber (eds.). Berlin Heidelberg: Springer-Verlag, Germany.
- Kloepper, J.W., Rodriguez-Kabana, R., Zehnder, G.W., Murphy, J., Sikora, E., and Fernandez, C. 1999. Plant root-bacterial interactions in biological control of soilborne diseases and potential extension to systemic and foliar diseases. *Austral. Plant. Pathol.* 28:27-33.
- Maurhofer, M., Reimmann, C., Schmidli-Sacherer, P., Heeb, S., Haas, D., and Defago, G. 1998. Salicylic acid biosynthetic genes expressed in *Pseudomonas fluorescens* strain P3 improve the induction of systemic resistance in tobacco against tobacco necrosis virus. *Phytopathology* 88:678-684.
- Ryu, C.M., Farag, M.A., Hu, C.H., Reddy, M.S., Wei, H.X., Pare, P.W., and Kloepper, J.W. 2003a. Bacterial volatiles promote growth in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA.* 100:4927-4932.
- Ryu, C.M., Hu, C.H., Reddy, M.S., and Kloepper, J.W. 2003b. Different signaling pathways of induced resistance by rhizobacteria in *Arabidopsis thaliana* against two pathovars of *Pseudomonas syringae*. *New Phytol.* 160:413-420.
- Ryu, C.M., Farag, M.A., Hu, C.H., Reddy, M.S., Kloepper, J.W., and Pare, P.W. 2004a. Bacterial volatiles induce systemic resistance in *Arabidopsis*. *Plant Physiol.* 134:1017-1026.
- Ryu, C.M., Murphy, J.F., Mysore, K.S., and Kloepper, J.W. 2004b. Plant growth-promoting rhizobacteria systemically protect *Arabidopsis thaliana* against Cucumber mosaic virus by a salicylic acid and NPR1-independent and jasmonic acid-dependent signaling pathway. *Plant J.* 39:381-392.
- Van Loon, L.C., Bakker, P.A.H.M., and Pieterse, C.M.J. 1998. Systemic resistance induced by rhizosphere bacteria. *Ann. Rev. Phytopath.* 36:453-483.
- Van Peer, R., and Schippers, B. 1992. Lipopolysaccharides of plant-growth promoting *Pseudomonas* sp. strain WCS417r induce resistance in carnation to fusarium wilt. *Neth. J. Plant Pathol.* 98:129-139.
- Yoon, S.S., and Mekalanos, J.J. 2006. 2,3-Butanediol synthesis and the emergence of the *Vibrio cholerae* El Tor biotype. *Infect. Immun.* 74:6547-6556.
- Zhang, H., Kim, M.S., Krishnamachari, V., Payton, P., Sun, Y., Grimson, M., Farag, M.A., Ryu, C.M., Allen, R., Melo, I.S., and Pare, P.W. 2007. Rhizobacterial volatile emissions regulate auxin homeostasis and cell expansion in *Arabidopsis*. *Planta* 226:839-51.

# Biology of Plant-Microbe Interactions, Volume 6

Edited by

Matteo Lorito  
Sheridan Lois Woo  
Felice Scala

Università Degli Studi Di Napoli  
Portici, Italy

Proceedings of the 13th International Congress  
on Molecular Plant-Microbe Interactions  
Sorrento (Naples), Italy  
July 21–27, 2007

Published by the  
International Society for Molecular Plant-Microbe Interactions  
St. Paul, Minnesota, USA

Articles may be referenced as follows:

[Authors' names]. 2008. [Paper title]. Paper [number] in: *Biology of Plant-Microbe Interactions*, Volume 6. M. Lorito, S. L. Woo, and F. Scala, eds. International Society for Molecular Plant-Microbe Interactions, St. Paul, MN.

Cover photo by Gianfranco Capodilupo; logo design by Antonino Balsamo

This CD-Rom has been reproduced directly from edited, computer-generated copy submitted by the authors to the editors of this volume. The editors have verified text formatting, references, and quality of text and figures. No editing or proofreading has been done by the publisher.

Reference in this publication to a trademark, proprietary product, or company name by personnel of the U.S. Department of Agriculture or anyone else is intended for explicit description only and does not imply approval or recommendation to the exclusion of others that may be suitable.

ISBN: 978-0-9654625-5-6

© 2008 by the International Society for Molecular Plant-Microbe Interactions

All rights reserved.

No part of this book may be reproduced in any form, including photocopy, microfilm, information storage and retrieval system, computer database or software, or by any other means, including electronic or mechanical, without written permission from the publisher.

Copyright is not claimed in any portion of this work written by U.S. government employees as part of their official duties.

International Society for Molecular Plant-Microbe Interactions  
3340 Pilot Knob Road  
St. Paul, Minnesota 55121 U.S.A.