

Polyphasic evidence supporting the reclassification of *Bradyrhizobium japonicum* group Ia strains as *Bradyrhizobium diazoefficiens* sp. nov.

Jakeline Renata Marçon Delamuta,^{1,2} Renan Augusto Ribeiro,^{1,2}
Ernesto Ormeño-Orrillo,³ Itamar Soares Melo,⁴
Esperanza Martínez-Romero³ and Mariangela Hungria^{1,2}

Correspondence

Mariangela Hungria
mariangela.hungria@embrapa.br
or biotecnologia.solo@hotmail.com

¹Embrapa Soja, C.P. 231, 86001-970 Londrina, Paraná, Brazil

²Universidade Estadual de Londrina, Dept of Microbiology, C.P. 60001, 86051-990 Londrina, Paraná, Brazil

³Centro de Ciencias Genómicas, Universidad Nacional Autónoma de México, Cuernavaca, Morelos, Mexico

⁴Embrapa Meio Ambiente, C.P. 69, 13820-000 Jaguariúna, São Paulo, Brazil

Bradyrhizobium japonicum was described from soybean root-nodule bacterial isolates. Since its description, several studies have revealed heterogeneities among rhizobia assigned to this species. Strains assigned to *B. japonicum* group Ia have been isolated in several countries, and many of them are outstanding soybean symbionts used in inoculants worldwide, but they have also been isolated from other legume hosts. Here, we summarize published studies that indicate that group Ia strains are different from the *B. japonicum* type strain USDA 6^T and closely related strains, and present new morphophysiological, genotypic and genomic evidence to support their reclassification into a novel species, for which the name *Bradyrhizobium diazoefficiens* sp. nov. is proposed. The type strain of the novel species is the well-studied strain USDA 110^T (=IAM 13628^T = CCRC 13528^T = NRRL B-4361^T = NRRL B-4450^T = TAL 102^T = BCRC 13528^T = JCM 10833^T = TISTR 339^T = SEMIA 5032^T = 3I1B110^T = ACCC 15034^T = CCT 4249^T = NBRC 14792^T = R-12974^T = CNPSo 46^T).

Biological fixation is the main source of nitrogen for natural and agricultural ecosystems. In agriculture, the symbioses of nitrogen-fixing bacteria, collectively known as rhizobia, with crops belonging to the family Leguminosae (=Fabaceae) are the most studied. Relatively high contributions to nitrogen nutrition have been demonstrated in pulses, fodders, green manures and trees (Ormeño-Orrillo *et al.*, 2013). Members of the genus *Bradyrhizobium* constitute an important group of rhizobia, some of which form symbioses with economically important crops, such as soybean [*Glycine max* (L.) Merr.].

A first systematic classification of rhizobia was proposed by Fred *et al.* (1932), based on the cross-inoculation concept with respect to the host legume, resulting in six species of

Rhizobium: *Rhizobium meliloti*, *R. trifolii*, *R. phaseoli*, *R. lupini*, *R. leguminosarum* and *R. japonicum*, the last of which is a symbiont of soybean. Fifty years later, the taxonomy was redefined using numerical criteria, including several morphophysiological and genetic properties. A new genus was created, *Bradyrhizobium*, to accommodate rhizobia with the typical property of slow growth rates in culture media, with only one defined species, *Bradyrhizobium japonicum* (Jordan, 1982). It is noteworthy that the division of rhizobia into fast and slow growers *in vitro* had been proposed earlier by Löhnis & Hansen (1921).

It has been suggested that *Bradyrhizobium* is the ancestor of the alpha-rhizobia, probably originating in the tropics (Lloret & Martínez-Romero, 2005; Norris, 1965; Provorov & Vorob'ev, 2000). However, despite its evolutionary position, significantly fewer species have been described in *Bradyrhizobium* in comparison with *Rhizobium*. One possible explanation is that the 'backbone' of modern taxonomy relies on the analysis of 16S rRNA gene sequences (Woese, 1987), which are highly conserved in

Abbreviations: DDH, DNA–DNA hybridization; ITS, intergenic transcribed spacer; MLSA, multilocus sequencing analysis.

The GenBank/EMBL/DDBJ accession numbers for the sequences determined in this study are detailed in Table S2.

Four supplementary figures and five supplementary tables are available with the online version of this paper.

Bradyrhizobium (Germano *et al.*, 2006; Menna *et al.*, 2006; So *et al.*, 1994; Urtz & Elkan, 1996; van Berkum & Fuhrmann, 2000; Vinuesa *et al.*, 1998; Willems *et al.*, 2001b). However, high diversity in a variety of morpho-physiological and genetic properties within *Bradyrhizobium* has been reported (Boddey & Hungria, 1997; Minamisawa, 1989; Minamisawa *et al.*, 1998; Tian *et al.*, 2012; Urtz & Elkan, 1996; van Berkum & Fuhrmann, 2001; Vinuesa *et al.*, 1998), and better resolution of species-level clades has been obtained by the analysis of other housekeeping genes using the MLSA (multilocus sequencing analysis) approach (e.g. Delamuta *et al.*, 2012; Menna *et al.*, 2009).

Since the subdivision of strains classified as *B. japonicum* in 1992, with the creation of *Bradyrhizobium elkanii* (Kuykendall *et al.*, 1992), no other change has been proposed for *B. japonicum*, despite several reports of subgroups within this species. These reports started in 1980 when DNA–DNA hybridization (DDH) analyses defined two relatedness groups for *B. japonicum* strains (Hollis *et al.*, 1981), I and Ia, later confirmed by Kuykendall *et al.* (1992). Urtz & Elkan (1996) also reported low DDH between the type strain USDA 6^T and the broadly studied group Ia strain USDA 110, proposed in our study as the type strain for a novel species. Other reports of heterogeneity within *B. japonicum* included differences in extracellular polysaccharide composition (Huber *et al.*, 1984), fatty acid profiles (fatty acid methyl esters) (Graham *et al.*, 1995; van Berkum & Fuhrmann, 2001), ribosomal and housekeeping genes (Delamuta *et al.*, 2012; Germano *et al.*, 2006; Menna *et al.*, 2006; van Berkum & Fuhrmann, 2000; Vinuesa *et al.*, 2008) and comparative genomics (Tian *et al.*, 2012), among others.

For the last two decades, our group has reported morpho-physiological, genetic and symbiotic diversity within *B. japonicum* (Boddey & Hungria, 1997; Delamuta *et al.*, 2012; Ferreira & Hungria, 2002; Germano *et al.*, 2006; Menna & Hungria, 2011; Menna *et al.*, 2006; Santos *et al.*, 1999), including a proposition, based on MLSA, that strains in group Ia might represent a novel species (Menna *et al.*, 2009).

In this study, we summarize differences reported by our group and other laboratories and, with additional analyses, we present new evidence to suggest that *B. japonicum* group Ia represents a novel species distinct from group I strains.

Four *Bradyrhizobium* strains used in this study, SEMIA 6059, SEMIA 5060, CPAC 7 and USDA 110 (Table 1), were chosen from a previous MLSA study by our group based on a high level of genetic similarity (Menna *et al.*, 2009). SEMIA 6059 was isolated from winged bean (*Psophocarpus tetragonolobus*), and the others from soybean growing in the USA, Japan and Brazil; other *Bradyrhizobium* strains used in this study are listed in Table 1. We always include comparisons with two strains well studied by our group because of their broad use in commercial inoculants in Brazil: group Ia strain CPAC 7, a natural variant of CB 1809 (Santos *et al.*, 1999), and group I strain CPAC 15, a

natural variant of SEMIA 566 (Mendes *et al.*, 2004). CPAC 15 and SEMIA 566 belong to the same highly competitive serogroup as USDA 123 (Mendes *et al.*, 2004). All strains from this study are deposited at the Culture Collection of Diazotrophic and Plant Growth Promoting Bacteria of Embrapa Soja (Londrina, Brazil) and at the Center for Genomic Sciences Culture Collection (Cuernavaca, Mexico).

The rhizobia were grown on yeast extract-mannitol agar (YMA) at 28 °C. Stocks were prepared on YMA and kept for long-term storage at –80 and –150 °C (in 30 % glycerol) and lyophilized, and at 4 °C as source cultures.

Genomic DNA extraction and fingerprints with the BOX-A1R primer were performed as described by Kaschuk *et al.* (2006). Cluster analyses, performed with the Bionumerics 4.6 software using the UPGMA algorithm and Jaccard similarity coefficients, revealed that all four group Ia strains clustered at an 85 % similarity level (Fig. S1, available in IJSEM Online). This group was separated from the group I *B. japonicum* strains USDA 6^T, CPAC 15 and SEMIA 566, as well as from other species of *Bradyrhizobium*.

The 16S rRNA gene and fragments of the *atpD*, *glnII*, *recA*, *gyrB*, *rpoB* and *dnaK* genes and of the 16S–23S rRNA intergenic transcribed spacer (ITS) were amplified using primers and conditions indicated in Table S1, and were sequenced by using dideoxy termination as described before (Menna *et al.*, 2009). Accession numbers for the sequences used in our study are given in Table S2. Multiple sequence alignment, alignment editing and phylogenetic analyses were performed with MEGA5 (Tamura *et al.*, 2011). Neighbour-joining (Saitou & Nei, 1987) and maximum-likelihood (Felsenstein, 1981) phylogenetic reconstructions gave similar results; therefore, only neighbour-joining phylograms are presented. For the ITS, a neighbour-joining dendrogram was reconstructed based on a matrix of uncorrected distances (Willems *et al.*, 2001a).

In the 16S rRNA gene phylogram, group Ia strains CPAC 7, SEMIA 6059, SEMIA 5060 and USDA 110 were clustered in a well-supported clade that was closely related to *Bradyrhizobium daqingense*, *B. canariense*, *B. yuanmingense*, *B. liaoningense* and *B. japonicum* (Fig. 1). The results confirm a cluster reported previously by our group with ten strains, all but SEMIA 6059 symbionts of soybean (Menna *et al.*, 2009). Sequences of group Ia strains were 99.8–100 % identical to each other and showed 96.1–99.6 % identity to those of other *Bradyrhizobium* strains (Table 2). Recently, two exact copies of the rRNA gene cluster (*rrn*) were identified on the chromosome of the group I strain USDA 6^T (Kaneko *et al.*, 2011) whereas, in group Ia strain USDA 110, a single *rrn* copy was present (Kaneko *et al.*, 2002). The same difference was observed in the genomes of the Brazilian strains CPAC 15 (*B. japonicum*) and CPAC 7 (group Ia); therefore, this might be another differential characteristic of the two groups.

High diversity within *B. japonicum* in analyses of the ITS sequence has been reported before (van Berkum &

Table 1. Strains used in this study

Strain	Other strain nomenclature	Host species	Geographical origin	Reference or source
<i>B. diazoefficiens</i> sp. nov.				
CPAC 7	SEMIA 5080, CNPSo 6	<i>Glycine max</i>	Brazil	Menna <i>et al.</i> (2009)
SEMIA 5060	J 507, CNPSo 1064	<i>Glycine max</i>	Japan	Menna <i>et al.</i> (2009)
SEMIA 6059	USDA 3309, CNPSo 1098	<i>Psophocarpus tetragonolobus</i>	USA	Menna <i>et al.</i> (2009)
USDA 110 ^T	TAL 102 ^T , TISTR 339 ^T , SEMIA 5032 ^T , CNPSo 46 ^T	<i>Glycine max</i>	USA	Kaneko <i>et al.</i> (2002)
<i>B. japonicum</i>				
USDA 6 ^T	ATCC 10324 ^T , CCUG 27876 ^T , CIP 106093 ^T , DSM 30131 ^T , HAMB1 2314 ^T , NBRC 14783 ^T , JCM 20679 ^T , LMG 6138 ^T , NRRL B-4507 ^T , NRRL L-241 ^T , VKM B-1967 ^T , CNPSo 158 ^T	<i>Glycine max</i>	USA	Jordan (1982)
CPAC 15	SEMIA 5079, DF 24, CNPSo 7	<i>Glycine max</i>	Brazil	Menna <i>et al.</i> (2009)
SEMIA 566	BR 40, CNPSo 17	<i>Glycine max</i>	USA	Menna <i>et al.</i> (2009)
<i>B. betae</i> LMG 21987 ^T	PL7HG1 ^T , CECT 5829 ^T , CNPSo 2079 ^T	<i>Beta vulgaris</i>	Spain	Rivas <i>et al.</i> (2004)
<i>B. canariense</i> LMG 22265 ^T	BTA-1 ^T , ATCC BAA-1002 ^T , CFNE 1008 ^T , CNPSo 2078 ^T	<i>Chamaecytisus proliferus</i>	Spain	Vinuesa <i>et al.</i> (2005)
<i>B. yuanmingense</i> LMG 21827 ^T	CCBAU 10071 ^T , CFNEB 101 ^T , NBRC 100594 ^T , CNPSo 2080 ^T	<i>Lespedeza</i> spp.	China	Yao <i>et al.</i> (2002)
<i>B. cytisi</i> CTAW11 ^T	LMG 25866 ^T , CECT 7749 ^T , CNPSo 2469 ^T	<i>Cytisus villosus</i>	Morocco	Chahboune <i>et al.</i> (2011)
<i>B. huanghuaihaiense</i> CCBAU 23303 ^T	LMG 26136 ^T , CGMCC 1.10948 ^T , HAMB1 3180 ^T , CNPSo 2458 ^T	<i>Glycine max</i>	China	Zhang <i>et al.</i> (2012)
<i>B. iriomotense</i> EK05 ^T	NBRC 102520 ^T , LMG 24129 ^T , CNPSo 2470 ^T	<i>Entada koshunensis</i>	Japan	Islam <i>et al.</i> (2008)
' <i>B. rifense</i> ' CTAW71	LMG 26781, CECT 8066, CNPSo 2468	<i>Cytisus villosus</i>	Morocco	Chahboune <i>et al.</i> (2012)
<i>B. liaoningense</i> LMG 18230 ^T	2281 ^T , ATCC 700350 ^T , CIP 104858 ^T , NBRC 100396 ^T , CNPSo 2455 ^T	<i>Glycine max</i>	China	Xu <i>et al.</i> (1995)




Fig. 1. Neighbour-joining phylogeny based on 16S rRNA gene sequences showing relationships between *Bradyrhizobium diazoefficiens* sp. nov. strains and other members of the genus *Bradyrhizobium*. Bootstrap values >70% are indicated at nodes. *Rhizobium tropici* CIAT 899^T was used as an outgroup. Bar, 1 substitution per 100 nucleotide positions.

Fuhrmann, 2000; Willems *et al.*, 2001a). Accordingly, in previous studies by our group with RFLP-PCR (Germano *et al.*, 2006) and sequencing of the ITS genomic region (Menna *et al.*, 2009), differences between groups I and Ia were pointed out. Here, the dendrogram based on ITS sequences also revealed that group Ia strains were distinct from other bradyrhizobia (Fig. S2). ITS sequence similarity among group Ia strains was greater than 99%, and less than 90% with other bradyrhizobia; similarities between group I and Ia strains ranged from 83.5 to 84% (Table 2). This observation is consistent with group Ia strains belonging to the same species; Willems *et al.* (2003) have shown that strains with more than 95.5% ITS similarity usually show at least 60% DDH and, therefore, belong to the same genospecies. Interestingly, group Ia strains were also characterized by a 37-nt insertion in the ITS (beginning 156 bp after the 16S rRNA gene in USDA 110) that was absent in group I strains. Therefore, this insertion may be characteristic of group Ia strains. Finally, it is noteworthy that, in a polyphasic analysis of the RFLP-PCR fragments of the 16S and 23S rRNA genes as well as the ITS, strains of groups I and Ia were positioned in different clusters (Germano *et al.*, 2006).

The MLSA approach has been used successfully in studies of *Bradyrhizobium* for over a decade (e.g. Menna *et al.*,

Table 2. Nucleotide identity (%) within *B. diazoefficiens* sp. nov. and between *B. diazoefficiens* strains and the type strains of other *Bradyrhizobium* species

Length of the aligned regions: 16S rRNA gene, 1303 bp; *atpD*, 435 bp; *glnII*, 513 bp; *recA*, 381 bp; ITS, 891 bp. Four strains of group Ia (*B. diazoefficiens* sp. nov.) were compared: CPAC 7, SEMIA 5060, SEMIA 6059 and USDA 110^T. NA, No available sequence for comparison.

Species comparison	16S rRNA	<i>atpD</i>	<i>glnII</i>	<i>recA</i>	ITS
Within <i>B. diazoefficiens</i>	99.8–100	99.5–100	99.4–100	95.5–99.7	99.4–99.7
Between <i>B. diazoefficiens</i> and:					
<i>B. japonicum</i>	99–99.2	95.8–96.2	97.4–97.6	94.7–96.5	83.5–84
<i>B. betae</i>	99.1–99.3	96.5–96.9	96.8–97.4	93.9–97.1	87.6–87.7
<i>B. canariense</i>	99–99.1	94.4–94.9	94.9–95.1	92.9–94.4	84.6–84.8
<i>B. yuanmingense</i>	98.6–98.7	95.1–95.6	93.5–93.7	91.3–92.6	82.7–82.9
<i>B. liaoningense</i>	99.1–99.3	94.9–95.3	95.7–95.9	93.1–94.4	85–85.3
<i>B. jicamae</i>	96.3–96.4	93.5–93.9	88.8–89	88.4–89.5	76.6–76.7
<i>B. huanghuaihaiense</i>	99.1–99.3	97.6–98.1	95.7–95.9	90.2–92.9	82.6–82.7
<i>B. cytisi</i>	99–99.2	94.2–94.6	94.9–95.1	91.8–94.2	NA
<i>B. daqingense</i>	99.4–99.6	95.8–96.2	94.5–94.7	90.5–91.8	85–85.3
<i>B. iriomotense</i>	98.9–99	94.6–95.1	94.9–95.1	89.5–90.5	80.3–80.5
' <i>B. rifense</i> '	99.3–99.4	97.2–97.6	95.5–95.7	91.3–92.9	NA
<i>B. elkanii</i>	96.1–96.3	93.5–93.9	89.2–89.6	91.8–93.4	73.9–74.2
<i>B. lablabi</i>	96.5–96.6	92.8–93.2	88.4–88.6	90.2–92.3	74.7–74.9
<i>B. pachyrhizi</i>	96.3–96.4	93.2–93.7	89–89.4	90.2–92.3	75.6–75.7

2009; Moulin *et al.*, 2004; Stepkowski *et al.*, 2005; Vinuesa *et al.*, 2008). Recently, sequences of at least three housekeeping genes were used for phylogenetic analysis and taxonomic classification of bradyrhizobia (Chahboune *et al.*, 2011, 2012; Chang *et al.*, 2011; Zhang *et al.*, 2012), and our results support the use of this methodology for such purposes. Group Ia strains formed distinct and well-supported clades in single-gene phylogenies of *atpD*, *recA* and *glnII*, for which sequences of all described *Bradyrhizobium* species are available (data not shown). They were closely related to *Bradyrhizobium betae* LMG 21987^T with *atpD* and *glnII* genes, and were most similar to *B. japonicum* group I for the *recA* gene. It should be noted that, in a previous study, we showed that *atpD* had limitations for the study of some strains of *B. elkanii*, but not with *B. japonicum* (Menna *et al.*, 2009). Now, in this study, group Ia strains shared higher identities in their *atpD* and *glnII* genes than with all other *Bradyrhizobium* species, but that was not the case for the *recA* gene (Table 2). Greater divergence in *recA* in comparison with other genes has also been found in the study of Chahboune *et al.* (2012). In the concatenated *atpD* + *glnII* + *recA* MLSA phylogram, group Ia strains also formed a distinct clade and were part of the group of species belonging to the *B. japonicum* 16S rRNA gene phylogenetic branch, having *B. betae* as its closest taxon (Fig. 2). Similar results were obtained when sequences of all six protein-encoding genes were analysed (*atpD*, *glnII*, *recA*, *gyrB*, *rpoB* and *dnaK*) (Fig. S3), confirming that sequences of three housekeeping genes seem to be sufficient for taxonomic classification of bradyrhizobia.

The draft genome sequence of group Ia strain CPAC 7, covering >99% of the genome (our unpublished data),

and the complete genome sequences of group Ia strain USDA 110 (Kaneko *et al.*, 2002) and *B. japonicum* USDA 6^T (Kaneko *et al.*, 2011) were used to estimate DDH,




Fig. 2. Neighbour-joining phylogeny based on concatenated *atpD*, *glnII* and *recA* gene sequences showing relationships between strains from the novel species (shown in bold) and other members of the genus *Bradyrhizobium*. Bootstrap values >70% are indicated at nodes. *Rhodopseudomonas palustris* CGA009 and *Xanthobacter autotrophicus* Py2 were used as outgroups. Bar, 2 substitutions per 100 nucleotide positions.

Table 3. Distinctive phenotypic features of *B. diazoefficiens* sp. nov. and phylogenetically related species

Strains: 1, SEMIA 5060; 2, CPAC 7; 3, SEMIA 6059; 4, USDA 110^T; 5, *B. japonicum* USDA 6^T; 6, *B. betae* LMG 21987^T; 7, *B. canariense* LMG 22265^T; 8, *B. yuanningense* LMG 21827^T; 9, *B. cytisi* CTAW11^T; 10, *B. huanghuaihaiense* CCBAU 23303^T; 11, *B. iriomotense* EK05^T; 12, '*B. rifense*' CTAW71; 13, *B. liaoningense* LMG 18230^T. Data were obtained in this study unless indicated. +, Growth; -, no growth; w, weakly positive; ND, not determined. Carbon-source utilization was evaluated with the API 50CH kit (bioMérieux).

Characteristic	1	2	3	4	5	6	7	8	9	10	11	12	13
Carbon-source utilization													
Glycerol	w	+	w	w	w	w	w	w	w	w	+	+	-
D-Arabinose	+	+	+	+	w	+	+	w	+	+	+	+	w
L-Arabinose	w	+	w	w	+	+	w	+	+	w	+	+	+
D-Ribose	+	+	+	+	+	+	+	+	w	+	+	+	+
D-Xylose	w	+	w	w	w	w	w	+	+	w	+	+	+
L-Xylose	+	+	+	+	w	+	+	+	+	+	+	+	w
D-Adonitol	-	w	w	-	-	-	w	-	w	-	w	+	-
Methyl β-D-xylopyranoside	w	w	w	w	w	-	w	w	w	w	w	w	-
D-Glucose	w	w	w	w	w	w	w	w	w	w	-	w	-
D-Fructose	w	w	w	w	w	w	w	w	w	w	+	w	-
D-Mannose	w	w	w	w	w	w	w	w	w	w	w	+	+
L-Rhamnose	w	w	w	w	w	w	+	w	w	w	w	w	w
Inositol	-	-	-	-	-	-	-	-	w	-	w	w	-
D-Mannitol	w	w	w	w	w	-	w	w	w	w	w	w	w
D-Sorbitol	w	w	w	w	w	-	w	-	w	-	-	-	-
N-Acetylglucosamine	-	-	-	-	-	-	-	-	-	-	-	w	-
Amygdalin	-	-	-	-	-	-	-	-	-	-	-	w	-
Arbutin	-	-	-	-	-	-	-	-	w	-	w	w	-
Aesculin ferric citrate	+	+	+	+	+	+	+	+	+	w	+	+	-
Salicin	-	-	-	-	-	-	-	-	w	-	w	w	-
Cellobiose	-	-	-	-	-	-	-	-	w	-	w	w	-
Maltose	-	-	-	-	-	-	-	w	-	-	-	-	-
Glycogen	-	-	-	-	-	+	+	-	+	+	+	+	-
Xylitol	-	w	w	-	-	-	-	-	-	-	w	w	-
Gentiobiose	-	-	-	-	-	-	-	-	w	-	w	+	-
L-Fucose	+	+	+	+	+	+	+	+	+	+	-	+	+
D-Arabitol	w	w	w	w	w	-	w	w	w	w	+	w	-
L-Arabitol	-	w	-	-	-	-	-	-	w	-	w	-	-
Potassium gluconate	-	+	+	-	-	-	-	-	-	+	+	-	w
Potassium 2-ketogluconate	-	-	-	-	-	+	-	-	-	-	-	-	-
Potassium 5-ketogluconate	-	-	+	-	-	-	-	-	+	+	-	-	-
Utilization as sole nitrogen source*													
Asparagine	+	+	+	+	-	-	w	w	+	+	+	+	+
Phenylalanine	+	+	+	+	-	w	w	-	w	+	+	+	-
Histidine	-	+	w	-	-	-	-	-	+	+	+	+	+
Leucine	+	+	+	+	w	-	-	w	+	+	+	+	w
Isoleucine	+	+	+	+	w	+	+	+	+	+	+	+	+
Growth in/at:													
37 °C	-	-	-	-	-	-	-	+	-	-	-	-	-
pH 4.5	+	+	+	+	+	+	+	w	+	+	+	w	w
pH 8	+	+	-	-	+	+	-	+	-	+	-	-	+
Resistance to (µg ml ⁻¹)													
Ampicillin (10)	-	+	w	w	-	-	w	-	+	+	+	+	-
Chloramphenicol (30)	+	+	+	+	+	+	+	+	-	+	-	+	w
Erythromycin (15)	+	+	+	-	+	+	+	+	+	+	+	+	-
Nalidixic acid (30)	+	+	+	+	+	+	+	+	+	+	w	+	+
Neomycin (30)	-	-	-	-	-	w	-	-	w	w	+	w	-
Penicillin G (10 U)	+	+	+	+	-	w	+	w	+	+	+	+	-
Tetracycline (30)	+	+	+	-	-	+	+	+	-	w	-	-	w
Streptomycin (10)	-	-	-	-	-	-	+	-	-	w	w	-	-

Table 3. cont.

Characteristic	1	2	3	4	5	6	7	8	9	10	11	12	13
Carbenicillin (500)	–	–	–	–	–	–	+	–	ND	ND	ND	ND	–
Colony size (mm) after 7–10 days of incubation on YMA†	1.3	1.5	1.2	1.3	0.7	0.7	0.9	1.2	<1 ^{a‡}	<1 ^b	0.2–1 ^c	<1 ^d	0.3

*Growth (OD₆₀₀) was considered positive if ≥ 0.09 , weak at 0.04–0.09, and negative at < 0.04 .

†Mean of three colonies.

‡Data taken from: a, Chahboune *et al.* (2011); b, Zhang *et al.* (2012); c, Islam *et al.* (2008); d, Chahboune *et al.* (2012).

average nucleotide identity (ANI) and G+C content. A digital DDH methodology using genomic sequences correlated well with experimental determinations (Auch *et al.*, 2010) and was used in our study to estimate the DDH of group Ia strains USDA 110 and CPAC 7, and of *B. japonicum* strains USDA 6^T and of CPAC 15, based on the draft genome sequence of this strain ($>99\%$ of the genome; our unpublished data) (Table S3). Based on published recommendations (Auch *et al.*, 2010), we used the BLAT program and formula 3 for calculations. The estimated DDH between group Ia strains USDA 110 and CPAC 7 was 89%, indicating that the two strains belong to the same species; in contrast, these two strains showed only 57.8 and 58.9% DDH, respectively, with *B. japonicum* USDA 6^T, strongly indicating that the group Ia strains do not belong to the species *B. japonicum*. Experimental DDH values reported in the literature between group Ia strain USDA 110 and *B. japonicum* USDA 6^T are 44% (Urtz & Elkan, 1996), 55.5% (Hollis *et al.*, 1981), 65% (Rivas *et al.*, 2004) and 65.5% (Willems *et al.*, 2001a), all below the recommended 70% threshold for species circumscription in prokaryotes (Stackebrandt & Goebel, 1994). In addition, low DDH values were also reported in the comparison of USDA 110 with the type strains of *B. liaoningense* (61%) and *B. betae* (63%) (Rivas *et al.*, 2004). Willems *et al.* (2001a) used a DDH value of 60% as a threshold for genospecies delineation in *Bradyrhizobium*; however, close inspection of their published DDH data indicates that this ‘relaxed’ value was probably used to accommodate group I and Ia strains into a single genospecies and a group of heterogeneous *Aeschynomene* strains into genospecies VI. The same authors recognized that genospecies VI seems to comprise at least two highly related subgroups (Willems *et al.*, 2001a), and that group I and Ia strains most likely correspond to different taxa (Willems *et al.*, 2001b). Interestingly, by using the approach of hybridization with cosmid profiles, Kuykendall *et al.* (1992) reported a mean value for the coefficient of similarity of only 0.36 between group I and Ia strains; in addition, the coefficients among group Ia strains (USDA 110, USDA 62, 61A50, 5631 and 8) were homogeneous at 0.88–0.94.

ANI of genome sequences has been proposed as an alternative to DDH in prokaryotic taxonomy. First, it has been proposed that an ANI $>94\%$ would correspond to 70% DDH (Konstantinidis & Tiedje, 2004) and, lately,

$>95\text{--}96\%$ ANI has been accepted as the threshold for species delineation (Richter & Rosselló-Móra, 2009). Group Ia strains USDA 110 and CPAC 7 shared an ANI value of 98.7%, indicating that this pair of strains belongs to the same species. *B. japonicum* strains USDA 6^T and CPAC 15 also shared a high ANI value (98.4%), characteristic of isolates from the same species. In contrast, when group Ia strains were compared with *B. japonicum* strains, low values were obtained ($\sim 91\%$), showing that these groups constitute different species (Table S3).

In conclusion, the results presented here emphasize DDH values lower than the 70% threshold, and, in the comparison of USDA 6^T and USDA 110, include compiled data from experiments reported in the literature ranging from 44 to 65.5%, as well as *in silico* results from our study for DDH (57.8%) and ANI (91.1%).

The G+C content was estimated by analysing genomic sequences with BioEdit (Hall, 1999). First, the sequences of 13 contigs representing the CPAC 7 genome were concatenated. The calculated DNA G+C content of group Ia strain CPAC 7 was 63.98 mol%, which falls within the 60–64 mol% range of experimental values reported for *Bradyrhizobium* species (Xu *et al.*, 1995), and is closer to the 64.1 mol% value for group Ia strain USDA 110 than to the 63.67 mol% value for *B. japonicum* USDA 6^T.

It is also interesting to comment on the symbiotic component of these strains. In a previous study from our group, Menna & Hungria (2011) reported high similarity of common (*nodA*) and host-specific (*nodZ*) nodulation genes for symbionts of soybean and belonging to groups I and Ia. In addition, group I and Ia soybean strains showed high similarity of nitrogen fixation genes, including *nifD* (Koppell & Parker, 2012; Parker *et al.*, 2002) and *nifH* (Menna & Hungria, 2011). However, strains from groups I and Ia but symbionts of other hosts, such as SEMIA 6059, isolated from *P. tetragonolobus* and unable to nodulate soybean, were positioned in different clusters in the study of Menna & Hungria (2011), and similar results were obtained for other strains used by Parker *et al.* (2002) and Koppell & Parker (2012); in this last case, divergence was also associated with biogeography. Therefore, these results indicate the existence of additional symbiovars besides *sv. glycineraum* in both group I and Ia strains.

To compare group Ia strains USDA 110, CPAC 7, SEMIA 6059 and SEMIA 5060 further with *B. japonicum* strains USDA 6^T, CPAC 15 and SEMIA 566, fatty acids were analysed. Profiles were determined using the MIDI Sherlock Microbial Identification System (MIDI, 2001) with the TSBA6 database after growth of strains on YMA for 5 days. All strains analysed showed C_{16:0} and summed feature 8 (C_{18:1ω7c} and/or C_{18:1ω6c}) as major fatty acids (Table S4), a typical characteristic of *Bradyrhizobium* (Tighe *et al.*, 2000). However, group Ia strains were distinguished from *B. japonicum* strains by having larger relative amounts of summed feature 8 and lacking C_{16:1ω5c} and 11-methyl C_{18:1ω7c} (Table S4). These results confirm differences reported previously in the composition of fatty acids for group Ia strain USDA 110 and group I strain USDA 6^T (Graham *et al.*, 1995; van Berkum & Fuhrmann, 2001).

Several phenotypic characteristics were determined for group Ia strains and compared with those of strains of *B. japonicum* and *B. betae*, the two most closely related species. *B. canariense*, *B. yuanmingense*, *B. cytisi*, *B. huanghuaihaiense*, *B. iriomotense*, *B. liaoningense* and 'B. rifense' strains were also included as representatives of more distant bradyrhizobial groups within the *B. japonicum* 16S rRNA gene phylogenetic branch. Unless indicated otherwise, all tests were performed at 28 °C. Colony morphology and size, acid/alkaline production and tolerance to pH and temperature were determined in liquid YM while tolerance to 1% NaCl on YMA, as described before (Hungria *et al.*, 2001). Ability to growth in liquid Luria–Bertani (LB) medium was also tested. Antibiotic resistance was evaluated using the disc diffusion method on YMA plates, or in liquid to test for high-level resistance to carbenicillin. Carbon-source utilization was determined using the API 50CH kit (bioMérieux), according to the manufacturer's instructions, using YM-minus-mannitol as the basal medium. Nitrogen source utilization was analysed in the defined medium of Brown & Dilworth (Bergersen, 1980) supplied with 0.25% glucose as the carbon source. Dissolved compounds were filter-sterilized and added to a final concentration of 0.7%. All tests were performed in triplicate. Group Ia strains analysed in this study showed similar reactions in most morphophysiological tests, although levels of carbon-source utilization were relatively variable (Table 3). We observed that, in general, bradyrhizobia showed weak carbon-source utilization when evaluated with the API 50CH kit. Nevertheless, at least five phenotypic differences were observed between group Ia strains and all tested *Bradyrhizobium* species (Table 3). Colony size was larger for group Ia strains than for the other species, except for *B. yuanmingense* (Table 3). A cluster analysis of strains analysed based on the phenotypic data showed that group Ia strains are more similar to each other than to strains of other species (Fig. S4).

In addition to the four strains from this study, several others from our previous studies, all symbionts of soybean, also fitted into group Ia (Chueire *et al.*, 2003; Germano *et al.*, 2006; Menna *et al.*, 2009) (Table S5).

Although strain USDA 6^T was not included in many studies from Japan and Thailand, undoubtedly several isolates from soybean also fit into group Ia (Minamisawa, 1989, 1990; Yokoyama *et al.*, 1999). Other group Ia strains from the pioneer studies of Hollis *et al.* (1981), Huber *et al.* (1984), Kuykendall *et al.* (1992) and van Berkum & Fuhrmann (2001) are listed in Table S5, together with strains from other recent studies.

Based on all the genotypic and phenotypic evidence presented in this study, we propose the reclassification of former *B. japonicum* group Ia strains into a novel species named *Bradyrhizobium diazoefficiens* sp. nov. The novel species includes several strains, the great majority isolated from soybean. It is worth mentioning that two of the strains used in our study are outstanding in their capacity to fix nitrogen with soybean: CPAC 7 is the most efficient strain in nitrogen fixation capacity used in commercial inoculants in Brazil (Hungria *et al.*, 2006; Mendes *et al.*, 2004), and USDA 110 has been used in commercial inoculants in USA and in other countries and is now being broadly evaluated in Africa, so far with excellent results (<http://www.n2africa.org/>). Other strains in this group are also very effective in fixing nitrogen, with especially high rates with the soybean crop.

Description of *Bradyrhizobium diazoefficiens* sp. nov.

Bradyrhizobium diazoefficiens (di.a.zo.ef.fi'ci.ens. L. inseparable particle *dis* twice, doubly; N.L. n. *azotum* nitrogen; N.L. pref. *diazo-* pertaining to dinitrogen; L. part. adj. *efficiens* efficient; N.L. part. adj. *diazoefficiens* dinitrogen efficient, referring to the efficiency in nitrogen fixation displayed by several strains of the species).

Cells are Gram-negative, aerobic rods, as for other species of the genus *Bradyrhizobium*. Colonies are 1.2–1.5 mm in diameter, circular, convex, opaque and slightly pink on YMA containing Congo red after 7 days of growth at 28 °C. All known strains produce an alkaline reaction in YM. The strains grow at pH 4.5, with optimum growth at pH 6.8. They grow optimally at 28 °C, and are unable to grow at 37 °C. They do not grow on YMA in the presence of 1% NaCl or in LB broth. Carbon-source utilization API tests were positive for D-arabinose, D-ribose, L-xylose, aesculin ferric citrate, starch, D-lyxose, D-fucose and L-fucose as carbon sources, weakly positive with glycerol, L-arabinose, D-xylose, methyl β-D-xylopyranoside, D-galactose, D-glucose, D-fructose, D-mannose, L-rhamnose, D-mannitol, D-sorbitol and D-arabitol and weak or negative with D-adonitol and xylitol. Strains can assimilate asparagine, glutamine, phenylalanine, leucine, methionine, isoleucine, proline, threonine, tryptophan and valine as nitrogen sources. Strains are resistant (μg per disc unless indicated otherwise) to the antibiotics nalidixic acid (30 μg), chloramphenicol (30 μg) and penicillin G (10 U), and sensitive to neomycin (30 μg), cefuroxime (30 μg), streptomycin (10 μg) and carbenicillin (500 μg).

ml⁻¹). Several strains are very effective in nodulating and fixing nitrogen when in symbiosis with soybean [*Glycine max* (L.) Merr.], but the species also includes symbionts of other host legumes.

The type strain is USDA 110^T (=IAM 13628^T =CCRC 13528^T =NRRL B-4361^T =NRRL B-4450^T =TAL 102^T =BCRC 13528^T =JCM 10833^T =TISTR 339^T =SEMIA 5032^T =3I1B110^T =ACCC 15034^T =CCT 4249^T =NBRC 14792^T =R-12974^T =CNPSO 46^T), isolated from an effective nodule of soybean [*Glycine max* (L.) Merr.] in 1959 in Florida, USA. Its genomic DNA G+C content is 63.98 mol%.

Acknowledgements

The study was partially supported by CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico, Brazil), Project Repensa (562008/2010-1). The authors acknowledge Ligia Maria O. Chueire, Dr Pâmela Menna (Embrapa Soja) and Márcia Maria Parma (Embrapa Meio Ambiente) for help in several steps of this work, Dr Allan R. J. Eaglesham for suggestions on the manuscript and Dr Manuel López-Muñoz and Professor Jean P. Euzéby for their help in the creation of the specific epithet. J. R. M. D. acknowledges a PhD fellowship from CAPES and M. H. is also a research fellow of CNPq (300547/2010-2). The laboratories had a bilateral collaboration project CNPq/CONACYT (490048/2009-9). Approved for publication by the Editorial Board of Embrapa Soja as manuscript 401/2012.

References

- Auch, A. F., von Jan, M., Klenk, H. P. & Göker, M. (2010). Digital DNA-DNA hybridization for microbial species delineation by means of genome-to-genome sequence comparison. *Stand Genomic Sci* 2, 117–134.
- Bergersen, F. J. (1980). *Methods for Evaluating Biological Nitrogen Fixation*. Chichester: Wiley.
- Boddey, L. H. & Hungria, M. (1997). Phenotypic grouping of Brazilian *Bradyrhizobium* strains which nodulate soybean. *Biol Fertil Soils* 25, 407–415.
- Chahboune, R., Carro, L., Peix, A., Barrijal, S., Velázquez, E. & Bedmar, E. J. (2011). *Bradyrhizobium cytisi* sp. nov., isolated from effective nodules of *Cytisus villosus*. *Int J Syst Evol Microbiol* 61, 2922–2927.
- Chahboune, R., Carro, L., Peix, A., Ramírez-Bahena, M. H., Barrijal, S., Velázquez, E. & Bedmar, E. J. (2012). *Bradyrhizobium rifense* sp. nov. isolated from effective nodules of *Cytisus villosus* grown in the Moroccan Rif. *Syst Appl Microbiol* 35, 302–305.
- Chang, Y. L., Wang, J. Y., Wang, E. T., Liu, H. C., Sui, X. H. & Chen, W. X. (2011). *Bradyrhizobium lablabi* sp. nov., isolated from effective nodules of *Lablab purpureus* and *Arachis hypogaea*. *Int J Syst Evol Microbiol* 61, 2496–2502.
- Chueire, L. M. O., Bangel, E. V., Mostasso, F. L., Campo, R. J., Pedrosa, F. O. & Hungria, M. (2003). Classificação taxonômica das estirpes de rizóbio recomendadas para as culturas da soja e do feijoeiro baseada no seqüenciamento do gene 16S rRNA. *Rev Bras Cienc Solo* 27, 833–840 (in Portuguese).
- Delamuta, J. R. M., Ribeiro, R. A., Menna, P., Bangel, E. V. & Hungria, M. (2012). Multilocus sequence analysis (MLSA) of *Bradyrhizobium* strains: revealing high diversity of tropical diazotrophic symbiotic bacteria. *Braz J Microbiol* 43, 698–710.
- Felsenstein, J. (1981). Evolutionary trees from DNA sequences: a maximum likelihood approach. *J Mol Evol* 17, 368–376.
- Ferreira, M. C. & Hungria, M. (2002). Recovery of soybean inoculant strains from uncropped soils in Brazil. *Field Crops Res* 79, 139–152.
- Fred, E. B., Baldwin, I. L. & McCoy, E. (1932). *Root Nodule Bacteria of Leguminous Plants*. Madison, WI: University of Wisconsin Press.
- Germano, M. G., Menna, P., Mostasso, F. L. & Hungria, M. (2006). RFLP analysis of the rRNA operon of a Brazilian collection of bradyrhizobial strains from 33 legume species. *Int J Syst Evol Microbiol* 56, 217–229.
- Graham, P. H., Sadowsky, M. J., Tighe, S. W., Thompson, J. A., Date, R. A., Howieson, J. G. & Thomas, R. (1995). Differences among strains of *Bradyrhizobium* in fatty-acid methyl-ester analysis. *Can J Microbiol* 41, 1038–1042.
- Hall, T. A. (1999). BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp Ser* 41, 95–98.
- Hollis, A. B., Kloos, W. E. & Elkan, G. H. (1981). DNA:DNA hybridization studies of *Rhizobium japonicum* and related *Rhizobiaceae*. *J Gen Microbiol* 123, 215–222.
- Huber, T. A., Agarwal, A. K. & Keister, D. L. (1984). Extracellular polysaccharide composition, *ex planta* nitrogenase activity, and DNA homology in *Rhizobium japonicum*. *J Bacteriol* 158, 1168–1171.
- Hungria, M., Chueire, L. M. O., Coca, R. G. & Megias, M. (2001). Preliminary characterization of fast growing rhizobial strains isolated from soybean nodules in Brazil. *Soil Biol Biochem* 33, 1349–1361.
- Hungria, M., Campo, R. J., Mendes, I. C. & Graham, P. H. (2006). Contribution of biological nitrogen fixation to the N nutrition of grain crops in the tropics: the success of soybean (*Glycine max* L. Merr.) in South America. In *Nitrogen Nutrition and Sustainable Plant Productivity*, pp. 43–93. Edited by R. P. Singh, N. Shankar & P. K. Jaiwal. Houston: Studium Press.
- Islam, M. S., Kawasaki, H., Muramatsu, Y., Nakagawa, Y. & Seki, T. (2008). *Bradyrhizobium iriomotense* sp. nov., isolated from a tumor-like root of the legume *Entada koshunensis* from Iriomote Island in Japan. *Biosci Biotechnol Biochem* 72, 1416–1429.
- Jordan, D. C. (1982). Transfer of *Rhizobium japonicum* Buchanan 1980 to *Bradyrhizobium* gen. nov., a genus of slow-growing, root nodule bacteria from leguminous plants. *Int J Syst Bacteriol* 32, 136–139.
- Kaneko, T., Nakamura, Y., Sato, S., Minamisawa, K., Uchiimi, T., Sasamoto, S., Watanabe, A., Idesawa, K., Iriguchi, M. & other authors (2002). Complete genomic sequence of nitrogen-fixing symbiotic bacterium *Bradyrhizobium japonicum* USDA110. *DNA Res* 9, 189–197.
- Kaneko, T., Maita, H., Hirakawa, H., Uchiike, N., Minamisawa, K., Watanabe, A. & Sato, S. (2011). Complete genome sequence of the soybean symbiont *Bradyrhizobium japonicum* strain USDA6^T. *Genes* 2, 763–787.
- Kaschuk, G., Hungria, M., Andrade, D. S. & Campo, R. J. (2006). Genetic diversity of rhizobia associated with common bean (*Phaseolus vulgaris* L.) grown under no-tillage and conventional systems in Southern Brazil. *Appl Soil Ecol* 32, 210–220.
- Konstantinidis, K. T. & Tiedje, J. M. (2004). Trends between gene content and genome size in prokaryotic species with larger genomes. *Proc Natl Acad Sci U S A* 101, 3160–3165.
- Koppell, J. H. & Parker, M. A. (2012). Phylogenetic clustering of *Bradyrhizobium* symbionts on legumes indigenous to North America. *Microbiology* 158, 2050–2059.
- Kuykendall, L. D., Saxena, B., Devine, T. E. & Udell, S. E. (1992). Genetic diversity in *Bradyrhizobium japonicum* Jordan 1982 and a

- proposal for *Bradyrhizobium elkanii* sp. nov. *Can J Microbiol* **38**, 501–505.
- Lloret, L. & Martínez-Romero, E. (2005).** Evolución y filogenia de rizobia. *Rev Latinoam Microbiol* **47**, 43–60 (in Spanish).
- Löhnis, F. & Hansen, R. (1921).** Nodule bacteria of leguminous plants. *J Agric Res* **20**, 543–556.
- Mendes, I. C., Hungria, M. & Vargas, M. A. T. (2004).** Establishment of *Bradyrhizobium japonicum* and *B. elkanii* strains in a Brazilian Cerrado oxisol. *Biol Fertil Soils* **40**, 28–35.
- Menna, P. & Hungria, M. (2011).** Phylogeny of nodulation and nitrogen-fixation genes in *Bradyrhizobium*: supporting evidence for the theory of monophyletic origin, and spread and maintenance by both horizontal and vertical transfer. *Int J Syst Evol Microbiol* **61**, 3052–3067.
- Menna, P., Hungria, M., Barcellos, F. G., Bangel, E. V., Hess, P. N. & Martínez-Romero, E. (2006).** Molecular phylogeny based on the 16S rRNA gene of elite rhizobial strains used in Brazilian commercial inoculants. *Syst Appl Microbiol* **29**, 315–332.
- Menna, P., Barcellos, F. G. & Hungria, M. (2009).** Phylogeny and taxonomy of a diverse collection of *Bradyrhizobium* strains based on multilocus sequence analysis of the 16S rRNA gene, ITS region and *glnII*, *recA*, *atpD* and *dnaK* genes. *Int J Syst Evol Microbiol* **59**, 2934–2950.
- MIDI (2001).** *Sherlock Microbial Identification System Operating Manual, version 4.0*. Newark: MIDI, Inc.
- Minamisawa, K. (1989).** Comparison of extracellular polysaccharide composition, rhizobitoxine production, and hydrogenase phenotype among various strains of *Bradyrhizobium japonicum*. *Plant Cell Physiol* **30**, 877–884.
- Minamisawa, K. (1990).** Division of rhizobitoxine-producing and hydrogen-uptake positive strains of *Bradyrhizobium japonicum* by *nifDKE* sequence divergence. *Plant Cell Physiol* **31**, 81–89.
- Minamisawa, K., Isawa, T., Nakatsuka, Y. & Ichikawa, N. (1998).** New *Bradyrhizobium japonicum* strains that possess high copy numbers of the repeated sequence RS alpha. *Appl Environ Microbiol* **64**, 1845–1851.
- Moulin, L., Béna, G., Boivin-Masson, C. & Stepkowski, T. (2004).** Phylogenetic analyses of symbiotic nodulation genes support vertical and lateral gene co-transfer within the *Bradyrhizobium* genus. *Mol Phylogenet Evol* **30**, 720–732.
- Norris, D. O. (1965).** Acid production by *Rhizobium* a unifying concept. *Plant Soil* **22**, 143–166.
- Ormeño-Orrillo, E., Hungria, M. & Martínez-Romero, E. (2013).** Dinitrogen-fixing prokaryotes. In *The Prokaryotes*, 4th edn, pp. 427–451. Edited by M. Dworkin, S. Falkow, E. Rosenberg, K.-H. Schleifer & E. Stackebrandt. Berlin & Heidelberg: Springer.
- Parker, M. A., Lafay, B., Burdon, J. J. & van Berkum, P. (2002).** Conflicting phylogeographic patterns in rRNA and *nifD* indicate regionally restricted gene transfer in *Bradyrhizobium*. *Microbiology* **148**, 2557–2565.
- Provorov, N. A. & Vorob'ev, N. I. (2000).** Evolutionary genetics of nodule bacteria: molecular and population aspects. *Russ J Genet* **36**, 1323–1335.
- Richter, M. & Rosselló-Móra, R. (2009).** Shifting the genomic gold standard for the prokaryotic species definition. *Proc Natl Acad Sci U S A* **106**, 19126–19131.
- Rivas, R., Willems, A., Palomo, J. L., García-Benavides, P., Mateos, P. F., Martínez-Molina, E., Gillis, M. & Velázquez, E. (2004).** *Bradyrhizobium betae* sp. nov., isolated from roots of *Beta vulgaris* affected by tumour-like deformations. *Int J Syst Evol Microbiol* **54**, 1271–1275.
- Saitou, N. & Nei, M. (1987).** The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* **4**, 406–425.
- Santos, M. A., Vargas, M. A. T. & Hungria, M. (1999).** Characterization of soybean *Bradyrhizobium* strains adapted to the Brazilian savannas. *FEMS Microbiol Ecol* **30**, 261–272.
- So, R. B., Ladha, J. K. & Young, J. P. (1994).** Photosynthetic symbionts of *Aeschynomene* spp. form a cluster with bradyrhizobia on the basis of fatty acid and rRNA analyses. *Int J Syst Bacteriol* **44**, 392–403.
- Stackebrandt, E. & Goebel, B. M. (1994).** Taxonomic note: a place for DNA-DNA reassociation and 16S rRNA sequence analysis in the present species definition in bacteriology. *Int J Syst Bacteriol* **44**, 846–849.
- Stepkowski, T., Moulin, L., Krzyżńska, A., McInnes, A., Law, I. J. & Howieson, J. (2005).** European origin of *Bradyrhizobium* populations infecting lupins and serradella in soils of Western Australia and South Africa. *Appl Environ Microbiol* **71**, 7041–7052.
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M. & Kumar, S. (2011).** MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol* **28**, 2731–2739.
- Tian, C. F., Zhou, Y. J., Zhang, Y. M., Li, Q. Q., Zhang, Y. Z., Li, D. F., Wang, S., Wang, J., Gilbert, L. B. & other authors (2012).** Comparative genomics of rhizobia nodulating soybean suggests extensive recruitment of lineage-specific genes in adaptations. *Proc Natl Acad Sci U S A* **109**, 8629–8634.
- Tighe, S. W., de Lajudie, P., Dipietro, K., Lindström, K., Nick, G. & Jarvis, B. D. W. (2000).** Analysis of cellular fatty acids and phenotypic relationships of *Agrobacterium*, *Bradyrhizobium*, *Mesorhizobium*, *Rhizobium* and *Sinorhizobium* species using the Sherlock Microbial Identification System. *Int J Syst Evol Microbiol* **50**, 787–801.
- Urtz, B. E. & Elkan, G. H. (1996).** Genetic diversity among *Bradyrhizobium* isolates that effectively nodulate peanut (*Arachis hypogaea*). *Can J Microbiol* **42**, 1121–1130.
- van Berkum, P. & Fuhrmann, J. J. (2000).** Evolutionary relationships among the soybean bradyrhizobia reconstructed from 16S rRNA gene and internally transcribed spacer region sequence divergence. *Int J Syst Evol Microbiol* **50**, 2165–2172.
- van Berkum, P. & Fuhrmann, J. J. (2001).** Characterization of soybean bradyrhizobia for which serogroup affinities have not been identified. *Can J Microbiol* **47**, 519–525.
- Vinuesa, P., Rademaker, J. L., de Bruijn, F. J. & Werner, D. (1998).** Genotypic characterization of *Bradyrhizobium* strains nodulating endemic woody legumes of the Canary Islands by PCR-restriction fragment length polymorphism analysis of genes encoding 16S rRNA (16S rDNA) and 16S-23S rDNA intergenic spacers, repetitive extragenic palindromic PCR genomic fingerprinting, and partial 16S rDNA sequencing. *Appl Environ Microbiol* **64**, 2096–2104.
- Vinuesa, P., León-Barrios, M., Silva, C., Willems, A., Jarabo-Lorenzo, A., Pérez-Galdona, R., Werner, D. & Martínez-Romero, E. (2005).** *Bradyrhizobium canariense* sp. nov., an acid-tolerant endosymbiont that nodulates endemic genistoid legumes (Papilionoideae: Genisteeae) from the Canary Islands, along with *Bradyrhizobium japonicum* bv. genistearum, *Bradyrhizobium* genospecies alpha and *Bradyrhizobium* genospecies beta. *Int J Syst Evol Microbiol* **55**, 569–575.
- Vinuesa, P., Rojas-Jiménez, K., Contreras-Moreira, B., Mahna, S. K., Prasad, B. N., Moe, H., Selvaraju, S. B., Thierfelder, H. & Werner, D. (2008).** Multilocus sequence analysis for assessment of the biogeography and evolutionary genetics of four *Bradyrhizobium* species that nodulate soybeans on the Asiatic continent. *Appl Environ Microbiol* **74**, 6987–6996.
- Willems, A., Coopman, R. & Gillis, M. (2001a).** Comparison of sequence analysis of 16S–23S rDNA spacer regions, AFLP analysis and DNA–DNA hybridizations in *Bradyrhizobium*. *Int J Syst Evol Microbiol* **51**, 623–632.

- Willems, A., Daignon-Bourcier, F., Goris, J., Coopman, R., de Lajudie, P., De Vos, P. & Gillis, M. (2001b).** DNA–DNA hybridization study of *Bradyrhizobium* strains. *Int J Syst Evol Microbiol* **51**, 1315–1322.
- Willems, A., Munive, A., de Lajudie, P. & Gillis, M. (2003).** In most *Bradyrhizobium* groups sequence comparison of 16S-23S rDNA internal transcribed spacer regions corroborates DNA-DNA hybridizations. *Syst Appl Microbiol* **26**, 203–210.
- Woese, C. R. (1987).** Bacterial evolution. *Microbiol Rev* **51**, 221–271.
- Xu, L. M., Ge, C., Cui, Z., Li, J. & Fan, H. (1995).** *Bradyrhizobium liaoningense* sp. nov., isolated from the root nodules of soybeans. *Int J Syst Bacteriol* **45**, 706–711.
- Yao, Z. Y., Kan, F. L., Wang, E. T., Wei, G. H. & Chen, W. X. (2002).** Characterization of rhizobia that nodulate legume species of the genus *Lespedeza* and description of *Bradyrhizobium yuanmingense* sp. nov. *Int J Syst Evol Microbiol* **52**, 2219–2230.
- Yokoyama, T., Ando, S. & Tsuchiya, K. (1999).** Serological properties and intrinsic antibiotic resistance of soybean bradyrhizobia isolated from Thailand. *Soil Sci Plant Nutr* **45**, 505–515.
- Zhang, Y. M., Li, Y., Jr, Chen, W. F., Wang, E. T., Sui, X. H., Li, Q. Q., Zhang, Y. Z., Zhou, Y. G. & Chen, W. X. (2012).** *Bradyrhizobium huanghuaihaiense* sp. nov., an effective symbiotic bacterium isolated from soybean (*Glycine max* L.) nodules. *Int J Syst Evol Microbiol* **62**, 1951–1957.