

RESEARCH ARTICLE

Irrigation management and phosphorus addition alter the abundance of carbon dioxide-fixing autotrophs in phosphorus-limited paddy soil

Xiaohong Wu^{1,2}, Tida Ge^{3,*}, Wende Yan^{1,2}, Juan Zhou³, Xiaomeng Wei³, Liang Chen^{1,2}, Xiangbi Chen³, Paolo Nannipieri⁴ and Jinshui Wu³

¹Faculty of Life Science and Technology, Central South University of Forestry and Technology, No 498, Shaoshan South Road, Tianxin District, Changsha 410004, Hunan, China, ²Key Laboratory of Agro-Ecological Processes in Subtropical Regions and Changsha Observation and Research Station for the Agricultural Environment, Institute of Subtropical Agriculture, Chinese Academy of Sciences, No 644, Yuanda Second Road, Furong District, Changsha 410125, Hunan, China, ³National Engineering Laboratory of Applied Technology for Forestry & Ecology in Southern China, No 498, Shaoshan South Road, Tianxin District, Changsha 410004, Hunan, China and ⁴Department of Agrifood Production and Environmental Sciences, University of Firenze, Viale delle Cascine 15, Florence 50144, Italy

*Corresponding author: Institute of Subtropical Agriculture, Chinese Academy of Sciences, Changsha City, Hunan Province 410125, China.

Tel: +86-731-84619736; Fax: +86-731-8461973; E-mail: gtd@isa.ac.cn

One sentence summary: P fertiliser significantly stimulated CO₂-fixation genes abundances in a P-limited paddy soil, and the response of CO₂-fixing autotrophs to P fertiliser depended on the irrigation management.

Editor: Angela Sessitsch

ABSTRACT

In this study, we assessed the interactive effects of phosphorus (P) application and irrigation methods on the abundances of marker genes (*cbbL*, *cbbM*, *accA* and *aclB*) of CO₂-fixing autotrophs. We conducted rice-microcosm experiments using a P-limited paddy soil, with and without the addition of P fertiliser (P-treated-pot (P) versus control pot (CK)), and using two irrigation methods, namely alternate wetting and drying (AWD) and continuous flooding (CF). The abundances of bacterial 16S rRNA, archaeal 16S rRNA, *cbbL*, *cbbM*, *accA* and *aclB* genes in the rhizosphere soil (RS) and bulk soil (BS) were quantified. The application of P significantly altered the soil properties and stimulated the abundances of Bacteria, Archaea and CO₂-fixation genes under CF treatment, but negatively influenced the abundances of Bacteria and marker genes of CO₂-fixing autotrophs in BS soils under AWD treatment. The response of CO₂-fixing autotrophs to P fertiliser depended on the irrigation management method. The redundancy analysis revealed that 54% of the variation in the functional marker gene abundances could be explained by the irrigation method, P fertiliser and the Olsen-P content; however, the rhizosphere effect did not have any significant influence. P fertiliser application under CF was more beneficial in improving the abundance of CO₂-fixing autotrophs compared to the AWD treatment; thus, it is an ideal irrigation management method to increase soil carbon fixation.

Keywords: CO₂-fixing autotrophs; marker gene abundance; P fertiliser; irrigation management; rhizosphere effect

INTRODUCTION

Phosphorus (P) is regarded as a limiting resource in rice cultivation. It is an essential macronutrient that determines rice growth and productivity (Fageria, Santos and Heinemann 2011); however, rice paddy fields are often P-deficient (Haefele, Nelson and Hijmans 2014), especially those located in the red-soil region of southern China (Li et al. 2012). To meet the P requirements of crops, P fertilisers have been increasingly applied in intensive rice-cropping systems (Shen et al. 2004; Lan et al. 2012), with P application rate rising up to 120 kg P ha⁻¹ (Wang et al. 2012). Applying P fertiliser not only increases the availability of the nutrient to rice crop but also directly or indirectly changes the chemical, physical and biological properties of soil (Griffiths, Spillies and Bonkowski 2012; Yan et al. 2016). Such changes are believed to have a significant effect on the growth rates and ecological functions of various soil microbes involved in the biogeochemical carbon cycle (Černá et al. 2009; Su et al. 2015).

On the other hand, as a water-intensive crop, rice is mostly grown under irrigated conditions; and therefore, water is another important limiting resource in this ecosystem (Cai and Chen 2000). The water regime and fertiliser uptake are related physiological processes (Suriyagoda, De Costa and Lambers 2014; Somaweera et al. 2016), which could interact to modify some soil properties. The bio-availability of soil P to plants is significantly related to the oxidation–reduction status of soil (Borggaard et al. 2005; Torrent 1997), which can be influenced by irrigation management practices (Haefele et al. 2006). Soil P availability, which is sensitive to water conditions, can influence plant growth (Hua et al. 2016), which, in turn, can affect nutrient turnovers in soil, especially in the rhizosphere soil (RS) (Jones, Nguyen and Finlay 2009; Richardson et al. 2009). It is possible that irrigation management together with P fertiliser application affects the activity, biomass and the composition of soil microbial communities.

Autotrophic microorganisms, capable of converting carbon dioxide (CO₂) into organic C, play an important role in the C cycle (Yuan et al. 2012a). This process is primarily catalyzed by the ribulose-1,5-biphosphate carboxylase/oxygenase (RubisCO) enzyme in the Calvin cycle (CBB) (Benson et al. 1950). The RubisCO enzyme exists in four forms (I–IV) (Tabita 1999); the *cbbL* and *cbbM* genes encode the large subunits of forms I and II of RubisCO, respectively, and are marker genes used in microbial ecology studies (Xiao et al. 2014; Wu et al. 2015; Lynn et al. 2017). In addition to the CBB cycle, four additional autotrophic CO₂-fixation pathways have been documented (Berg 2011). The reductive tricarboxylic acid (rTCA) cycle is a CO₂-fixation pathway of microaerophiles and anaerobes (Evans, Buchanan and Arnon 1966). The β -ATP citrate lyase enzyme is involved in this cycle and encoded by the marker gene *acIb* (Campbell, Stein and Cary 2003). The reductive acetyl-CoA pathway is considered the oldest pathway of strictly anaerobic bacteria and archaea (Wood, Ragsdale and Pezacka 1986; Berg 2011), with the key enzyme being CO dehydrogenase/acetyl-CoA synthase, but without a proper marker gene. The 3-hydroxypropionate/malyl-CoA cycle is only found in a few species of the genus *Chloroflexus* (Holo 1989) and the carboxylation reaction of this cycle is catalysed by biotin-dependent acetyl-CoA/propionyl-CoA carboxylase (Hügler 2003). The marker gene *accC* encodes the subunit of biotin-dependent acetyl-CoA carboxylase and the primers developed to target this gene have low specificity (Auguet et al. 2008). The 3-hydroxypropionate/4-hydroxybutyrate pathway is a modified version of the 3-hydroxypropionate/malyl-CoA pathway exclusively found in *Crenarchaeota* (Berg et al. 2007). The acetyl CoA carboxylase (ACCase), encoded by the *accA* gene, is

one of the key enzymes involved in CO₂ fixation in this cycle (Yakimov, Cono and Denaro 2009).

The CO₂-fixation by the autotrophic community depends on various factors, including fertiliser management (Yuan et al. 2012b; Su et al. 2015), soil type (Lynn et al. 2017), crop rotation (Wu et al. 2015), tillage (Ge et al. 2016), rhizosphere effect (Yousuf et al. 2013) and growth stage (Xiao et al. 2014). However, no previous study has assessed the effects of P fertiliser addition combined with irrigation management on the abundance of CO₂-fixation autotrophs. Understanding these effects can provide insights on the relationship between some agricultural management practices and CO₂ fixation by microbial communities of paddy soils.

We hypothesize that the interactive effect of P fertiliser and irrigation management on CO₂-fixing autotrophs can be significant. To test this hypothesis, we conducted experiments with rice microcosms to investigate the effects of the addition of P fertiliser on CO₂-fixing autotrophs in a P-limited paddy soil under continuous flooding (CF) or alternative wetting and drying treatment (AWD). We quantified the copy numbers of the bacterial 16S rRNA gene, archaeal 16S rRNA gene and four functional genes for three CO₂-fixation pathways (the CBB cycle, the rTCA cycle and the 3-hydroxypropionate/4-hydroxybutyrate cycle) in both bulk and rhizosphere soils. In addition, we assessed the soil physicochemical properties and their potential influence on CO₂-fixing autotrophic microorganisms.

MATERIALS AND METHODS

Soil sampling

Surface (0–20 cm) soil was sampled in a P-limited paddy field on 4 April 2015 during the paddy fallow season; the field is located at the Changsha Agricultural and Environmental Monitoring Station (113°19'52"E, 28°33'04"N), Hunan Province, China. This region is characterized by a subtropical humid monsoon climate, with an annual precipitation of 1330 mm, air temperature of 17.5°C, 1663 h of sunshine and a frost-free period of 274 days. The paddy soil, derived from granite red soil, is classified as Stagnic Anthrosols (Gong, Zhang and Chen 2007). The main soil properties were: 5.43 pH, 14.26 g kg⁻¹ soil organic matter (SOC), 1.45 g kg⁻¹ total N (TN), 0.75 g kg⁻¹ total P (TP), 4.96 mg kg⁻¹ available P (Olsen-P) and 7.71 cmol kg⁻¹ cation exchange capacity (CEC). After removing all visible plant roots, the soil was sieved (<2 mm), flooded with distilled water and pre-incubated for 14 days at 25°C prior to use.

Rice microcosm

We added 160 mg K kg⁻¹ dry soil as potassium chloride and 250 mg N kg⁻¹ dry soil as ammonium sulphate to the pre-incubated paddy soil as base fertilisers. Subsequently, the soil sample was divided into two parts; one part was treated with monopotassium phosphate to reach a P content of 80 mg kg⁻¹ dry soil and the other part was used as the control soil sample and no P was added. Subsequently, we packed either 1.0 kg of the P-treated soil or 1.0 kg of the control soil into PVC pots (diameter 17.2 cm and height 20 cm), corresponding to the P-treated-pot (P) or to the control pot (CK), respectively. Nylon rhizosphere bags (\varnothing 5 cm \times 17 cm, mesh 48 μ m), each containing 500 g soil, with or without added P, were placed in the centre of the corresponding P or CK pot to create the rhizosphere soil. Two-week-old rice seedlings were subsequently transplanted into the rhizosphere bags, with or without added P, with each pot having three rice seedlings. The rice pots were then flooded with distilled water to develop 2–3 cm deep standing water.

All the rice pots were placed randomly in a growth chamber (80 × 250 cm, height 120 cm) and incubated for 14 days as described by Ge et al. (2012). Half of the CK and P-treated rice pots were flooded continuously (CF) with 2–3 cm water above the soil surface throughout the incubation, whereas the other half were subjected to alternate wetting and drying (AWD). The latter soil was covered with 2–3 cm of water at the beginning of the experiment. Then, the soil was air-dried to achieve 70%–75% of water holding capacity (WHC) for 3–4 days and it was re-flooded again. The drying–rewetting cycle was repeated three times during the experiment. Therefore, we had a total of four treatments: (1) rice grown in soil with no added P under CF; (2) rice grown in soil with no added P under AWD; (3) rice grown in P-added soil under CF; and (4) rice grown in P-added soil under AWD. Each treatment was repeated three times. Destructive soil sampling was conducted 14 days after rice transplanting. The rice grew well and did not suffer from nutrient limitation during sampling (Fig. S1, Supporting Information). We sampled RS (the soil in the rhizosphere bag) and the bulk soil (BS) (the soil out of the bag). Each soil sample was divided into two sub-samples, one was stored at 4°C for physicochemical analysis and the other was frozen at –80°C for molecular analysis.

Soil physicochemical analysis

The soil pH was determined in a 1:2.5 (wt/vol) soil-to-water suspension using a pH meter (Delta 320, Mettler-Toledo Instruments Ltd., China). The soil organic carbon (SOC) and total nitrogen (TN) contents were measured by dry combustion using an elemental analyser (Vario MAX C/N, Elementar, Germany). Exchangeable ammonium (N-NH₄⁺) and nitrate (N-NO₃⁻) were extracted with 1 M KCl and determined by an automated continuous flow analyser (FIAstar 5000, Foss, Sweden). Olsen-P was extracted with NaHCO₃ (0.5 M, pH 8.5) and measured by a colorimetric method (Murphy and Riley 1962). Soil microbial biomass carbon (MBC) was measured by the chloroform fumigation-extraction method described by Wu et al. (1990).

Soil DNA extraction

Soil DNA was extracted from 0.5 g of frozen soil (–80°C) using the FastDNA Spin Kit for soil (MP Biomedicals, OH, USA), following the manufacturer's instructions. Subsequently, the DNA extracts were diluted with nuclease-free water before the integrity and quality check. The integrity of the DNA was evaluated with 1% agarose gel. The DNA concentration and quality were determined using NanoDrop spectrophotometer (NanoDrop ND-1000, PeqLab, Germany).

Table 1. Three-way analysis of the influences of P addition (P), rhizosphere effect (R), irrigation management (W), and their interactions on the abundances of 16S rRNA genes and autotrophic CO₂ fixation genes.

Source of variance	Variable					
	Bacterial 16s rRNA	Archaeal 16S rRNA	<i>cbbL</i>	<i>cbbM</i>	<i>accA</i>	<i>aclB</i>
P	0.020	0.000	0.147	0.179	0.006	0.000
R	0.004	0.001	0.000	0.082	0.818	0.532
W	0.054	0.983	0.001	0.000	0.008	0.004
P × R	0.555	0.133	0.000	0.006	0.060	0.054
P × W	0.000	0.183	0.000	0.000	0.232	0.013
R × W	0.030	0.028	0.000	0.056	0.080	0.500
P × R × W	0.119	0.074	0.003	0.982	0.012	0.043

Significant values are shown in bold ($P < 0.05$).

Quantification of genes

The abundances of the bacterial 16S rRNA gene, archaeal 16S rRNA gene and functional genes (*cbbL*, *cbbM*, *accA* and *aclB*) involved in autotrophic CO₂-fixation pathways were estimated by quantitative PCR (qPCR), using 341f/515r (López-Gutiérrez et al. 2004), 364f/934r (Großkopf, Janssen and Liesack 1998), K2f/V2r (Tolli and King 2005), *cbbM*-f/*cbbM*-r (Alfreider et al. 2003), Crena.529F/Crena.981R (Yakimov, Cono and Denaro 2009) and 892F/1204R (Campbell, Stein and Cary 2003) primers, respectively. The detailed sequences of the primer sets used in this study are listed in Table S1, Supporting Information.

We carried out qPCR in 10 μL reaction mixtures containing 1× SYBR Premix Ex Taq (Takara, Dalian, China), 1 μL diluted template DNA of approximately 5 ng, 0.15 μM respective primer, and nuclease-free water. Reactions were repeated three times for each sample using an ABI 7900 Real-Time PCR System (PerkinElmer, Applied Biosystems, USA). The conditions were as follows: initial denaturation at 95°C for 30 s, 40 cycles of denaturation at 95°C for 15 s, annealing at 50°C–62°C for 30 s (Table S1, Supporting Information) and extension at 72°C for 20 s. Series of 10-fold dilutions of plasmids that contained cloned genes of bacterial 16S rRNA, archaeal 16S rRNA, *cbbL*, *cbbM*, *accA* and *aclB* were run in parallel with the template DNA to prepare the respective standard curves for quantification. At the end of each qPCR run, melting curve analysis was performed to check the specificity of the amplification. The copy numbers of the relative genes were calculated using the SDS 2.3 software within the qPCR system.

Statistical analysis

The effects of environmental variables (P addition, irrigation management and rhizosphere effect) and their interactions on the abundances of Bacteria, Archaea and CO₂ fixing autotrophs were tested by univariate analysis of variance (UANOVA) in the SPSS 13.0 for Windows (IBM, Armonk, NY, USA), considering the significant difference at $P < 0.05$. The influence of soil properties (pH, MBC, DOC, N-NH₄⁺, N-NO₃⁻ and Olsen-P) and the environmental variables on the abundances of microbial functional genes was evaluated by redundancy analysis (RDA) using the CANOCO 5.0 for Windows (Microcomputer Power, Ithaca, NY, USA). A forward selection procedure was conducted to select soil properties and environmental variables that significantly affected microbial functional gene abundances. Following the forward selection, the pure contribution of each significant parameter, as well as their interactive contributions to changes in the abundances of microbial functional communities was




Figure 1. Abundance of (a) bacterial and (b) archaeal 16S rRNA genes in soils without (CK) or with P addition (P) under alternate wetting and drying (AWD) and continuous flooding (CF) irrigation management. Both bulk soil (BS) and rhizosphere soil (RS) were analyzed. Error bars represent standard errors of triplicate samples. Asterisks indicate significant differences.




Figure 2. Abundances of (a) *cbbL* gene, (b) *cbbM* gene, (c) *accA* gene and (d) *acbB* gene in soils without (CK) or with P addition (P) under alternate wetting and drying (AWD) and continuous flooding (CF) irrigation management. Both bulk soil (BS) and rhizosphere soil (RS) were analyzed. Error bars represent standard errors of triplicate samples. Asterisks indicate significant differences.

further determined with the variance partitioning analysis by the CANOCO 5.0. The significance test was conducted using the Monte Carlo permutation test, with 999 unrestricted permutations ($P < 0.05$).

RESULTS

Abundances of Bacteria and Archaea

The number of bacterial 16S rRNA gene copies outnumbered the archaeal 16S rRNA gene copies by one to two orders of magnitude. The abundances of both genes were influenced significantly by P addition (Table 1). The copy numbers of the archaeal 16S rRNA genes in BS and RS were higher in P treatment than in CK treatment, irrespective of irrigation methods, with an average increase of 167% (Fig. 1). On the contrary, the effect of P addition on the number of bacterial 16S rRNA gene copies depended on the irrigation management method. Under CF treatment, P addition significantly increased the bacterial abundance (101% on average), regardless of the rhizosphere effect (Fig. 1), whereas the opposite (an average decrease of 21%) was observed in the AWD treatment.

The UANOVA showed a statistically significant rhizosphere effect on the abundances of the bacterial 16S rRNA and archaeal 16S rRNA genes (Table 1). RS harboured a significantly higher number of bacterial 16S rRNA gene copies than BS, whereas an

opposite pattern was observed for the number of archaeal 16S rRNA gene copies, which was on average 44% lower in RS than in BS (Fig. 1). Although no significant differences in gene copy numbers were observed between the two types of irrigation management methods, the interaction of the water treatment method and the rhizosphere effect significantly affected the gene abundances (Table 1).

Abundances of marker genes related to autotrophic CO₂ fixation

In all treatments, the *cbbL* gene copies, ranging from 1.86×10^8 to 1.93×10^9 copies g^{-1} dry soil, were higher than those of the *cbbM* gene (5.49×10^7 to 1.70×10^8 copies g^{-1} dry soil), *accA* gene (6.96×10^7 to 2.88×10^8 copies g^{-1} dry soil) and *aclB* gene (4.30×10^6 to 2.15×10^7 copies g^{-1} dry soil). The P application changed the abundances of all marker genes involved in autotrophic CO₂ fixation, according to the irrigation treatment (Table 1 and Fig. 2). As regards CF treatment, the abundances of CO₂ fixation genes were increased significantly by P addition 14 days after transplanting ($P < 0.05$), 68% and 156% increase for the *cbbL* gene, 2% and 99% for the *cbbM* gene, 82% and 71% for the *accA* gene and 149% and 142% for the *aclB* gene in BS and RS, compared to the CK treatment. However, in the AWD treatment, the effects of P fertiliser on the CO₂-fixation functional genes differed between BS and RS. Phosphorus application decreased the




Figure 3. RDA analysis of autotrophic CO₂ fixation gene abundances, soil properties and environmental variables that significantly affected microbial variations. Pink and blue colours denote alternate wetting and drying (AWD) and continuous flooding (CF), respectively. The circle and square indicate bulk soil (BS) and rhizosphere soil (RS), respectively. The hollow and solid symbols indicate soils without (CK) and with P addition (P), respectively.




Figure 4. Variation partitioning analysis of the pure effects of Olsen-P (O), irrigation management (W), P addition (P) and the interactions of a combination of these factors ($O \times W$, $O \times P$, $W \times P$ and $O \times W \times P$) on the abundances of the autotrophic CO_2 fixing community. The circles on the edges of the triangle show the variation due to each factor alone. The square on the sides of the triangle and the circle at the centre of the triangle show the interactions of any two factors and the interactions among all the factors, respectively. The unexplained portion is shown by the square at the bottom of the triangle. The geometric areas of each diagram are proportional to the respective percentages of the explained variation by a single factor or a combination of factors.

number of *cbbL*, *cbbM*, *accA* and *aclB* genes in BS of the CK treatment, whereas it increased the gene abundances in RS compared to the CK treatment, except for the *cbbM* gene (Fig. 2).

The influence of the rhizosphere effect was only significant for the *cbbL* gene (Table 1), with the gene abundance being on average 260% greater in RS than in BS (Fig. 2). The irrigation method significantly affected the abundances of all functional genes involved in the autotrophic CO_2 fixation (Table 1). For example, the abundances of the *cbbL*, *accA* and *aclB* genes in BS treated with P fertiliser were higher in the CF than in the AWD treatment. However, the *cbbM* gene showed an opposite pattern (Fig. 2). The UANOVA showed that the interaction between the added P, the rhizosphere effect and the irrigation management method significantly affected the abundances of most functional genes involved in autotrophic CO_2 fixation (Table 1).

Relationships among microbial functional genes, management practices, and soil properties

The first two axes of the RDA explained 42.97% and 16.44%, respectively, of the total variation in autotrophic CO_2 -fixation gene abundances (Fig. 3). The Olsen-P ($P = 0.001$), irrigation management method ($P = 0.001$) and P addition ($P = 0.001$) significantly influenced the abundances of autotrophic CO_2 -fixing microorganisms. Variance partitioning analysis showed that these variables independently explained 12.8%, 42.1% and 17.3% of the total observed variation, respectively, leaving 46%

of the variation unexplained (Fig. 4). The interactions among Olsen-P, irrigation management method and P addition had a smaller effect than each individual property, except for the shared variation between Olsen-P and irrigation management method (−13.7%).

DISCUSSION

To quantify the response of the abundances of microbial communities involved in the CO_2 -fixation to P fertiliser addition may help understand the potential ecological functions of these communities (Su et al. 2015). The addition of P fertiliser can change the soil properties such as enzyme activities and the amount of available nutrients, which may affect the size, activity and composition of soil microbial communities sensitive to such changes (Malik, Marschner and Khan 2012; Krey et al. 2013; Van Geel et al. 2016). Therefore, the differences in soil properties, such as nutrient concentrations (Table S2, Supporting Information), may explain the differences in the abundances of bacterial, archaeal and CO_2 -fixing autotrophic communities in CK and P soils as shown by the qPCR and CCA analyses. The increase in the abundance of functional microbial genes in soils amended with P fertiliser is likely caused by higher organic C inputs to soil owing to the stimulation of root growth and root exudation by added fertiliser (Tan et al. 2013; Wang et al. 2016). In this study, the application of P increased the abundances of autotrophic CO_2 -fixing microorganisms in RS except the *cbbM* genes but decreased the copy numbers of functional marker genes in BS under the AWD




Figure 5. A conceptual diagram of the pure and interacted contributions of phosphorus addition, irrigation management and rhizosphere effect to the abundances of the autotrophic CO₂ fixing community.

treatment. Probably the abundance of *cbbM* genes decreased in RS following fertiliser application in the AWD treatment because of the competition among different functional microorganisms in the microenvironment (Hibbing et al. 2010). In AWD soil, P application might have mitigated the P shortage pressure on functional groups, such as denitrifiers, and stimulated their growth under aerobic conditions (Wei et al. 2017). It might have inhibited the growth of the *cbbM*-harbouring autotrophs probably because these species were less competitive. However, the underlying mechanisms need further investigation.

The effect of P application on the abundance of CO₂-fixing autotrophic communities also depended on the irrigation condition. The irrigation management method explained 42.1% of the variations in the abundances of the four marker genes. It is well established that water conditions can affect activity, biomass and composition of soil microbial communities (Kavamura et al. 2013; Tian et al. 2013a). Irrigation condition is a crucial factor that impacts microbial biomass, through its effects on the dynamics of available C in rice cropped soils (Tian et al. 2013b). The available C may also affect the biomass and the distribution of CO₂-fixing autotrophic communities because most of these populations are facultative autotrophs, capable of using organic substrates as a carbon and an energy source (Hügler and Sievert 2011). In addition, water conditions may affect the activity of some enzymes involved in microbial CO₂ fixation, showing different affinities for oxygen (Hügler and Sievert 2011); for ex-

ample, the ATP-citrate lyase is sensitive to oxygen and the Ru-bisCO is tolerant to oxygen (Berg 2011). Accordingly, the marker genes involved in four CO₂-fixation pathways responded differently to irrigation management methods. Compared to P fertiliser addition, irrigation management contributed more to variations in the abundances of CO₂-fixing autotrophs (Fig. 5), probably because the irrigation management might have influenced the turnover and bio-availability of soil P (Haefele et al. 2006), which are quite important for the growth of CO₂-fixing autotrophs.

The Olsen-P concentration in soil was another important factor influencing the composition of CO₂-fixing autotrophic communities, probably because, as any microorganism, they require P for energy metabolism and nucleic acids and cell membrane synthesis (Rodríguez and Fraga 1999; Richardson et al. 2009). Indeed, the increase in the Olsen-P content can stimulate the growth of CO₂-fixing autotrophs, as this form of P can be directly utilized by microorganisms (Malik, Marschner and Khan 2012). Consistent with a previous study (Saleque et al. 2004), we observed a significant increase in the Olsen-P content in all soils after the application of inorganic P (Table S2, Supporting Information), but this increase was not followed by the increase in the abundances of CO₂-fixing microorganisms in BS under the AWD treatment. It may be because the AWD treatment (the soil water content reached 70%–75% of the WHC after drying for 3–4 days) limited the availability of Olsen-P to the soil microorganisms. In

addition, although the Olsen-P is considered a labile form to be easily used by microorganisms, it refers to a P extraction pool; thus, it is possible that some of this P pool may not be easily degradable by soil microbes (Malik, Marschner and Khan 2012). The Olsen-P becomes less available when bound to metal oxides in soil (Stewart and Tiessen 1987). In contrast to a previous study on CO₂ fixation gene abundances (Xiao et al. 2014), the rhizosphere effect was not significant in our study as revealed by the RDA analysis. P application can stimulate the growth of rice roots; and thus, the increase of rhizodeposition (Zhong and Cai 2007), which may increase the microbial abundance and activity in RS compared to BS. However, this is not likely to occur in our short-term experiment because of the low root biomass (Fig. S1, Supporting Information). Future research is needed to verify this hypothesis by monitoring the quality and quantity of rhizodeposition.

A considerable percentage (46%) of the variation in the abundances of CO₂-fixation genes was not explained by the measured properties. It is possible that an undetermined soil property is important in affecting the abundances of the four marker genes. For instance, the quality and quantity of root exudation can affect the abundances of functional genes in RS (Baraniya et al. 2016; Pathan et al. 2015; Yuan et al. 2016; Zhu, Vivanco and Manter 2016) and this may also happen for the CO₂ fixing microorganisms that are facultative autotrophic. The quality and quantity of root exudation depend on different types and rates of fertiliser application and irrigation management (Zhong and Cai 2007). Both acid and alkaline phosphomonoesterase activities of soils depend on the type and the rate of fertiliser application and irrigation treatment (Nannipieri et al. 2011; Tan et al. 2013) and they may also play a role in affecting the abundance of CO₂-fixing autotrophs because both enzyme activities mineralize organic P to inorganic P.

CONCLUSIONS

The addition of P significantly changed the soil properties and the abundances of four marker genes involved in CO₂ fixation in P-limited paddy soils. The application of P increased the abundances of CO₂-fixing autotrophs in the CF treatment, but decreased the copy numbers of the functional marker genes in BS under the AWD treatment. This indicates that the response of CO₂-fixing autotrophs to P fertiliser can be mediated by changes in the irrigation management system. Our findings revealed the positive and interactive effect of P application and CF on the abundance of CO₂-fixing autotrophs, suggesting that this combined management may potentially be more effective in increasing soil C sequestration than other tested conditions in P-limited paddy soils. However, the detection of genes does not imply that they are active; therefore, further research, including transcriptomics and proteomics studies, should be conducted to determine the activity and the expression of these CO₂ fixation genes.

SUPPLEMENTARY DATA

Supplementary data are available at [FEMSEC](https://academic.oup.com/femsec/article-abstract/93/1/2/1544604/778) online

ACKNOWLEDGEMENT

We especially thank the Public Service Technology Center, Institute of Subtropical Agriculture, Chinese Academy of Sciences for technical assistance.

FUNDING

This study was supported by the grants from the National Natural Science Foundation of China (41503081 and 41522107), National Key Research and Development Program (2016YFE0101100) and Youth Innovation Team Project of ISA, CAS (2017QNCXTD.GTD).

Conflict of interest. None declared.

REFERENCES

- Alfreider A, Vogt C, Hoffmann D et al. Diversity of ribulose-1,5-bisphosphate carboxylase/oxygenase large-subunit genes from groundwater and aquifer microorganisms. *Microb Ecol* 2003;**45**:317–28.
- Auguet JC, Borrego CM, Bañeras L et al. Fingerprinting the genetic diversity of the biotin carboxylase gene (*accC*) in aquatic ecosystems as a potential marker for studies of carbon dioxide assimilation in the dark. *Environ Microbiol* 2008;**10**: 2527–36.
- Baraniya D, Puglisi E, Ceccherini MT et al. Protease encoding microbial communities and protease activity of the rhizosphere and bulk soils of two maize lines with different N uptake efficiency. *Soil Biol Biochem* 2016;**96**:176–9.
- Benson AA, Bassham JA, Calvin M et al. The path of carbon in photosynthesis. V. Paper chromatography and radioautography of the products. *J Am Chem Soc* 1950;**72**:1710–8.
- Berg IA, Kockelkorn D, Buckel W et al. A 3-hydroxypropionate/4-hydroxybutyrate autotrophic carbon dioxide assimilation pathway in Archaea. *Science* 2007;**318**:1782–6.
- Berg IA. Ecological aspects of the distribution of different autotrophic CO₂ fixation pathways. *Appl Environ Microbiol* 2011;**77**:1925–36.
- Borggaard OK, Raben-Lange B, Gimsing AL et al. Influence of humic substances on phosphate adsorption by aluminium and iron oxides. *Geoderma* 2005;**127**:270–9.
- Cai H, Chen Q. Rice production in China in the early 21st century. *Chin Rice Res Newsl* 2000;**8**:14–16.
- Campbell BJ, Stein JL, Cary SC. Evidence of chemolithoautotrophy in the bacterial community associated with *Alvinella pompejana*, a hydrothermal vent polychaete. *Appl Environ Microbiol* 2003;**69**:5070–8.
- Černá B, Rejmánková E, Snyder JM et al. Heterotrophic nitrogen fixation in oligotrophic tropical marshes: changes after phosphorus addition. *Hydrobiologia* 2009;**627**:55–65.
- Evans MC, Buchanan BB, Arnon DI. A new ferredoxin-dependent carbon reduction cycle in a photosynthetic bacterium. *P Natl Acad Sci USA* 1966;**55**:928–34.
- Fageria N, Santos A, Heinemann A. Lowland rice genotypes evaluation for phosphorus use efficiency in tropical lowland. *J Plant Nutr* 2011;**34**:1087–95.
- Ge T, Wu X, Liu Q et al. Effect of simulated tillage on microbial autotrophic CO₂ fixation in paddy and upland soils. *Sci Rep-UK* 2016;**6**:19784.
- Ge TD, Yuan HZ, Zhu HH et al. Biological carbon assimilation and dynamics in a flooded rice—soil system. *Soil Biol Biochem* 2012;**48**:39–46.
- Gong ZT, Zhang GL, Chen ZC. *Pedogenesis and Soil Taxonomy*. Beijing, China: Science Press, 2007, 613–26.
- Griffiths BS, Spilles A, Bonkowski M. C:N:P stoichiometry and nutrient limitation of the soil microbial biomass in a grazed grassland site under experimental P limitation or excess. *Ecol Process* 2012;**1**:6.

- Großkopf R, Janssen PH, Liesack W. Diversity and structure of the methanogenic community in anoxic rice paddy soil microcosms as examined by cultivation and direct 16S rRNA gene sequence retrieval. *Appl Environ Microbiol* 1998;**64**:960–9.
- Haefele SM, Naklang K, Harnpichitvitaya D et al. Factors affecting rice yield and fertilizer response in rainfed lowlands of northeast Thailand. *Field Crop Res* 2006;**98**:39–51.
- Haefele SM, Nelson A, Hijmans RJ. Soil quality and constraints in global rice production. *Geoderma* 2014;**235–6**:250–259.
- Hibbing ME, Fuqua C, Parsek MR et al. Bacterial competition: surviving and thriving in the microbial jungle. *Nat Rev Microbiol* 2010;**8**:15–25.
- Holo H. Chloroflexus aurantiacus secretes 3-hydroxypropionate, a possible intermediate in the assimilation of CO₂ and acetate. *Arch Microbiol* 1989;**151**:252–6.
- Hua KK, Zhang WJ, Guo ZB et al. Evaluating crop response and environmental impact of the accumulation of phosphorus due to long-term manuring of vertisol soil in northern China. *Agr Ecosyst Environ* 2016;**219**:101–10.
- Hügler M, Krieger RS, Jahn M et al. Characterization of acetyl-CoA/propionyl-CoA carboxylase in *Metallosphaera sedula*. Carboxylating enzyme in the 3-hydroxypropionate cycle for autotrophic carbon fixation. *Eur J Biochem* 2003;**270**:736–44.
- Hügler M, Sievert SM. Beyond the Calvin cycle: autotrophic carbon fixation in the ocean. *Annu Rev Mar Sci* 2011;**3**:261–89.
- Jones DL, Nguyen C, Finlay RD. Carbon flow in the rhizosphere: carbon trading at the soil-root interface. *Plant Soil* 2009;**321**:5–33.
- Kavamura VN, Taketani RG, Lançon MD et al. Water regime influences bulk soil and rhizosphere of *Cereus jamacaru* bacterial communities in the Brazilian Caatinga biome. *PLoS One* 2013;**8**:e73606.
- Krey T, Vassilev N, Baum C et al. Effects of long-term phosphorus application and plant-growth promoting rhizobacteria on maize phosphorus nutrition under field conditions. *Eur J Soil Biol* 2013;**55**:124–30.
- Lan ZM, Lin XJ, Wang F et al. Phosphorus availability and rice grain yield in a paddy soil in response to long-term fertilization. *Biol Fert Soils* 2012;**48**:579–88.
- Li Y, Wu JS, Liu SL et al. Is the C:N:P stoichiometry in soil and soil microbial biomass related to the landscape and land use in southern subtropical China? *Global Biogeochem Cy* 2012;**26**:GB4002.
- López-Gutiérrez JC, Henry S, Hallet S et al. Quantification of a novel group of nitrate-reducing bacteria in the environment by real-time PCR. *J Microbiol Meth* 2004;**57**:399–407.
- Lynn TM, Ge TD, Yuan HZ et al. Soil carbon-fixation rates and associated bacterial diversity and abundance in three natural ecosystems. *Microb Ecol* 2017;**73**:1–13.
- Malik MA, Marschner P, Khan KS. Addition of organic and inorganic P sources to soil—Effects on P pools and microorganisms. *Soil Biol Biochem* 2012;**49**:106–13.
- Murphy J, Riley JP. A modified single solution method for the determination of phosphate in natural waters. *Anal Chimica Acta* 1962;**27**:31–36.
- Nannipieri P, Giagnoni L, Landi L et al. Role of phosphatase enzymes in soil. In: Bunemann EK, Oberson A, Frossard E et al. (eds). *Phosphorus in Action*. Soil Biology. Berlin, Heidelberg: Springer Verlag, 2011;**26**:215–41.
- Pathan SI, Ceccherini MT, Hansen MA et al. Maize lines with different nitrogen use efficiency select bacterial communities with different β -glucosidase-encoding genes and glucosidase activity in the rhizosphere. *Biol Fert Soils* 2015;**51**:995–1004.
- Richardson AE, Barea J, McNeill AM et al. Acquisition of phosphorus and nitrogen in the rhizosphere and plant growth promotion by microorganisms. *Plant Soil* 2009;**321**:305–39.
- Rodríguez H, Fraga R. Phosphate solubilizing bacteria and their role in plant growth promotion. *Biotechnol Adv* 1999;**17**:319–39.
- Saleque M, Naher U, Islam A et al. Inorganic and organic phosphorus fertilizer effects on the phosphorus fractionation in wetland rice soils. *Soil Sci Soc Am J* 2004;**68**:1635–44.
- Shen J, Li R, Zhang F et al. Crop yields, soil fertility and phosphorus fractions in response to long-term fertilization under the rice monoculture system on a calcareous soil. *Field Crop Res* 2004;**86**:225–38.
- Somaweera KATN, Suriyagoda LDB, Sirisena DN et al. Accumulation and partitioning of biomass, nitrogen, phosphorus and potassium among different tissues during the life cycle of rice grown under different water management regimes. *Plant Soil* 2016;**401**:169–83.
- Stewart JWB, Tiessen H. Dynamics of soil organic phosphorus. *Biogeochemistry* 1987;**4**:41–60.
- Su JQ, Ding LJ, Xue K et al. Long-term balanced fertilization increases the soil microbial functional diversity in a phosphorus-limited paddy soil. *Mol Ecol* 2015;**24**:136–50.
- Suriyagoda L, De Costa WAJM, Lambers H. Growth and phosphorus nutrition of rice when inorganic fertiliser application is partly replaced by straw under varying moisture availability in sandy and clay soils. *Plant Soil* 2014;**384**:53–68.
- Tabita FR. Microbial ribulose 1,5-bisphosphate carboxylase/oxygenase: a different perspective. *Photosynth Res* 1999;**60**:1–28.
- Tan H, Barret M, Mooij MJ et al. Long-term phosphorus fertilisation increased the diversity of the total bacterial community and the *phoD* phosphorus mineraliser group in pasture soils. *Biol Fert Soils* 2013;**49**:661–72.
- Tian J, Dippold M, Pausch J et al. Microbial response to rhizodeposition depending on water regimes in paddy soils. *Soil Biol Biochem* 2013a;**65**:195–203.
- Tian J, Pausch J, Fan MS et al. Allocation and dynamics of assimilated carbon in rice-soil system depending on water management. *Plant Soil* 2013b;**363**:273–85.
- Tolli J, King GM. Diversity and structure of bacterial chemolithotrophic communities in pine forest and agroecosystem soils. *Appl Environ Microbiol* 2005;**71**:8411–8.
- Torrent J. Interactions between phosphate and iron oxide. In: Auerswald K, Stanjek H, Bigham JM (eds). *Soils and Environment. Advances in Geocology*. Reiskirchen, Germany: Catena Verlag, 1997;**30**:321–44.
- Van Geel M, De Beenhouwer M, Ceulemans T et al. Application of slow-release phosphorus fertilizers increases arbuscular mycorrhizal fungal diversity in the roots of apple trees. *Plant Soil* 2016;**402**:291–301.
- Wang SQ, Zhao X, Xing GX et al. Phosphorus pool in paddy soil and scientific fertilization in typical areas of Taihu Lake Watershed. *Soils* 2012;**44**:158–62.
- Wang Y, Zhao X, Wang L et al. A five-year P fertilization pot trial for wheat only in a rice-wheat rotation of Chinese paddy soil: interaction of P availability and microorganism. *Plant Soil* 2016;**399**:305–18.
- Wei XM, Hu YJ, Peng PQ et al. Effect of P stoichiometry on the abundance of nitrogen-cycle genes in phosphorus-limited paddy soil. *Biol Fert Soils* 2017;**7**:767–76.
- Wood HG, Ragsdale SW, Pezacka E. The acetyl-CoA pathway: a newly discovered pathway of autotrophic growth. *Trends Biochem Sci* 1986;**11**:14–18.

- Wu JS, Joergensen RG, Pommerening B et al. Measurement of soil microbial biomass C by fumigation-extraction—an automated procedure. *Soil Biol Biochem* 1990;**22**:1167–9.
- Wu XH, Ge TD, Wang W et al. Cropping systems modulate the rate and magnitude of soil microbial autotrophic CO₂ fixation in soil. *Front Microbiol* 2015;**6**:237–46.
- Xiao KQ, Nie SA, Bao P et al. Rhizosphere effect has no effect on marker genes related to autotrophic CO₂ fixation in paddy soils? *J Soils Sediment* 2014;**14**:1082–7.
- Yakimov MM, Cono VL, Denaro R. A first insight into the occurrence and expression of functional *amoA* and *accA* genes of autotrophic and ammonia-oxidizing bathypelagic *Crenarchaeota* of Tyrrhenian Sea. *Deep-Sea Res Part II: Topical Studies in Oceanography* 2009;**56**:748–54.
- Yan C, Zhan HQ, Yan SS et al. Effects of straw retention and phosphorous fertilizer application on available phosphorus content in the soil solution during rice growth. *Paddy Water Environ* 2016;**14**:61–69.
- Yousuf B, Keshri J, Mishra A et al. Application of targeted metagenomics to explore abundance and diversity of CO₂-fixing bacterial community using *cbbL* gene from the rhizosphere of *Arachis hypogaea*. *Gene* 2013;**506**:18–24.
- Yuan HZ, Ge TD, Chen CY et al. Significant role for microbial autotrophy in the sequestration of soil carbon. *Appl Environ Microbiol* 2012a;**78**:2328–36.
- Yuan HZ, Ge TD, Wu XH et al. Long-term field fertilization alters the diversity of autotrophic bacteria based on the ribulose-1,5-biphosphate carboxylase/oxygenase (RubisCO) large-subunit genes in paddy soil. *Appl Environ Biotechnol* 2012b;**95**:1061–71.
- Yuan HZ, Zhu ZK, Liu SL et al. Microbial utilization of rice root exudates: ¹³C labeling and PLFA composition. *Biol Fert Soils* 2016;**52**:615–27.
- Zhong WH, Cai ZC. Long-term effects of inorganic fertilizers on microbial biomass and community functional diversity in a paddy soil derived from quaternary red clay. *Appl Soil Ecol* 2007;**36**:84–91.
- Zhu S, Vivanco JM, Manter DK. Nitrogen fertilizer rate affects root exudation, the rhizosphere microbiome and nitrogen-use-efficiency of maize. *Appl Soil Ecol* 2016;**107**:324–33.