

Chapter 9

Role of Phosphatase Enzymes in Soil

P. Nannipieri, L. Giagnoni, L. Landi, and G. Renella

9.1 Introduction

Phosphatases have been extensively studied in soil, as shown by some reviews (Ramirez-Martinez 1968; Speir and Ross 1978; Malcom 1983; Tabatabai 1994), because they catalyse the hydrolysis of ester-phosphate bonds, leading to the release of phosphate (P), which can be taken up by plants or microorganisms (Cosgrove 1967; Halstead and McKercher 1975; Quiquampoix and Mousain 2005). It has been shown that the activities of phosphatases (like those of many hydrolases) depend on several factors such as soil properties, soil organism interactions, plant cover, leachate inputs and the presence of inhibitors and activators (Speir and Ross 1978).

Phosphatases are enzymes catalysing the hydrolysis of both esters and anhydrides of phosphoric acid (Schmidt and Laskowski 1961) and, according to the Nomenclature Committee of the International Union of Biochemistry and Molecular Biology, they can be classified as phosphoric monoester hydrolases or phosphomonoesterases (EC 3.1.3), phosphoric diester hydrolases or phosphodiesterases (EC 3.1.4), triphosphoric monoester hydrolases (EC 3.1.5) and enzymes acting on phosphoryl-containing anhydrides (EC 3.6.1) and on P–N bonds (EC 3.9). Phosphatases can also be subdivided according to their regulation (e.g. calmodulin), the requirements of metal cations for their activity (e.g. Mg^{2+} and Ca^{2+}) and their sensitivity to various phosphatase inhibitors. Phosphomonoesterases include acid and alkaline phosphomonoesterase (which hydrolyse monoester bonds including mononucleotides and sugar phosphates), phosphoprotein phosphatases (which hydrolyse phosphoester bonds of phosphoserines, phosphothreonines or phosphotyrosines), phytases (EC 3.1.3.26 for 4-phytase and EC 3.1.3.8 for 3-phytase, which hydrolyse all six phosphate groups from inositol hexaphosphate) and nucleotidases. Acid and alkaline phosphomonoesterases do not hydrolyse phosphates of phytic

P. Nannipieri (✉), L. Giagnoni, L. Landi, and G. Renella
Department of Plant, Soil and Environmental Sciences, University of Firenze, Piazzale delle
Cascine 18, 50144 Firenze, Italy
e-mail: paolo.nannipieri@unifi.it

acid (*myo*-inositol hexaphosphates) but they can hydrolyse lower-order inositol phosphates (Cosgrove 1980). Phosphodiesterases hydrolyse one or two ester bonds in phosphodiester compounds and include nucleases, which catalyse the hydrolysis of phosphodiester bonds of nucleic acids to produce nucleotide units or mononucleotides but not inorganic phosphates. Phospholipases hydrolyse phospholipids. We shall also discuss inorganic pyrophosphatase (pyrophosphate phosphohydrolases, EC 3.6.1.1), the enzyme that hydrolyses pyrophosphate to inorganic P, because pyrophosphate can be used as a fertilizer (Dick and Tabatabai 1978).

The aim of this review is to discuss the role of phosphatases in P mineralisation in soil and the response of these enzyme activities to changes in environmental factors, agricultural management and pollution. Particular attention will be given to phosphomonoesterase activities, which have been studied most among soil phosphatases. The meaning of measuring phosphatase activities and the drawbacks of the current protocols for enzyme assays have a central role because we think that a better understanding of the role of phosphatases (like that of any enzyme activity in soil) depends on improvement of the present enzyme assays by separating the contribution of extracellular stabilised phosphatase activities from the contribution of activities of phosphatases associated with active microbial cells. The effects of organic amendments, fertilizers and pollutants will be discussed by considering the present drawbacks of the currently used enzyme assays rather than listing all reports on the subject and underlining the contradictory data. We suggest that the reader also considers the review by Speir and Ross (1978) because it discusses the first reports and includes an extensive bibliography of the 1950s, 1960s and part of the 1970s on the effects of sterilisation, air drying, storage of soil samples before measurements, pH, temperature, soil properties, soil depth, fertilizers, trace elements, activators and inhibitors. We also refer to the reviews by Malcom (1983) and Tabatabai (1994) for a more detailed discussion of the analytical problems and for the state-of-the-art on kinetic properties and the effects of inhibitors, activators and soil properties on phosphatase activities.

9.2 Determination of Soil Phosphatase Activities

Activities of soil phosphomonoesterases have been the most studied, although phospholipids and nucleic acids, whose degradation is catalysed by phosphodiesterases, are among the major sources of fresh organic P inputs to soil (Cosgrove 1967). Before the advent of the simple, accurate and rapid enzyme assay based on the use of *p*-nitrophenyl phosphate (pNPP) by Tabatabai and Bremner (1969), phosphatase assays used natural substrates such as β -glycerolphosphate and nucleic acids (Speir and Ross 1978; Malcom 1983; Tabatabai 1994). The use of artificial substrates began in the early 1960s with phenyl phosphate (Hofmann 1963), phenolphthalein phosphate (Dubovenko 1964; Geller and Ginzburg 1979), pNPP (Bertrand and de Wolf 1968), α -naphthyl phosphate (Hochstein 1962) and β -naphthyl phosphate (Ramirez-Martinez and McLaren 1966). The choice of artificial

substrates eliminated the determination of released phosphate, which is easily adsorbed by soil particles (Tabatabai 1994). The success of the pNPP assay stems also from the fact that hydrolysis of pNPP is much more rapid than that of natural substrates such as nucleic acids. The pNPP is hydrolysed to *p*-nitrophenol (pNP), which is usually determined spectrophotometrically at 400 nm under alkaline conditions. Soluble organic compounds can interfere with the quantification of the pNP (Vuorinen and Saharinen 1996). For this reason, Gerritse and van Dijk (1978) suggested the separation of pNP from pNPP and other soluble organic compounds, extracted from organic soils or animal wastes, by high pressure liquid chromatography on a cellulose column. They also observed a marked reduction of both acid and alkaline phosphomonoesterase activity by phosphate concentrations greater than 0.1 mM and therefore suggested using pNPP concentrations of 0.01–0.1 mM in the enzyme assay.

Because the phosphate group in pNPP is attached to the aromatic chromophore, hydrolysis may not reflect the activity of alkyl phosphomonoesterases. To determine this enzyme activity in soil, Avidov et al. (1993) proposed an assay based on the hydrolysis of 4-(*p*-nitrophenoxy)-1,2-butanediol phosphate with successive oxidation of the reaction product 4-(*p*-nitrophenoxy)-1,2-butanediol to pNP.

The hydrolysis of 4-methylumbelliferyl phosphate (MUP) to 4-methylumbelliferone (MU) has also been used to assay phosphomonoesterase activity in soil by determining the fluorescence of the MU. This assay circumvents interferences by soluble organic compounds because fluorescence is measured for emission wavelengths after specific excitation (Marx et al. 2001). The MUP assay gave higher values than the pNPP assay, but the two enzyme activities (measured in modified universal buffer adjusted to the soil pH value) were significantly correlated ($P < 0.001$) when expressed on the basis of C content but not when expressed on the basis of dry soil weight (Drouillon and Merckx 2005). The MUP assay gave lower K_m values than the pNPP assay (Table 9.1) (Marx et al. 2001) and this may suggest that the former substrate mimics the hydrolysis of naturally occurring soil organic phosphate esters more closely (Freeman et al. 1995).

The phosphodiesterase assay is similar to the phosphomonoesterase assays because it is based on the release of pNP from bis-*p*-nitrophenyl phosphate (bpNPP) when the soil slurry is incubated with the substrate at pH 8.0 for 1 h (Browman and Tabatabai 1978). The bpNPP was first used by Ishii and Hayano (1974). Ohmura and Hayano (1986) showed that the optimum pH of phosphodiesterase activity of 15 soils ranged from 4.5 to 9.5, a broader pH optimum than that suggested in the assay by Browman and Tabatabai (1978). In addition, the enzyme activity was significantly correlated with soil pH.

Phosphotriesterase activity of soil has been determined by hydrolysis of tris-*p*-nitrophenyl phosphate, which is insoluble in water, to pNP (Eivazi and Tabatabai 1977).

According to Turner et al. (2002a, b), soil phytase has been poorly studied because it has been determined by the release of phosphate from phytate and not by using suitable artificial substrates (Yadav and Tarafdar 2003). Berry et al. (2007) proposed measuring the phytase activity of soil by using a chromophoric substrate

Table 9.1 Some K_m values of phosphatases

Enzyme	K_m (mM)	Substrate concentration (mM)	Temperature (°C)	Buffer	pH	References
Acid phosphomonoesterase	0.94–1.75	1–20 ^a	37	MUB	6.5	Tabatabai and Bremner (1971)
Acid phosphomonoesterase	0.35–5.40 ^b	–	37	Acetate	4.7	Cervelli et al. (1973)
Acid phosphomonoesterase	1.11–3.40	1–20 ^a	37	MUB	6.5	Eivazi and Tabatabai (1977)
Acid phosphomonoesterase	0.1	0.05–0.50	30	Acetate	5.0	Gerritse and van Dijk (1978)
Acid phosphomonoesterase	1.71–6.99 ^b	3.2–23.0 ^b	30	MUB	5–6	Trasar-Cepeda and Gil-Sotres (1988)
Alkaline phosphomonoesterase	0.7	0.05–0.50	30	Tris	8.0	Gerritse and van Dijk (1978)
Alkaline phosphomonoesterase	0.44–4.94	1–20 ^a	37	MUB	11	Eivazi and Tabatabai (1977)
Phosphodiesterase	0.25–1.25	1–15 ^a	37	MUB	10	Eivazi and Tabatabai (1977)
Pyrophosphatase	21–51 ^c	10–60 ^a	37	MUB	–	Dick and Tabatabai (1978)

MUB modified universal buffer

Substrates were *p*-nitrophenyl phosphate for phosphomonoesterases, bis-*p*-nitrophenyl phosphate for phosphodiesterases, and pyrophosphate for pyrophosphatase

^aSoil solution bases

^bCorrected for the adsorption of the substrate

^cLineweaver–Burk plot

analogue of phytic acid whose disappearance can be monitored by high-performance liquid chromatography with UV detection. However, the method has not yet been set up for determining enzyme activity in soil.

Dick and Tabatabai (1977, 1978) set up an accurate method for determining inorganic pyrophosphatase activity of soil at pH 8.0 using pyrophosphate as the substrate and an improved determination of the released phosphate. This enzyme activity can be important from an agricultural point of view because pyrophosphate is a fertilizer P. According to Dick and Tabatabai (1978), the previous assays presented various drawbacks such as the adsorption of enzymatically released inorganic P by soil particles, hydrolysis of pyrophosphate to inorganic P after extraction from soil due to other reactions than that catalysed by pyrophosphatase, and interference of pyrophosphate on the determination of inorganic P.

The hydrolysis of polyphosphates in soil has been determined by Dick and Tabatabai (1986). One of the polyphosphates used in agriculture is trimetaphosphate, a cyclic polyphosphate (Busman and Tabatabai 1985). The assay for determining trimetaphosphatase (trimetaphosphate hydrolase, EC 3.6.1.2) activity was set up by Busman and Tabatabai (1985). It involves the incubation of soil with trimetaphosphate at pH 8.0 for 5 h, followed by precipitation of residual trimetaphosphate, pyrophosphate and triphosphate. Phosphate is not precipitated and can then be determined. Trimetaphosphate is hydrolysed by trimetaphosphatase to triphosphate, which is then hydrolysed by triphosphatases to pyrophosphate and phosphate (Tabatabai 1994). Finally, pyrophosphate is hydrolysed to phosphate by pyrophosphatase. Therefore, the interpretation of the data obtained by this assay is complicated by the fact that the enzyme assay measures the activity of three enzymes, trimetaphosphatase, triphosphatase and pyrophosphatase (Tabatabai 1994).

9.3 Range and Kinetic Properties

Table 9.2 shows the range of phosphatase activities measured in soil with current assay procedures. Acid phosphomonoesterase activities in soil have been frequently measured at pH 6.5; however, at this pH the measured enzyme activity may include acid and alkaline phosphomonoesterase activity (Malcom 1983). Acid phosphomonoesterase activity generally prevails in acidic soils, whereas alkaline phosphomonoesterase activity prevails in alkaline soils, and for this reason the activities of the two enzymes are negatively correlated (Juma and Tabatabai 1978). Pang and Kolenko (1986) found a pH optimum of 7.0 for phosphomonoesterase activity in two forest soils. In comparing phosphatase activities (as for any other enzyme activity in soil) it is important to consider the period of the year in which soil sampling is done because enzyme activities of soil can change throughout the year (Schneider et al. 2001). Grierson and Adams (2000) observed that acid phosphomonoesterase activity of Jarrah (*Eucalyptus marginata* Donn ex Sm) forest soils

Table 9.2. Range of measured phosphatase activities

Enzyme	Substrate	Substrate concentration (mM)	Temperature (°C)	Duration of assay (h)	Buffer	pH	Range of enzyme activity ($\mu\text{mol product g}^{-1} \text{h}^{-1}$)	References
Phosphomonoesterase	pNPP	50	20	1	MUB	At soil pH of 3.2–8.1	0.05–5.22	Drouillon and Merckx (2005)
Phosphomonoesterase	MUP	16 and 25	20	0.25	MUB	At soil pH of 3.2–8.1	0.48–10.41	Drouillon and Merckx (2005)
Acid Phosphomonoesterase	pNPP	10	37	0.5	0.5 M Tris maleate	6.5	10.4–307	Turner et al. (2002b)
Acid Phosphomonoesterase	pNPP	5	37	1	0.5 M Tris maleate	6.5	2.62–12.19	Turner and Haygarth (2005)
Acid Phosphomonoesterase	–	25	37	1	MUB	6.5	1.03–10.38 in air dried soils; 2.11–27.07 in moist soils	Baligar et al. (1988)
Acid phosphomonoesterase	MUP	0.01–0.40	–	–	Water	At soil pH of 4.0–4.3	0.57–1.08	Santruckova et al. (2004)
Acid phosphomonoesterase	pNPP	50	37	1	MUB	6.5	0.85–14.9	Dick et al. (1988)
Acid phosphomonoesterase	pNPP	50	37	1	MUB	6.5	0.31–3.15	Zornoza et al. (2009)
Acid phosphomonoesterase	pNPP	5	37	1	MUB	6.5	0.35–0.88	Eivazi and Tabatabai (1977)
Acid phosphomonoesterase	–	115	37	1	MUB	6.5	0.06–0.13	Ho (1979)
Alkaline phosphomonoesterase	pNPP	5	37	1	MUB	11	0.06–1.60	Eivazi and Tabatabai (1977)
Alkaline phosphomonoesterase	pNPP	50	37	1	MUB	11	0.34–5.50	Dick et al. (1988)
Phosphodiesterase	bpNPP	5	37	1	MUB	9–11	0.10–0.55	Eivazi and Tabatabai (1977)
Phosphotriesterase	tpNPP (insoluble)	5	37	1	MUB	10	0.01–0.08	Eivazi and Tabatabai (1977)

pNPP *p*-nitrophenyl phosphate, *MUP* 4-methyl umbelliferyl phosphate, *MUB* modified universal buffer, *bpNPP* bis-*p*-nitrophenyl phosphate, *tpNPP* tris-*p*-nitrophenyl phosphate

ranged from 30 to 40 $\mu\text{mol pNP g}^{-1} \text{h}^{-1}$ in winter and spring when soil was moist, whereas it was below 10 $\mu\text{mol pNP g}^{-1} \text{h}^{-1}$ in the dry summer.

Activities of phosphodiesterases are lower than acid and alkaline phosphomonoesterase activities (Criquet et al. 2007) because the production of P monoesters from P diesters may stimulate the microbial synthesis of phosphomonoesterases (Turner and Haygarth 2005). It is reasonable to hypothesise that phosphodiesterase and phosphomonoesterase activities act sequentially in soil (Fig. 9.1). Phosphotriesterase activity was also lower than acid and alkaline phosphomonoesterase activities of soil (Eivazi and Tabatabai 1977).

Table 9.1 shows the K_m (the Michaelis–Menten constant) values of phosphatases in soil. Although phosphatases, like other hydrolases in soil, can derive from different sources and thus have different kinetic constants, the K_m value of phosphatase activity of a soil can be calculated. As discussed by Nannipieri and Gianfreda (1998), the calculated values probably represent a weighted average of the various constants of enzymes involved in the measured enzyme activity, with an unknown weighting factor. However, in the case of acid phosphomonoesterases, at least two enzymes with markedly different K_m values were found in pyrophosphate extracts from two soils by applying the Eadie–Scatchard plot (rate of reaction V vs. the substrate concentration S) (Nannipieri et al. 1982).

Brams and McLaren (1974) observed a marked deviation from linearity at higher substrate concentration for soil phosphomonoesterase (pH 6.90), and Irving and Cosgrove (1976) suggested that diffusional effects and adsorption of substrate by soil colloids were responsible for the fact that acid phosphomonoesterase of a Krasnozem did not follow Michaelis–Menten kinetics. Cervelli et al. (1973) proposed calculating the K_m value by considering the adsorption of the substrate (pNPP) by the Freundlich law; the corrected K_m value of acid phosphomonoesterase was lower than




Fig. 9.1 Mechanisms of phosphodiesterase and alkaline phosphomonoesterase reaction involving molecular rearrangement (*Reaction A*) and incorporation of oxygen atoms (●) in the phosphate molecule (*Reactions B and C*). Redrawn from Blake et al. (2005)

the uncorrected value. The same was observed for the K_m values of acid phosphomonoesterase in acid and organic soils from Galicia, Spain (Trasar-Cepeda and Gil-Sotres 1988). The shaking of soil slurries can accelerate the diffusion of the substrate towards the enzymes. Indeed, K_m values of phosphomonoesterases measured at pH 6.9 were 2.5 times greater when measured in soil columns than when measured in a batch-type system with shaking (Brams and McLaren 1974).

9.4 Limitations of the Present Enzyme Assays

Although the drawbacks of the currently used enzyme assays have been extensively discussed (Skujins 1978; Burns 1978, 1982; Nannipieri 1994; Tabatabai 1994; Nannipieri et al. 2002; Gianfreda and Ruggiero 2006), they are still frequently neglected when soil enzyme activities are interpreted. Firstly, the present enzyme assays measure potential and not real enzyme activities (Fig. 9.2) because the assay conditions are different (optimal pH, optimal temperature, substrate present at saturating concentration, presence of buffer to control pH during the assay, soil slurry, shaking) from those occurring in situ (fluctuations of temperature and moisture of the soil; pH and substrate concentration are rarely at the optimum for enzyme activity, etc.). Secondly, we do not know which enzymes contributed to the




Fig. 9.2 Contribution of activities of phosphomonoesterase differently located in the soil matrix to the measured enzyme activity, and drawbacks of the currently used assays

measured enzyme activity. According to Burns (1982), enzymes catalysing the measured reaction can be:

1. Associated with active microbial cells, either intracellular or attached to the outer cell surface
2. Associated with cell debris or dead cells
3. Associated with resting cells, such as bacterial spores
4. Released as truly extracellular enzymes to degrade high molecular weight or insoluble substrates
5. Present as extracellular enzymes of enzyme–substrates complexes
6. Present as free extracellular enzymes
7. Present as extracellular enzymes stabilised by their association with surface-reactive particles (e.g. clay minerals, iron oxides and hydroxides)
8. Entrapped by humic matter (the humus–enzyme complexes)

The activity of enzyme-like catalysts is probably significant under extreme environmental conditions where these catalysts are present, whereas activities of free extracellular enzymes (6), enzymes associated with substrates (5), and enzymes of cell debris and dead cells (2) are probably short-lived because they can be rapidly degraded unless they are adsorbed by soil particles. The contribution of enzymes associated with resting cells is probably insignificant (Nannipieri et al. 2002). Therefore, it is reasonable to hypothesise that the measured enzyme activity depends on the activity of enzymes associated with active microbial cells, including enzyme activities of plant cells in the rhizosphere and detritosphere soil, and on the activity of extracellular enzymes stabilised by soil colloids (Fig. 9.2) (Nannipieri 1994; Nannipieri et al. 2002). The extracellular stabilised enzyme activity is not affected by changes in composition, abundance or activity of the soil microflora; thus the intracellular enzyme activity of active cells should be used as an indicator of nutrient dynamics and changes in soil functioning due to agricultural management and ecological factors because it is well established that microbial activities of soil are more sensitive to these changes than other soil properties (Nannipieri et al. 2003). However, the separation of stabilised extracellular enzyme activity and enzyme activity associated with active microbial cells and plant cells is not possible with the present enzyme assays (Nannipieri et al. 2002). Most reports on soil enzymes assume that the present short assays only determine extracellular and stabilised enzyme activity. Despite this assumption, the measured enzyme activities are often taken as indicators of soil quality, which is strictly related to microbial activity and thus to intracellular enzyme activity. In addition, it is often assumed that changes in enzyme activities only reflect the response of microbiota to environmental factors, neglecting the fact that the measured enzyme activity also depends on the activity of stabilised extracellular enzymes.

Microbial inhibitors such as toluene have been used to inhibit the enzyme activity associated with active microbial cells, but this can create artefacts; for example, toluene can increase the permeability of cell membranes and thus the access of the urea substrate to intracellular ureases (Nannipieri et al. 2002). In addition, microorganisms can also use toluene as a substrate (Kaplan and

Hartenstein 1979). Toluene did not affect phosphodiesterase or acid and alkaline phosphomonoesterase activities but increased the phosphotriesterase activity of soil (Eivazi and Tabatabai 1977).

The so-called physiological response method is based on the measurements of enzyme activities and microbial biomass of soil during the period when microbial growth is stimulated by adding glucose and a nitrogen source to soil (Nannipieri et al. 2002). If enzyme activity is plotted against biomass, there is generally a significant and positive correlation between the enzyme activity (y -axis) and microbial biomass (x -axis). The extrapolation to zero of microbial biomass gives a positive intercept of the plot on the y -axis, which is the extracellular stabilised enzyme activity. This approach has been used to calculate extracellular acid phosphomonoesterase activity of a moist soil treated with different rates of sewage sludges, with measurement of microbial biomass by measuring ATP. Extracellular phosphomonoesterase activity of soil was 14.9, 5.3 and 4.3 $\mu\text{mol pNPP g}^{-1} \text{h}^{-1}$ after addition of 0, 50 and 100 tons of sewage sludge per hectare, respectively (Nannipieri et al. 1996a). This approach can only work for constitutive but not for inducible or repressible enzymes, such as phosphomonoesterase, whose synthesis is generally repressed by inorganic phosphate (Nannipieri 1994). Indeed, changes in inducible or repressible enzyme activities are not related to changes in microbial biomass. In addition, the percentage of glucose-utilising microorganisms depends on soil type, management and pollution (Nannipieri et al. 2002). Acid phosphomonoesterase activity of two eucalypt forest soils was significantly correlated with ergosterol content and microbial P when all these properties were measured throughout the year, the relative plots giving a positive intercept on the y -axis (Grierson and Adams 2000). Obviously, the intercept obtained with the first correlation cannot represent the extracellular acid phosphomonoesterase activity because ergosterol content only determines the fungal biomass. However, the approach by Grierson and Adams (2000) does not involve the stimulation of microbial growth by adding easily degradable organic compounds to soil, and thus does not present the above-mentioned drawbacks (Nannipieri et al. 2002).

Chloroform fumigation has also been used to distinguish enzyme activity associated with active microbial and plant cells from the extracellular enzyme activity stabilised in soil (Klose and Tabatabai 1999). This method assumes that the present short-term enzyme assays measure the stabilised extracellular enzyme activity and that the increase in enzyme activity after CHCl_3 fumigation is due to the intracellular enzyme activity. Therefore, the intracellular enzyme activity of soil can be calculated by subtracting the enzyme activity before fumigation from that after fumigation. As discussed by Nannipieri et al. (2002), this approach presents the following problems: (1) CHCl_3 fumigation does not kill all microbial cells and the efficiency of cell lysis depends on soil structure (Arnebrant and Schnurer 1990), and (2) the assumption that the present short-term enzyme assays determine only the extracellular enzyme activity has never been proven. The fact that enzyme activities, including acid and alkaline phosphomonoesterase activities, can increase with microbial biomass when easily degradable organic compounds, such as glucose, are added to soil (Nannipieri et al. 1978, 1979, 1983; Renella et al. 2006a, b, 2007b)

suggests that the present enzyme assays also measure the contribution of enzyme activities associated with active microbial cells of soil, and (3) proteases are active during the CHCl_3 fumigation period and degrade urease and both phosphomonoesterase enzymes (Renella et al. 2002); thus, protease activity needs to be inhibited during soil fumigation.

The use of sonication can increase the enzyme activity of soil. Indeed, the activity of acid phosphomonoesterase was 156% higher with soil sonication than without it (De Cesare et al. 2000). The increase probably depended on the release of extracellular enzymes stabilised by soil colloids and not on cell lysis, because the release of enzymes by sonication was not related to the release of ATP.

9.5 Role of Phosphatase in Organic P Mineralisation in Soil and the Effect of Inorganic P

As already mentioned, phosphodiesterase and phosphomonoesterase activities may act sequentially (Fig. 9.1). Pant and Warman (2000) observed that acid phosphomonoesterase (from wheat germ), alkaline phosphomonoesterase (from calf intestinal mucosa), phospholipase (from *Clostridium perfringens*) and nuclease (from *Staphylococcus aureus*), all immobilised on positively charged supports, were able to mineralise (at pH 7.0) organic P extracted from different soils by water or NaOH. The activities of both phosphomonoesterases were generally increased when these enzymes were used with one of the two phosphodiesterases.

Soil acid phosphomonoesterase activity was higher at low inorganic P content of soil than at high content, and the enzyme activity of the low-P soil was significantly correlated with herbage yield, probably due to the importance of organic P mineralisation for plant P nutrition (Speir and Cowling 1991). Santruckova et al. (2004) found that higher enzymatic hydrolysis of organic P depended on the higher microbial P immobilisation but not on the higher mineralisation of organic P compounds.

Application of inorganic P can repress the synthesis of phosphomonoesterases in soil because it inhibits the expression of *PHO* genes (Oshima et al. 1996) and, indeed, phosphate inhibits the phosphatase activities of soil (Halstead 1964; Juma and Tabatabai 1977, 1978; Lima et al. 1996; Moscatelli et al. 2005; Nannipieri et al. 1978; Olander and Vitousek 2000; Spiers and McGill 1979). However, the absence of a response of phosphatase activities to P addition has also been reported. For example, the application of triple superphosphate to an oak soil in 1992 did not affect acid phosphomonoesterase activity of soil samples taken in 1993 and 1994 (Schneider et al. 2001). Addition of phosphate with glucose and inorganic N did not stimulate the phosphomonoesterase activity (pH 6.5) of soil, whereas the stimulation occurred in the respective soil treated only with glucose and inorganic N (Nannipieri et al. 1978). Presumably, the enzyme activity was not decreased by phosphate due to the presence of extracellular phosphomonoesterases stabilised by

soil colloids or due to the presence of constitutive microbial phosphomonoesterase in soil. Enzyme assays discriminating the activities of extracellular stabilised enzymes from activities of enzymes associated with soil microorganisms would permit an understanding of the underlying mechanisms (Fig. 9.2).

9.6 Phosphatase Activities of Bulk and Rhizosphere Soil and the Origin of Phosphatases in Soil

It is well established that enzyme activities are higher in rhizosphere than bulk soil (Skujins 1978; Tarafdar and Chhonkar 1978; Dinkelaker and Marschner 1992). Both acid and alkaline phosphomonoesterase activities of soil were increased near the rhizoplane of *Brassica oleracea*, *Allium cepa*, *Triticum aestivum* and *Trifolium alexandrinum* and such an increase depended on plant species, soil type and plant age (Tarafdar and Jungk 1987). Probably, the increase with plant age was due to the gradual formation of the rhizosphere microflora and to the release of plant phosphomonoesterases. The distance from the rhizoplane at which the rhizosphere effect on enzyme activities was observed was higher for acid (from 2 to 3.1 mm) than for alkaline (from 1.2 to 1.6 mm) phosphomonoesterase. There was an inverse and significant correlation between the acid or the alkaline phosphomonoesterase activity and the content of organic P of the rhizosphere soil sampled from *Triticum aestivum* and *Trifolium alexandrinum*, whereas the content of inorganic P increased towards the rhizoplane. Increases in both acid and alkaline phosphomonoesterase activities near the rhizoplane of maize were accompanied by changes in the composition of bacterial communities as determined by PCR-DGGE (polymerase chain reaction-denaturing gradient gel electrophoresis) (Kandeler et al. 2002). Some agroforestry species (*Tithonia diversifolia*, *Tephrosia vogelii* and *Crotalaria grahamiana*) stimulated acid phosphomonoesterase activity of rhizosphere soil, whereas maize stimulated alkaline phosphomonoesterase activity of rhizosphere soil (George et al. 2002). Higher soil phosphomonoesterase activities were found under invader plant species than under grass and forbs, presumably due to the higher P uptake of the invading plants (Neal 1973). Izaguirre-Mayoral et al. (2002) found that nodulated legumes species growing in acid savanna soils stimulated the acid phosphomonoesterase activity of rhizosphere soil.

Interactions between soil microorganisms and plant species can also affect phosphomonoesterase activities of rhizosphere soil. It has been suggested that arbuscular mycorrhizal (AM) fungi stimulate the release of acid phosphomonoesterase from roots of subterranean clover (*Trifolium subterraneum* L.) (Joner and Jakobsen 1995). However, mycorrhizal infection of cucumber (*Cucumis sativus* L.) did not affect acid and alkaline phosphomonoesterase activities of soil (Joner et al. 1995). More information on the relationship between phosphatases and mycorrhizae is given by Jansa et al. (2011).

Obviously, it is difficult to interpret the measurement of phosphomonoesterase activities of rhizosphere soil if the contribution of plant and microbial phosphatases to the measured enzyme activity are not separated. Colvan et al. (2001) suggested that acid phosphomonoesterase activity of hay meadow soils was due to enzyme released by plants, because the enzyme activity was high and microbial P was low in soils never treated with fertilizer or treated with N or K fertilizer for 100 years. However, the measured acid phosphomonoesterase activities could also have been at least partly due to enzymes synthesised by the soil microflora in response to P-deficient conditions (Nannipieri 1994). A significant correlation between acid phosphomonoesterase activity of an oak soil and fine root length density of oak has been found (Schneider et al. 2001). The response of activities of hydrolases, including acid and alkaline phosphomonoesterase and phosphodiesterase, in rhizosphere soil depends on the type of root exudate stimulating microbial activity (Renella et al. 2006b, 2007a), which suggests that active microbial phosphomonoesterases are probably the major contributing enzymes to the measured enzyme activity of the rhizosphere soil.

Phosphorus nutrition of transgenic *Nicotiana tabacum* expressing a chimeric phytase gene (*ex::ph A*) from soil fungus *Aspergillus niger* was better in neutral than in acid soils, because the adsorption by soil of phytase released from roots was lower at neutral than at acid pH values (George et al. 2005a, b). The expression of phytase in the transgenic plant had no effect on the structure of microbial communities of rhizosphere soil compared to the wild type (George et al. 2009).

No correlations have been found between acid phosphomonoesterase activity and acid phosphomonoesterase-active bacterial colonies, and between alkaline phosphomonoesterase activity and alkaline phosphomonoesterase-active bacterial colonies of forest soils (Hysek and Sarapatka 1998), but this is not surprising since culturable bacteria only represent 1–10% of the bacteria inhabiting soil (Torsvik et al. 1996). Acid phosphomonoesterase was secreted by roots of three cereals (wheat, pearl millet and sorghum), three legumes (mung bean, moth bean and cluster bean), and three oil seed crops (groundnut, sesame and mustard) when these plants were grown in culture solution under P-deficient conditions (Yadav and Tarafdar 2001). The secretion pattern depended on the plant species, and the amount of root exudates increased with plant age and was higher with phytin than with leucithin and glycerolphosphate, each present as the sole source of P. Because acid phosphomonoesterases (like any enzyme molecule) diffuse poorly into the soil matrix, plant-released enzymes probably mineralise organic P from sloughed off or damaged cells rather than from native soil organic P (Lefebvre et al. 1990; Yadav and Tarafdar 2001).

Alkaline phosphomonoesterase activity has not been detected in plants (Dick et al. 1983; Juma and Tabatabai 1988a, b, c) and for this reason microbial cells supposedly synthesise most of the soil alkaline phosphomonoesterases (Tabatabai 1994). Both soil bacteria and soil microorganisms other than AM fungi (Joner and Jakobsen 1995) are thought to contribute to the measured soil alkaline phosphomonoesterase activity. Indeed, alkaline phosphatase activity of both rhizosphere and bulk soil depend on the composition of bacterial communities harbouring alkaline phosphatase genes, as determined by DGGE after amplification of extracted DNA by polymerase chain

reaction with specific primers (Sakurai et al. 2008). Increases in acid and alkaline phosphomonoesterase activities due to the addition of ryegrass residues to soils were related to changes in the composition of bacterial communities, as determined by DGGE (Renella et al. 2006a). Changes in the ergosterol content of Jarrah forest soils explained 50% of the changes in acid phosphomonoesterase activity in each season, whereas the ergosterol content of soil sampled under *Banksia grandis*, an understorey dominant plant growing in dense thickets in the absence of fire, explained 74% of the changes in enzyme activity during the dry season (summer) but only 10% in moist soils (Grierson and Adams 2000). Both phytase- and phosphomonoesterase-producing fungi isolated from arid and semiarid soils of India belonged to the genera *Aspergillus*, *Emmericella* and *Penicillium* (Yadav and Tarafdar 2003). In conclusion, the present evidence confirms that bacteria are the main source of alkaline phosphomonoesterase activity in soil, whereas acid phosphomonoesterase and phytase can derive from plants, fungi and bacteria. However, studies such as that by Sakurai et al. (2008) involving detection of genes codifying these enzymes are necessary.

Renella et al. (2007b) estimated the production and persistence of acid and alkaline phosphomonoesterase and phosphodiesterase activities in soils with a wide range of properties, by stimulating microbial growth through the addition of glucose and inorganic N to soil. Phosphatase activities of the soil increased, with microbial biomass reaching a peak value, but then both declined on prolonging the incubation time. Enzyme production (Pr) was calculated by the relationship $Pr = H/t_H$, where H is the peak of the enzyme activity and t_H is the time of the peak after adding glucose plus N to soil. Enzyme persistence (Pe) was calculated by the relationship $Pe = (r/H)\Delta t$, where r indicates the residual enzyme activity at the end of the incubation time and Δt is the time interval between the peak value (H) and the residual activity (r). The Pr values of acid phosphomonoesterase activity were highest in soils under forest and set-aside management, whereas Pr values of alkaline phosphomonoesterase and phosphodiesterase activities were highest in alkaline and neutral soils. The Pe values of acid phosphomonoesterase activity were highest in acid soils, whereas no relationship was found between alkaline phosphomonoesterase or phosphodiesterase activity and the soil pH or management (Renella et al. 2007b).

Both phosphodiesterases and pyrophosphatases are ubiquitous in animal, plant and microbial cells because they are involved in the degradation of nucleic acids and in several basic metabolic pathways of cells, respectively (Browman and Tabatabai 1978; Cooperman et al. 1992; Tabatabai 1994).

9.7 Effects of Soil Handling, Soil Properties, Agricultural Management and Pollutants on Soil Phosphatase Activities

Here, we shall only discuss reports after the late 1970s because Speir and Ross (1978) have extensively reviewed the effects of soil sampling, handling and storage, soil

properties, different agricultural managements, forest practices and pollutants on phosphatase activities.

9.7.1 Effects of Soil Handling and Soil Properties on Phosphatase Activities

Both air-drying and freeze-drying often decrease acid and alkaline phosphomonoesterase activity of soil (Gerritse and van Dijk 1978; Baligar et al. 1988; Adams 1992). However, Eivazi and Tabatabai (1977) found an increase in acid phosphomonoesterase and phosphotriesterase activities and a decrease in alkaline phosphomonoesterase and phosphodiesterase activities after air-drying of soil. Acid phosphomonoesterase activities in moist soils stored at 4°C and in the respective air-dried soils were significantly correlated (Baligar et al. 1988). Air-drying also decreased pyrophosphatase activity of soil, and the best storage conditions were to keep field-moist soils at 5°C (Tabatabai and Dick 1979). Probably the best strategy is to keep moist soils at 4°C and measure the enzyme activity as soon as possible. Kandeler (2007) suggests that if the determination of the enzyme activity requires storage periods longer than 3 weeks at 4°C, it is better to store the samples at -20°C than at 4°C. At the end of the storage period, soil samples are allowed to thaw at 4°C for about 2 days before the determination of the enzyme activity.

Steam sterilisation at 121°C for 1 h completely inactivated alkaline phosphomonoesterase, phosphodiesterase and phosphotriesterase activity, but increased acid phosphomonoesterase activity (Eivazi and Tabatabai 1977). Heating above 60°C inactivated the pyrophosphatase activity of soil (Tabatabai and Dick 1979).

It is well established that phosphatase activities are correlated with the content of organic matter and decrease with soil depth (Speir and Ross 1978; Tabatabai and Dick 1979; Prado et al. 1982; Pang and Kolenko 1986; Tabatabai 1994). Factors affecting phosphatase activity, measured by using sodium phenyl phosphate as a substrate in different woodland soils, could be ranked as rock type = vegetation type > soil type = season > soil depth (Harrison 1983). Some 20% of the variation of both acid and alkaline phosphomonoesterase activities of semiarid woodland soils depended on soil microclimate and surface depth (0–10 cm); soil temperature together with soil water potential was a better predictor of phosphatase activities than either factor alone (Kramer and Green 2000).

Humic acids competitively inhibited wheat phytase activity measured at 55°C and pH 5.15 (Pereira 1971). For forest soils, phosphomonoesterase activity of litter (pH in water 3.6–3.7), humus (pH in water 3.4–3.5) and mineral layers (pH in water 4.0–4.3), measured with water and MUP as the substrate, were correlated with the contents of organic C, microbial N and microbial P, and with soil respiration (Joergensen and Scheu 1999). Santruckova et al. (2004) observed that changes in soil pH and contents of total organic C, total N, total P, oxalate-soluble reactive and organic P, as well as oxalate-soluble Al and Fe can affect phosphomonoesterase activity and microbial biomass P in various ways, leading either to a surplus or a

deficiency in available soil phosphate. The relationship between phosphomonoesterase activities and the distribution of P forms is not clear. Acid phosphomonoesterase and phosphodiesterase activities of Karri forest soils were not related to soil P fractions (non-occluded Fe- and Al-bound P, P sorbed by carbonates, occluded P, and Ca-bound P) or total P (Adams 1992).

Alkaline phosphomonoesterase activity, measured using disodium phenylphosphate as substrate, was mainly associated with silt and clay fractions of a Haplic Chernozem (Kandeler et al. 1999). By contrast, in a calcareous and in an acid soil, both alkaline and acid phosphomonoesterase activities were associated with larger soil fractions (100–2,000 μm particle diameter) containing plant debris and less humified organic matter and were characterised by the highest mineralisation of organic P among the soil fractions (Rojo et al. 1990).

9.7.2 Effect of Agricultural Management, Forest Practices and Fire on Soil Phosphatases

Hay meadow soils treated with farmyard manure for about 100 years had higher acid and alkaline phosphomonoesterase and phosphodiesterase activities and higher $\text{NH}_4\text{F-HCl}$ -extractable P than those receiving mineral P (Colvan et al. 2001). However, both phosphomonoesterases and phosphodiesterase activities were positively ($P < 0.1$) correlated with extractable P in soils treated with farmyard manure or with phosphate, whereas acid phosphomonoesterase activity was negatively ($P < 0.05$) correlated with extractable P. Acid phosphomonoesterase activity was negatively correlated with alkaline phosphomonoesterase ($P < 0.05$) and with phosphodiesterase ($P < 0.05$) activities when considering all treatments, but was always higher than the alkaline phosphomonoesterase activity because all soils were acidic.

Straw burning instead of straw soil incorporation decreased acid but not alkaline phosphomonoesterase activity of the soil (Dick et al. 1988). Sewage sludges applied to soils increased phosphodiesterase and acid and alkaline phosphomonoesterase activities, which subsequently decreased with time (Criquet et al. 2007). Long-term experiments have shown that repeated applications of manure increases both acid and alkaline phosphomonoesterase activities, particularly immediately after manure addition to the soil (Dick et al. 1988; Colvan et al. 2001), due to the stimulation of microbial growth. When the monitoring period is prolonged, stimulation of microbial synthesis of enzymes by easily degradable organic substrates decreases (Garcia et al. 1993; Nannipieri 1994). However, the interpretation of changes in enzyme activity of soils treated with organic materials is difficult because these materials add exogenous enzymes associated with microorganisms and extracellularly stabilised enzymes to the soil.

No-till systems usually have higher enzyme activities in the surface soils than tilled soils because of the increase in soil organic matter content (Nannipieri 1994). However, this did not occur in an organic soil (Bergstrom et al. 1998).

In forest soils, fertilisation with urea or phosphate reduced acid phosphomonoesterase activity (Pang and Kolenko 1986). Fumigation of forest soils with methylbromide and chloropicrin for 24 h decreased phosphomonoesterase activity (pH 7.0), but there was a recovery when the fumigant was removed and the moist soils were incubated under controlled conditions (Pang and Kolenko 1986).

Controlled fire did not affect acid phosphomonoesterase activity of forest surface (0–5 cm) soil because the temperature never exceeded 50°C, whereas a marked reduction occurred with uncontrolled wildfires (Saa et al. 1993). Also, logging and/or burning operations almost eliminated acid phosphomonoesterase and phosphodiesterase activities of Karri forest soils (Adams 1992). Acid phosphomonoesterase activity did not recover after incubation of a wildfire-affected soil for 11 weeks under controlled conditions, probably because the high inorganic P repressed enzyme synthesis by soil microorganisms (Saa et al. 1998).

9.7.3 Effects of Pollutants on Soil Phosphatase Activities

Both acid and alkaline phosphomonoesterase activities have been monitored to evaluate the effects of several pollutants on organic P mineralisation in soil. The ecological dose (ED₅₀), i.e. the concentration of the pollutant that reduces the enzyme activity by 50%, has been calculated to quantify some of these effects.

Acid phosphomonoesterase activity of a blanket peat, an organic grassland soil and a calcareous grassland soil were high due to the P limitation induced by long-term atmospheric nitrogen deposition (Turner et al. 2002b), whereas sulfur pollution decreased the acid phosphomonoesterase activity of ectomycorrhizal roots in Norway spruce (Rejsek 1991).

The acid phosphomonoesterase activity of either moist or air-dried acid mesic fibrisols and histosols decreased after addition of copper to soil (Mathur and Rayment 1977; Mathur and Sanderson 1978). Juma and Tabatabai (1977) observed (depending on the soil type) the highest inhibition by Hg(II), As(V), W(VI) and Mo(VI) in the case of the acid phosphomonoesterase activity, and by Ag(I), Cd(II), V(IV) and As(V) in the case of the phosphomonoesterase activity. A negative and significant correlation was found between the sum of Cu and Zn total concentration in the soil and phosphomonoesterase activity determined using phenyl phosphate as the substrate (Tyler 1976).

Short-term laboratory incubations might not reflect the toxic effects in long-term heavy metal polluted soils. Alkaline phosphomonoesterase activity was still reduced in soils contaminated with Cd (concentration ranging from 0 to 0.36 nmol Cd kg⁻¹) in 1988–1990 and sampled in 2001, despite very low Cd availability, as determined either by water extraction or by the BIOMET bacterial biosensor system (Renella et al. 2004). In contrast, acid phosphomonoesterase activity and the composition of the bacterial community, determined either by plate counts or by DGGE, were unaffected, probably because the Cd pollution caused physiological adaptations rather than the selection of metal-resistant culturable bacteria. Addition of dry milled

ryegrass to these long-term Cd-contaminated soils increased both microbial biomass and acid and alkaline phosphomonoesterase activities (Renella et al. 2005a). However, the ratio of alkaline phosphomonoesterase activity to ATP decreased while that of acid phosphomonoesterase activity to ATP was unaffected compared to the respective uncontaminated soils. Both acid and alkaline phosphomonoesterase activities and the respective ratios of hydrolase to microbial biomass C were reduced in soils contaminated on the long term with Ni and Cd, but not in those with a Mn and Zn contamination; the former contamination also changed the composition of the bacterial community, as determined by DGGE (Renella et al. 2005b).

Dose–response curves can combine the effects of pollutants and soil physico-chemical properties on soil microflora (Babich et al. 1983) and can be used to calculate the effective ecological dose (ED_{50}) of enzyme activity of soil in response to pollution by heavy metals (Doelman and Haastra 1989). The ED_{50} values for Cd, Cu and Zn were 2.6 mmol kg⁻¹ soil in sandy soils and 45 mmol kg⁻¹ soil in clayey soils. Generally, the Cd toxicity was higher when observed 1.5 years after addition of heavy metal salts to soils than after 6 weeks. The presence of Cu or Zn increased the toxicity of Cd on acid and alkaline phosphomonoesterase activity of contrasting forest soils, as observed by comparing the relative ED_{50} values determined by the kinetic model (Renella et al. 2003). The toxicity was higher in sandy than in finer textured soils. The sensitivity of acid phosphomonoesterase activity was higher in alkaline than in acid or neutral soils, and the sensitivity of alkaline phosphomonoesterase activity showed an opposite behaviour (Renella et al. 2003).

Recovery of both acid and alkaline phosphomonoesterase activities occurred 7 years after the in situ remediation of soils contaminated with sludge-borne metals. Inorganic amendments were used, such as 5% (w/w) beringite (a coal fly ash) or 1% (w/w) zerovalent iron grid; both treatments reduced the heavy metal availability, whereas the composition of the bacterial community as determined by DGGE was not affected (Mench et al. 2006). The treatment of soils that were vegetated with a grass and herb mixture with alkaline fly ash and peat reduced leaching of Cu and Pb and increased the phosphodiesterase activity and the acid and alkaline phosphomonoesterase activities of the soil (Kumpiene et al. 2009). The treatment of an As-contaminated loamy sand soil with beringite (with or without zerovalent iron grid) reduced the extractable As, uptake of As by lettuce, and acid phosphomonoesterase activity because the treatment increased soil pH, whereas both alkaline phosphomonoesterase and phosphodiesterase activities were increased (Ascher et al. 2009). The composition of bacterial and fungal communities determined by DGGE both changed, with a decrease in microbial diversity induced by the treatments.

The interpretation of the effect of any pollutant on soil phosphatase activities (as for any soil enzyme activity) is problematic because of the limits of the enzyme assays currently used (as already discussed) and the presence of direct and indirect effects on the target enzyme (Nannipieri et al. 2002). For example, inhibition of enzyme activity by a pollutant may be masked by the growth of surviving microorganisms with expression of genes codifying the enzyme; the microbial growth can be caused by the use of microbial debris (derived from microbial cells killed by the pollutant) by the surviving microorganisms.

9.8 Stabilisation of Extracellular Phosphatases in Soil by Interaction with Surface-Reactive Particles or by Entrapment Within Humic Molecules

Three approaches have been followed to study the stabilisation of phosphatases in soil: (1) the use of pure enzymes to create model enzyme complexes, either with inorganic minerals such as clay, or with humus like materials, (2) the extraction and characterisation of phosphatases from soil, and (3) the visualisation of extracellular phosphatases in the soil matrix.

Enzymes, like any protein, are rapidly (within a few hours) adsorbed to clays and can be partially desorbed by washing the clay–protein complex; the molecules that cannot be desorbed by washing are referred to as “bound” proteins (Stotzky 1986; Nielsen et al. 2006). Protein adsorption depends on clay properties such as surface area, cation exchange capacity, charge density, type of saturating cation and degree of clay swelling (Stotzky 1986; Nannipieri et al. 1996b; Nielsen et al. 2006). The type of protein is also important because adsorption is generally maximal at pH values within the range of the protein’s isoelectric point and thus involves an ion-exchange mechanism. However, hydrogen bonding, van der Waals forces and hydrophobic effects are also involved in the adsorption of proteins by clay minerals.

Reduced enzyme activity after the adsorption can occur due to modification in the tertiary structure of the protein or due to reduced accessibility of the substrate to the active site. Adsorption of alkaline phosphomonoesterase by illite reduced the enzyme activity more than adsorption of the enzyme by montmorillonite or kaolinite (Makboul and Ottow 1979). Inhibition of enzyme upon adsorption may depend on its function. Intracellular phytase was completely inhibited when adsorbed by clays, whereas extracellular phytase retained its catalytic activity (Quiquampoix and Mousain 2005). Usually, the adsorption of the enzyme by clay minerals increases V_{\max} and K_m values (Nannipieri and Gianfreda 1998). Both values were increased when alkaline phosphomonoesterase was adsorbed by Ca-montmorillonite whereas they were decreased when the enzyme was adsorbed by Ca-illite (Makboul and Ottow 1979). The adsorption of acid phosphomonoesterase and pyrophosphatase by Ca-illite and Ca-montmorillonite did not affect the kinetic constants of enzymes (Dick and Tabatabai 1987). The protein adsorption by clays may also improve stability against thermal denaturation, wetting and drying cycles, and proteolysis; the resistance against proteolysis occurs if the protein penetrates the interlayer space of montmorillonite (Stotzky 1986; Nannipieri et al. 1996b; Nielsen et al. 2006). Indeed, acid phosphomonoesterase adsorbed on kaolinite was less resistant to proteolysis than that adsorbed on montmorillonite against proteolysis and thermal denaturation (Sarkar et al. 1989) and alkaline, whereas alkaline phosphomonoesterase adsorbed on Ca-illite was more resistant to proteolysis than the free enzyme (Makboul and Ottow 1979). Both acid and alkaline phosphomonoesterase bound to homo-ionic clays were degraded by soil microorganisms but degradation rates were higher with kaolinite complexes than with bentonite and vermiculate complexes, and higher with Ca-clays than with Al-clays (Chhonkar and Tarafdar 1985).

Acid phosphomonoesterase from sweet potato was more inhibited by tannic acid than urease and invertase (Rao et al. 1998).

Humus–enzyme complexes have been prepared by oxidative coupling of phenols in the presence of the enzyme to be immobilised (Nannipieri et al. 1996b). Acid phosphomonoesterase was immobilised in a resorcinol polymer synthesised by peroxidase; the enzyme was not linked to the resorcinol moiety by covalent bonds and it was more resistant to denaturation by pH, temperature and proteolysis than the free enzyme (Garzillo et al. 1996).

Acid phosphomonoesterase from potato adsorbed on Ca-polygalacturonate (a polymer simulating mucigel of the root–soil interface) by electrostatic interactions showed increased stability, but a decreased resistance against proteolytic and thermal denaturation (Marzadori et al. 1998).

Phosphatases have been extracted from soil using different solutions (Tabatabai and Fu 1992; Nannipieri et al. 1996b). Both acid and alkaline phosphomonoesterase were extracted by shaking litter with 1 M CaCl₂, 0.05% Tween 80 and polyvinylpolypyrrolidone, and the enzyme activities were determined after dialysis and concentration of the extract (Criquet et al. 2004). Soil moisture was the most important factor affecting the production of acid phosphomonoesterase. However, principal component analysis and multiple regressions showed that both temperature and the number of culturable heterotrophic bacteria also affected the dynamics of acid and alkaline phosphomonoesterase activities and organic P mineralisation.

Free extracellular alkaline phosphomonoesterase extracted from two soils by water was less resistant (i.e. had a lower inactivation temperature) and showed lower K_m values than the respective humic–enzyme complexes extracted by the chelating resin Chelex (Kandeler 1990).

Mayaudon (1986) suggested that several enzymes, including phosphomonoesterases and phosphodiesterases extracted from soil by phosphate-EDTA at pH 7–8, are fungal glycoenzymes associated with bacterial lipopolysaccharides, which are linked to humic compounds by Ca bridges. Humus–phosphomonoesterase complexes have been extracted from soil by pyrophosphate (Nannipieri et al. 1996b). Successive exhaustive ultrafiltration divided the soil extract into two fractions: one with molecular weights higher than 100,000 (A_I), and the other with molecular weights between 10,000 and 100,000 (A_{II}). Gel chromatography of the A_I fraction gave three peaks of enzyme activity, whereas gel chromatography of the A_{II} fraction gave only one peak. The kinetic behaviour of some of the fractions showed the existence of two enzymes (or two forms of the same enzyme) catalysing the same reaction with markedly different kinetic properties. In addition, humus–phosphomonoesterase complexes with higher molecular weight were more resistant to thermal and proteolytic denaturation than those with lower molecular weight (Nannipieri et al. 1996b). It was suggested that humus–enzyme complexes of higher molecular weight are likely to possess the molecular arrangement proposed by Burns et al. (1972), in which enzymes are surrounded by a network of humic molecules with pores large enough to permit the passage of substrates and products of the enzyme reaction, but not that of proteolytic enzymes.

The yield of phosphodiesterases extracted from a forest soil using a 0.1 M phosphate buffer was increased when KCl and EDTA were added to the buffer, probably because the extracting solution desorbed extracellular enzymes adsorbed on the surface of soil colloids by ionic bonding (Hayano 1977). The treatment of the soil extract with protamine sulfate removed brown-coloured substances (probably humic molecules), and the partially purified enzyme showed a pH optimum in the range of 5.2–6.0 and hydrolysed either the 3'- or the 5'-phosphodiester bond of deoxythymidine *p*-nitrophenyl phosphate. Two phosphodiesterase fractions extracted from an A horizon of a larch forest Andosol showed high affinity to adenosine 3'- and 5'-mononucleotides (Hayano 1988), whereas a third fraction was 2',3'-cyclic nucleotide 2'-phosphodiesterase (EC 3.1.4.16) or 2',3'-cyclic nucleotide 3'-phosphodiesterase (EC 3.1.4.37), with a pH optimum of 5.0 (Hayano 1987). All three phosphodiesterase active fractions were inhibited by Hg²⁺.

Approaches based on the study of clay–enzyme or humic–enzyme complexes, either prepared with pure enzymes or based on the extraction of humus–enzyme complexes from soil, present several drawbacks. In soil, neither enzymes nor surface-reactive particles, such as clays, are pure; for example, enzymes are released after cell death and lysis together with other cellular remains, which can affect adsorption of proteins to soil particles and the resistance of the formed complexes to thermal denaturation and proteolysis (Nannipieri et al. 1996b; Nielsen et al. 2006). Cell lysis with release of intracellular enzymes and artifacts due to interaction between enzymes and co-extracted soil components can occur during soil extraction.

The visualisation of the stabilised extracellular enzymes in the soil matrix by electron microscopy can give insights into the formation of stabilised enzyme complexes. Ultracytochemical tests have detected acid phosphomonoesterase activity in soil microbial cells and in fragments of microbial membranes as small as 7 × 20 nm, using electron microscopy. However, these tests were not able to locate enzymes in electron-dense minerals such as clays, or in soil components such as humic materials, reacting with counterstains such as OsO₄ (Ladd et al. 1996).

Techniques such as those based on the use of enzyme-labelled fluorescence (ELF)-97 phosphate seem to be promising for localising active phosphatases in the soil matrix because fluorescence is emitted after the enzymatic hydrolysis of the substrate (Wasaki et al. 2008; Wasaki and Maruyama 2011). This technique has detected acid phosphomonoesterase activity in roots of plants grown under P-deficient conditions, but its use in soil needs to be tested.

9.9 Conclusions and Future Research Needs

Probably, phosphodiesterase and phosphomonoesterase activities act sequentially in the mineralisation of organic P to inorganic P, which can be taken up by plant roots or by soil microorganisms (Fig. 9.1). These enzyme activities are higher in rhizosphere than in bulk soil and this suggests an important role for these enzyme activities in plant P nutrition. Acid phosphomonoesterase activity is more important in acid soils than is alkaline phosphomonoesterase activity, and vice versa in

alkaline soils. Effects of soil properties, agrochemicals, soil tillage, forest practices and pollutants have been extensively studied. However, a better understanding of the underlying mechanisms as well as the role of phosphatases (as that of any other enzyme in soil) requires setting up assays that discriminate the activities of enzymes associated with soil microorganisms from those of extracellular stabilised enzymes. A better understanding of the role of phosphatases in soil can be gained from studies on oxygen isotopes (isotope fractionation) of the phosphate group because alkaline phosphomonoesterase, unlike pyrophosphatase, catalyses a unidirectional reaction, producing kinetic isotope effects (Blake et al. 2005; Frossard et al. 2011).

The problem of measuring real rather than soil potential phosphatase activities may be solved by comparing the activities obtained from currently used enzyme assays with the organic P mineralisation (using isotope dilution methods as described by Frossard et al. 2011) in soils with a broad spectrum of properties, and by using the quantitatively most important soil organic P esters as substrates.

Extracellular enzyme activity of soil has been considered in the model describing decomposition of organic matter (Schimel and Weintraub 2003). However, the validation of this model requires determination of soil extracellular enzyme activity.

Phosphatases originate mainly from soil microorganisms, but in the rhizosphere and detritosphere they can also originate from plant cells. Changes in phosphatase activities have been related to changes in the composition of microbial communities, as determined by molecular techniques, in order to better understand the origin of phosphatases in soil. However, further insights into the origin of phosphatases in soil require relating the phosphatase activities to the expression of genes codifying these enzymes. The research carried out by Sakurai et al. (2008) involving the detection of alkaline phosphomonoesterase genes should be extended to soils with a broad spectrum of properties. Detection of the other genes expressing the various phosphatases is needed together with a relative comparison of the activity of these enzymes with the target gene.

References

- Adams MA (1992) Phosphatase activity and phosphorus fractions in karri (*Eucalyptus diversicolor* F. Muell.) forest soils. *Biol Fertil Soils* 14:200–204
- Arnebrant K, Schnurer J (1990) Changes in ATP content during and after chloroform fumigation. *Soil Biol Biochem* 22:875–877
- Ascher J, Ceccherini MT, Landi L, Mench M, Pietramellara G, Nannipieri P, Renella G (2009) Composition, biomass and activity of microflora, and leaf yield and foliar elemental concentrations of lettuce, after in situ stabilization of an arsenic-contaminated soil. *Appl Soil Ecol* 41:351–359
- Avidov E, Dick WA, Racke KD (1993) Proposed substrate for evaluating alkyl phosphomonoesterase activity in soil. *Soil Biol Biochem* 25:763–768
- Babich H, Bewley RJF, Stotzky G (1983) Application of “ecological dose” concept to the impact of heavy metals in some microbe mediated ecological processes in soil. *Arch Environ Contam Toxicol* 12:421–426

- Baligar VC, Wright RJ, Smedley MD (1988) Acid phosphatase activity in soil of the Appalachian Region. *Soil Sci Soc Am J* 52:1612–1616
- Bergstrom DW, Monreal CM, King DJ (1998) Sensitivity of soil enzyme activities to conservation practices. *Soil Sci Soc Am J* 62:1286–1295
- Berry DF, Shang C, Waltham Sajdak CA, Zelazny LW (2007) Measurement of phytase activity using tethered phytic acid as an artificial substrate: methods development. *Soil Biol Biochem* 39:361–367
- Bertrand D, de Wolf A (1968) Effect of microelement applied as complementary fertilizers on the soil microflora. *C R Acad Agric Fr* 54:1130–1133
- Blake RE, O’Neil JRO, Surkov AV (2005) Biogeochemical cycling of phosphorus: insights from oxygen isotope effects of phosphoenzymes. *Am J Sci* 305:596–620
- Brams WH, McLaren AD (1974) Phosphatase reactions in columns of soil. *Soil Biol Biochem* 6:183–189
- Browman MG, Tabatabai MA (1978) Phosphodiesterase activity of soils. *Soil Sci Soc Am J* 42:284–290
- Burns RG (1978) Enzyme activity in soil: some theoretical and practical considerations. In: Burns RG (ed) *Soil enzymes*. Academic, New York, pp 295–340
- Burns RG (1982) Enzyme activity in soil: location and possible role in microbial ecology. *Soil Biol Biochem* 14:423–427
- Burns RG, Pukite AH, McLaren AD (1972) Concerning the location and persistence of soil urease. *Soil Sci Soc Am Proc* 36:308–311
- Busman LM, Tabatabai MA (1985) Hydrolysis of trimetaphosphate in soils. *Soil Sci Soc Am J* 49:630–636
- Cervelli S, Nannipieri P, Sequi P (1973) Michaelis constant of soil acid phosphatase. *Soil Biol Biochem* 5:841–845
- Chhonkar PK, Tarafdar JC (1985) Degradation of clay-enzyme complexes by soil microorganisms. *Zbl Mikrobiol* 140:471–474
- Colvan SR, Syers JK, O’Donnell AG (2001) Effect of long-term fertiliser use on acid and alkaline phosphomonoesterase and phosphodiesterase activities in managed grassland. *Biol Fertil Soils* 34:258–263
- Cooperman BS, Baykov AA, Lahti R (1992) Evolutionary conservation of the active site of soluble inorganic pyrophosphatase. *Trends Biochem Sci* 7:262–266
- Cosgrove DJ (1967) Metabolism of organic phosphates in soil. In: McLaren AD, Peterson GH (eds) *Soil biochemistry*, vol 1. Marcel Dekker, New York, pp 216–228
- Cosgrove DJ (1980) Inositol phosphates. Their chemistry, biochemistry and physiology. Elsevier, Amsterdam
- Criquet S, Ferre E, Farnet AM, Le Petit J (2004) Annual dynamics of phosphatase activities in an evergreen oak litter: influence of biotic and abiotic factors. *Soil Biol Biochem* 36:1111–1118
- Criquet S, Braud A, Neble S (2007) Short-term effects of sewage sludge application on phosphatase activities and available P fractions in mediterranean soil. *Soil Biol Biochem* 39:921–929
- De Cesare F, Garzillo AM, Buonocore V, Badalucco L (2000) Use of sonication for measuring acid phosphatase activity in soil. *Soil Biol Biochem* 32:825–832
- Dick WA, Tabatabai MA (1977) Determination of orthophosphate in aqueous solutions containing labile organic and inorganic phosphorus compounds. *J Environ Qual* 6:82–85
- Dick WA, Tabatabai MA (1978) Inorganic pyrophosphatase activity of soil. *Soil Biol Biochem* 10:59–65
- Dick RP, Tabatabai MA (1986) Hydrolysis of polyphosphate in soils. *Soil Sci* 142:132–140
- Dick WA, Tabatabai MA (1987) Kinetic and activities of phosphatase-clay complexes. *Soil Sci* 143:5–15
- Dick WA, Juma NG, Tabatabai MA (1983) Effects of soils on acid phosphatase and inorganic pyrophosphatase of corn roots. *Soil Sci* 136:19–25

- Dick RP, Rasmussen PE, Kerle EA (1988) Influence of long-term residue management on soil enzyme activities in relation to soil chemical properties of a wheat-fallow system. *Biol Fertil Soils* 6:159–164
- Dinkelaker B, Marschner H (1992) In vivo demonstration of acid phosphatase activity in the rhizosphere of soil-grown plants. *Plant Soil* 144:199–205
- Doelman P, Haastra L (1989) Short- and long-term effects of heavy metals on phosphatase activity in soil: an ecological dose-response model approach. *Biol Fertil Soils* 8:235–241
- Drouillon M, Merckx R (2005) Performance of *para*-nitrophenyl phosphate and 4-methylumbelliferyl phosphate as substrate analogues for phosphomonoesterase in soil with different organic matter content. *Soil Biol Biochem* 37:1527–1534
- Dubovenko EK (1964) Phosphatase activity of different soils. *Chem Abstr* 62:13794f
- Eivazi F, Tabatabai MA (1977) Phosphatases in soils. *Soil Biol Biochem* 9:167–177
- Freeman C, Liska G, Ostle NJ, Jones SE, Lock MA (1995) The use of fluorogenic substrates for measuring enzyme activity in peatlands. *Plant Soil* 175:147–152
- Frossard E, Achat DL, Bernasconi SM, Bünemann EK, Fardeau J-C, Jansa J, Morel C, Rabeharisoa L, Randriamanantsoa L, Sinaj S, Tamburini F, Oberson A (2011) The use of tracers to investigate phosphate cycling in soil–plant systems. In: Bünemann EK, Oberson A, Frossard E (eds) *Phosphorus in action: biological processes in soil phosphorus cycling*. *Soil biology*, vol 26. Springer, Heidelberg. doi:10.1007/978-3-642-15271-9_3
- Garcia C, Hernandez T, Costa C, Ceccanti B, Masciandaro G, Ciardi C (1993) A study of biochemical parameters of composted and fresh municipal wastes. *Bioresour Technol* 44:17–23
- Garzillo AMV, Badalucco L, De Cesare F, Grego S, Buonocore V (1996) Synthesis and characterization of an acid phosphatase-polyresorcinol complex. *Soil Biol Biochem* 9:1155–1161
- Geller IT, Ginzburg KY (1979) Determination of the phosphatase activity of various soil groups. *Pochvovedeniye* 1:97–105
- George TS, Gregory PJ, Wood M, Read D, Buresh RJ (2002) Phosphatase activity and organic acids in the rhizosphere of potential agroforestry species and maize. *Soil Biol Biochem* 34:1487–1494
- George TS, Richardson AE, Simpson RJ (2005a) Behaviour of plant-derived extracellular phytase upon addition to soil. *Soil Biol Biochem* 37:977–978
- George TS, Richardson AE, Hadobas PA, Simpson RJ (2005b) Expression of fungal phytase gene in *Nicotiana tabacum* improves phosphorus nutrition of plants grown in amended soils. *Plant Biotechnol J* 3:129–140
- George TS, Richardson AE, Li SS, Gregory PJ, Daniell TJ (2009) Extracellular release of heterologous phytase from root of transgenic plants: does manipulation of rhizosphere biochemistry impact microbial community structure. *FEMS Microbiol Ecol* 70:433–445
- Gerritse RG, van Dijk H (1978) Determination of phosphatase activities of soils and animal wastes. *Soil Biol Biochem* 10:545–551
- Gianfreda L, Ruggiero P (2006) Enzyme activities in soil. In: Nannipieri P, Smalla K (eds) *Nucleic acids and proteins in soil*. Springer, Heidelberg, pp 257–311
- Grierson PF, Adams MA (2000) Plant species affect acid phosphatase, ergosterol and microbial P in a Jarrah (*Eucalyptus marginata* n ex SM) forest in south-western Australia. *Soil Biol Biochem* 32:1817–1827
- Halstead RL (1964) Phosphatase activity of soils as influenced by lime and other treatments. *Can J Soil Sci* 44:137–144
- Halstead RL, McKercher RB (1975) Biochemistry and cycling of phosphorus. In: Paul EA, McLaren AD (eds) *Soil biochemistry*, vol 4. Marcel Dekker, New York, pp 31–63
- Harrison AF (1983) Relationship between intensity of phosphatase activity and physico-chemical properties in woodland soils. *Soil Biol Biochem* 15:93–99
- Hayano K (1977) Extraction and properties of phosphodiesterase from a forest soil. *Soil Biol Biochem* 9:221–223

- Hayano K (1987) Characterization of two phosphodiesterase components in an extract of a larch forest soil. *Soil Sci Plant Nutr* 34:393–403
- Hayano K (1988) Characterization of a phosphodiesterase component in a forest soil extract. *Biol Fertil Soils* 3:159–164
- Ho I (1979) Acid phosphatase activity in forest soil. *Forest Sci* 25:567–568
- Hochstein L (1962) The fluorometric assay of soil enzymes. *Chem Abstr* 63:15164g
- Hofmann E (1963) The analysis of enzymes in soil. In: Linskens HF, Tracey MV (eds) *Moderne methoden der pflanzenanalyse*, vol VI. Springer, Berlin, pp 416–423
- Hysek J, Sarapatka B (1998) Relationship between phosphatase active bacteria and phosphatase activities in forest soils. *Biol Fertil Soils* 26:112–115
- Irving GJ, Cosgrove DJ (1976) The kinetics of soil acid phosphatase. *Soil Biol Biochem* 8:335–349
- Ishii T, Hayano K (1974) Method for the estimation of phosphodiesterase activity of soil. *Chem Abstr* 83:8023u
- Izaguirre-Mayoral ML, Flores S, Carballo O (2002) Determination of acid phosphatase and dehydrogenase activities in the rhizosphere of nodulated legume species native to two contrasting savanna sites in Venezuela. *Biol Fertil Soils* 35:470–472
- Jansa J, Finlay R, Wallander H, Smith FA, Smith SE (2011) Role of mycorrhizal symbioses in phosphorus cycling. In: Bünemann EK, Oberson A, Frossard E (eds) *Phosphorus in action: biological processes in soil phosphorus cycling*. Soil biology, vol 26. Springer, Heidelberg. doi:10.1007/978-3-642-15271-9_6
- Joergensen RG, Scheu S (1999) Response of soil microorganisms to the addition of carbon, nitrogen and phosphorus in a forest Rendzina. *Soil Biol Biochem* 31:859–866
- Joner EJ, Jakobsen I (1995) Growth and extracellular phosphatase activity of arbuscular mycorrhizal hyphae as influenced by soil organic matter. *Soil Biol Biochem* 9:1153–1159
- Joner EJ, Magid J, Gahoonia TS, Jakobsen I (1995) P depletion and activity of phosphatases in the rhizosphere of mycorrhizal and non-mycorrhizal cucumber (*Cucumis sativus* L.). *Soil Biol Biochem* 27:1145–1151
- Juma NG, Tabatabai MA (1977) Effects of trace elements on phosphatase activity in soils. *Soil Sci Soc Am J* 41:343–346
- Juma NG, Tabatabai MA (1978) Distribution of phosphomonoesterases in soils. *Soil Sci* 126:101–108
- Juma NG, Tabatabai MA (1988a) Hydrolysis of organic phosphates by corn and soybean roots. *Plant Soil* 107:31–38
- Juma NG, Tabatabai MA (1988b) Phosphatase activity in corn and soybean roots: conditions for assay and effects of metals. *Plant Soil* 107:39–47
- Juma NG, Tabatabai MA (1988c) Comparison of kinetic and thermodynamic parameters of phosphomonoesterases of soils and of corn and soybean roots. *Soil Biol Biochem* 20:533–539
- Kandeler E (1990) Characterization of free and adsorbed phosphatases in soils. *Biol Fertil Soils* 9:199–202
- Kandeler E (2007) Physiological and biochemical methods for studying soil biota and their function. In: Paul EA (ed) *Soil microbiology, ecology and biochemistry*. Academic, Amsterdam, pp 53–83
- Kandeler E, Palli S, Stemmer M, Gerzabeck MH (1999) Tillage changes in microbial biomass and enzyme activities in particle-size fractions of a Haplic Chernozem. *Soil Biol Biochem* 31:1253–1264
- Kandeler E, Marschner P, Tschirko D, Gahoonia TS, Nielsen NE (2002) Microbial community composition and functional diversity in the rhizosphere of maize. *Plant Soil* 238:310–312
- Kaplan DL, Hartenstein R (1979) Problems with toluene and the determination of extracellular enzyme activity in soils. *Soil Biol Biochem* 11:335–338
- Klose S, Tabatabai MA (1999) Urease activity of microbial biomass in soils. *Soil Biol Biochem* 31:205–211

- Kramer S, Green DM (2000) Acid and alkaline phosphatase dynamics and their relationship to soil microclimate in a semiarid woodland. *Soil Biol Biochem* 32:179–188
- Kumpiene J, Guerri G, Landi L, Pietramellara G, Nannipieri P, Renella G (2009) Microbial biomass, respiration and enzyme activities after in situ aided phytostabilization of a Pb- and Cu-contaminated soil. *Ecotoxicol Environ Saf* 72:115–119
- Ladd JN, Foster R, Nannipieri P, Oades JM (1996) Soil structure and biological activity. In: Stotzky G, Bollag J-M (eds) *Soil biochemistry*, vol 9. Marcel Dekker, New York, pp 23–78
- Lefebvre DD, Duff MG, Fife CA, Julien-Inalsigh C, Plaxton WC (1990) Response of phosphate deprivation in *Brassica nigra* suspension cells. *Plant Physiol* 93:504–511
- Lima JA, Nahas E, Gomes AC (1996) Microbial populations and activities in sewage sludge and phosphate fertilizer-amended soil. *Appl Soil Ecol* 4:75–82
- Makboul HE, Ottow JCG (1979) Alkaline phosphatase and Michaelis constant in the presence of different clay minerals. *Soil Sci* 128:129–135
- Malcom RE (1983) Assessment of phosphatase activity in soils. *Soil Biol Biochem* 15:403–408
- Marx MC, Wood M, Jarvis SC (2001) A microplate fluorimetric assay for the study of enzyme diversity in soils. *Soil Biol Biochem* 33:1633–1640
- Marzadori C, Gessa C, Ciurli S (1998) Kinetic properties and stability of potato phosphatase immobilized on Ca-polygalacturonate. *Soil Biol Biochem* 27:97–103
- Mathur SP, Rayment AF (1977) Influence of trace element fertilization on the decomposition rate and phosphatase activity of a mesic fibrisol. *Can J Soil Sci* 57:397–408
- Mathur SP, Sanderson RB (1978) Relationship between copper contents, rates of soil respiration and phosphatase activities of some histosols in an area of southwestern Quebec in the summer and the fall. *Can J Soil Sci* 58:125–134
- Mayaudon J (1986) The role of carbohydrates in the free enzymes in soil. In: Fuchsman CH (ed) *Peat and water*. Elsevier, New York, pp 263–308
- Mench M, Renella G, Gelsomino A, Landi L, Nannipieri P (2006) Biochemical parameters and bacterial species richness in soils contaminated by sludge-borne metals and remediated with inorganic soil amendments. *Environ Pollut* 14:24–31
- Moscatelli MC, Lagomarsino A, De Angelis O, Grego S (2005) Seasonality of soil biological properties in a poplar plantation growing under elevated atmospheric CO₂. *Appl Soil Ecol* 30:162–173
- Nannipieri P (1994) The potential use of soil enzymes as indicators of productivity, sustainability and pollution. In: Pankhurst CE, Doube BM, Gupta VVSR, Grace PR (eds) *soil biota: management in sustainable farming systems*. CSIRO, East Melbourne, pp 238–244
- Nannipieri P, Gianfreda L (1998) Kinetics of enzyme reactions in soil environments. In: Huang PM, Buffle SN, Senesi N, Buffle J (eds) *Structure and surface reactions of soil particles*. IUPAC series on analytical and physical chemistry of environmental systems, vol 4. Wiley, New York, pp 449–479
- Nannipieri P, Johanson RL, Paul EA (1978) Criteria for measurement of microbial growth and activity in soil. *Soil Biol Biochem* 10:223–229
- Nannipieri P, Pedrazzini F, Arcara PG, Piovanelli C (1979) Changes in amino acids, enzyme activities and biomasses during soil microbial growth. *Soil Sci* 127:26–34
- Nannipieri P, Ceccanti B, Cervelli S, Conti C (1982) Hydrolases extracted from soil: kinetic parameters of several enzymes catalysing the same reaction. *Soil Biol Biochem* 14:429–432
- Nannipieri P, Muccini L, Ciardi C (1983) Microbial biomass and enzyme activities: production and persistence. *Soil Biol Biochem* 15:679–685
- Nannipieri P, Sastre J, Landi L, Lobo MC, Pietramellara G (1996a) Determination of extracellular neutral phosphomonoesterase activity in soil. *Soil Biol Biochem* 28:107–112
- Nannipieri P, Sequi P, Fusi P (1996b) Humus and enzyme activity. In: Piccolo A (ed) *Humic substances in terrestrial ecosystems*. Elsevier, Amsterdam, pp 293–328
- Nannipieri P, Kandeler E, Ruggiero P (2002) Enzyme activities and microbiological and biochemical processes. In: Burns RG, Dick RP (eds) *Enzymes in the environment. Activity, ecology and applications*. Marcel Dekker, New York, pp 1–3

- Nannipieri P, Ascher J, Ceccherini MT, Landi L, Pietramellara G, Renella G (2003) Microbial diversity and soil functions. *Eur J Soil Sci* 54:655–670
- Neal JL (1973) Influence of selected grasses and forbs on soil phosphatase activity. *Can J Soil Sci* 53:119–121
- Nielsen KM, Calamai L, Pietramellara G (2006) Stabilization of extracellular DNA and proteins by transient binding to various soil components. In: Nannipieri P, Smalla K (eds) *Nucleic acids and proteins in soil*. Springer, Heidelberg, pp 141–188
- Ohmura H, Hayano K (1986) Effect of pH on phosphomonoesterase activity in soil. *Tochigi Agric Exp Stn Natl Inst Agro-Environ Sci*: 55–61 (Japanese article with a summary in English)
- Olander LP, Vitousek PM (2000) Regulation of soil phosphatase and chitinase activity by N and P availability. *Biogeochemistry* 49:175–190
- Oshima Y, Ogawa N, Harashima S (1996) Regulation of phosphatase synthesis in *Saccharomyces cerevisiae* – a review. *Gene* 179:171–177
- Pang PCK, Kolenko H (1986) Phosphomonoesterase activity in forest soils. *Soil Biol Biochem* 18:35–40
- Pant HK, Warman PR (2000) Enzymatic hydrolysis of soil organic phosphorus by immobilized phosphatases. *Biol Fertil Soils* 30:306–311
- Pereira M (1971) The effect of humic acids on the activity of phytase. *Humus Planta* 357–364
- Prado R, Tena M, Pinilla J-A (1982) Relationship between soil phosphatase activity and organic matter content. *Agronomie* 2:539–544
- Quiquampoix H, Mousain D (2005) Enzymatic hydrolysis of organic phosphorus. In: Turner BL, Frossard E, Baldwin DS (eds) *Organic phosphorus in the environment*. CABI, Wallingford, pp 89–112
- Ramirez-Martinez JR (1968) Organic phosphorus mineralization and phosphatase activity in soils. *Folia Microbiol* 13:161–174
- Ramirez-Martinez JR, McLaren AD (1966) Determination of soil phosphatase activity by a fluorimetric technique. *Enzymologia* 30:243–253
- Rao MA, Violante MA, Gianfreda L (1998) Interactions between tannic acid and acid phosphatase. *Soil Biol Biochem* 30:11–112
- Rejsek K (1991) Acid phosphomonoesterase activity of ectomycorrhizal roots in Norway spruce pure stands exposed to pollution. *Soil Biol Biochem* 23:667–671
- Renella G, Landi L, Nannipieri P (2002) Hydrolase activities during and after the chloroform fumigation of soil as affected by protease activity. *Soil Biol Biochem* 34:51–60
- Renella G, Ortigoza ALR, Landi L, Nannipieri P (2003) Additive effects of copper and zinc on cadmium toxicity on phosphatase activities and ATP content of soil as estimated by the ecological dose (ED₅₀). *Soil Biol Biochem* 35:1203–1210
- Renella G, Munch M, van del Lilie D, Pietramellara G, Ascher J, Ceccherini MT, Landi L, Nannipieri P (2004) Hydrolase activity, microbial biomass and community structure in long-term Cd-contaminated soils. *Soil Biol Biochem* 36:443–451
- Renella G, Mench M, Landi L, Nannipieri P (2005a) Microbial activity and hydrolase synthesis in long-term Cd-contaminated soils. *Soil Biol Biochem* 37:133–139
- Renella G, Mench M, Gelsomino A, Landi L, Nannipieri P (2005b) Functional activity and microbial community structure in soils amended with bimetallic sludges. *Soil Biol Biochem* 37:1498–1506
- Renella G, Landi L, Ascher MT, Ceccherini MT, Pietramellara G, Nannipieri P (2006a) Phosphomonoesterase production and persistence and composition of bacterial communities during plant material decomposition in soils in soil with different pH values. *Soil Biol Biochem* 38:795–802
- Renella G, Egamberdiyeva D, Landi L, Mench M, Nannipieri P (2006b) Microbial activity and hydrolase activities during decomposition of root exudates released by an artificial root surface in Cd-contaminated soils. *Soil Biol Biochem* 38:702–708
- Renella G, Landi L, Valori F, Nannipieri P (2007a) Microbial and hydrolase activity after release of low molecular weight organic compounds by a model root surface in a clayey and a sandy soil. *Appl Soil Ecol* 36:124–129

- Renella G, Szukcis U, Landi L, Nannipieri P (2007b) Quantitative assessment of hydrolase production and persistence in soil. *Biol Fertil Soils* 44:321–329
- Rojo MJ, Carcedo SG, Mateos MP (1990) Distribution and characterization of phosphatase and organic phosphorus in soil fractions. *Soil Biol Biochem* 22:169–174
- Saa A, Trasar-Cepeda MC, Gil-Sotres F, Carballas T (1993) Change in soil phosphorus and acid phosphatase activity immediately following forest fires. *Soil Biol Biochem* 25:1223–1230
- Saa A, Trasar-Cepeda MC, Gil-Sotres F, Carballas T (1998) Soil P status and phosphomonoesterase activity of recently burnt and unburnt soil following laboratory incubation. *Soil Biol Biochem* 30:419–428
- Sakurai M, Wasaki J, Tomizawa Y, Shinano T, Osaki M (2008) Analysis of bacterial communities on alkaline phosphatase gene in organic matter applied soil. *Soil Biol Biochem* 54:62–71
- Santruckova H, Vrba J, Picek T, Kopacek J (2004) Soil biochemical activity and phosphorus transformations and losses from acidified forest soils. *Soil Biol Biochem* 36:1569–1576
- Sarkar JM, Leonowicz A, Bollag JM (1989) Immobilization of enzymes on clay and soils. *Soil Biol Biochem* 21:223–230
- Schimel JP, Weintraub MN (2003) The implications of exoenzyme activity on microbial carbon and nitrogen limitation in soil: a theoretical model. *Soil Biol Biochem* 35:549–563
- Schmidt G, Laskowski M Sr (1961) Phosphatase ester cleavage (survey). In: Boyer PD, Lardy H, Myrback K (eds) *The enzymes*, 2nd edn. Academic, New York, pp 3–35
- Schneider K, Turrión M-B, Grierson PF, Gallardo JF (2001) Phosphatase activity, microbial phosphorus, and fine root growth in forest soils in the Sierra de Gata, western Spain. *Biol Fertil Soils* 34:151–155
- Skujins J (1978) History of abiotic soil enzyme research. In: Burns RG (ed) *Soil enzymes*. Academic, London, pp 1–49
- Speir TW, Cowling JC (1991) Phosphatase activities of pasture plants and soils: relationship with plant productivity and P fertility indices. *Biol Fertil Soils* 12:189–194
- Speir TW, Ross DJ (1978) Soil phosphatase and sulphatase. In: Burns RG (ed) *Soil enzymes*. Academic, London, pp 197–250
- Spiers GA, McGill WB (1979) Effects of phosphorus addition and energy supply on acid phosphatase production and activity in soils. *Soil Biol Biochem* 11:3–8
- Stotzky G (1986) Influence of soil mineral colloids and metabolic processes, growth adhesion, and ecology of microbes and viruses. In: Huang M, Schnitzer M (eds) *Interactions of soil minerals with natural organics and microbes*, special publication 17. Soil Science Society of America, Madison, pp 305–428
- Tabatabai MA (1994) Soil enzymes. In: Weaver RW, Angle S, Bottomley P, Bezdicek D, Smith S, Tabatabai A, Wollum A (eds) *Methods of soil analysis. Part 2. Microbiological and biochemical properties*. Soil Science Society of America, Madison, pp 775–833
- Tabatabai MA, Bremner JM (1969) Use of *p*-nitrophenyl phosphate for assay of soil phosphatase activity. *Soil Biol Biochem* 1:301–307
- Tabatabai MA, Bremner JM (1971) Michaelis constant of soil enzymes. *Soil Biol Biochem* 3:317–323
- Tabatabai MA, Dick WA (1979) Distribution and stability of phosphatase in soil. *Soil Biol Biochem* 11:655–659
- Tabatabai A, Fu M (1992) Extraction of enzymes from soil. In: Stotzky G, Bollag J-M (eds) *Soil biochemistry*, vol 7. Marcel Dekker, New York, pp 197–227
- Tarafdar JC, Chhonkar PK (1978) Status of phosphatases in the root-soil interface of leguminous and non-leguminous crops. *Z Pflanz Bodenkunde* 141:347–351
- Tarafdar JC, Jungk A (1987) Phosphatase activity in the rhizosphere and its relation to the depletion of soil organic phosphorus. *Biol Fertil Soils* 3:199–204
- Torsvik V, Sørheim R, Gorksoyr J (1996) Total bacterial diversity in soil and sediment communities – a review. *J Ind Microbiol* 17:170–178
- Trasar-Cepeda MC, Gil-Sotres F (1988) Kinetics of acid phosphatase activity in various soils of Galicia (NW Spain). *Soil Biol Biochem* 20:275–280

- Turner BL, Haygarth PM (2005) Phosphatase activity in temperate pasture soils: potential regulation of labile organic phosphorus turnover by phosphodiesterase activity. *Sci Total Environ* 344:27–36
- Turner BL, Paphazy MJ, Haygarth PM, McKelvie ID (2002a) Inositol phosphates in the environment. *Philos Trans R Soc Lond B Biol Sci* 357:449–469
- Turner BL, Baxter R, Whitton BA (2002b) Seasonal phosphatase activity in three characteristic soils of the English uplands polluted by long-term atmospheric nitrogen deposition. *Environ Pollut* 120:313–317
- Tyler G (1976) Heavy metal pollution, phosphatase activity and mineralization of organic phosphorus in forest soils. *Soil Biol Biochem* 8:327–332
- Vuorinen AH, Saharinen MH (1996) Effects of soil organic matter extracted from soil on acid phosphomonoesterase. *Soil Biol Biochem* 28:1477–1481
- Wasaki J, Maruyama H (2011) Molecular approaches to the study of biological phosphorus cycling. In: Bünemann EK, Oberson A, Frossard E (eds) *Phosphorus in action: biological processes in soil phosphorus cycling*. Soil biology, vol 26. Springer, Heidelberg. doi: 10.1007/978-3-642-15271-9_4
- Wasaki J, Kojima S, Maruyama H, Haase S, Osaki M, Kandeler E (2008) Localization of acid phosphatase activities in the roots of white lupin plants grown under phosphorus-deficient conditions. *Soil Sci Plant Nutr* 54:95–102
- Yadav RS, Tarafdar JC (2001) Influence of organic and inorganic phosphorus supply on the maximum secretion of acid phosphatase by plants. *Biol Fertil Soils* 34:140–143
- Yadav RS, Tarafdar JC (2003) Phytase of fungi in arid and semi-arid soils and their efficiency in hydrolysing different organic P compounds. *Soil Biol Biochem* 35:1–7
- Zornoza R, Mataix-Solera J, Guerrero C, Arcenegui V, Mataix-Beneyto J (2009) Comparison of soil physical, chemical and biochemical properties among native forest, maintained and abandoned almond orchards in mountainous areas of eastern Spain. *Arid Land Res Manag* 23:267–282