

Occurrence of nitrogen-fixing *Azospirillum* in vesicular-arbuscular mycorrhizal fungi

K.V.B.R. TILAK¹, C.Y. LI² and IWAN HO²

¹Division of Microbiology, Indian Agricultural Research Institute, New Delhi-110012, India and ²US Department of Agriculture, Forest Service, Pacific Northwest Research Station, Forestry Sciences Laboratory, 3200 SW Jefferson Way, Corvallis, OR 97331, USA

Received 31 May 1988. Revised February 1989

Key words: *Azospirillum*, *Endogone*, *Gigaspora*, *Glomus*, spores

Abstract

Nitrogenase activity measured by acetylene reduction, was detected when surface-sterilized spores of vesicular-arbuscular (VA) mycorrhizal fungi (*Glomus fasciculatum*, *G. intraradices*, *G. scientillans*, *G. mosseae*, *Gigaspora gilmorei* and *Endogone dusii*) were inoculated into nitrogen-free liquid medium containing malic acid and incubated under microaerophilic conditions (99% N₂ + 1% O₂) at 30°C. *Azospirillum* species were isolated from the nitrogenase-active cultures.

Introduction

Through the years there have been reports of bacteria isolated from fungal spores. Several heterotrophic, fermentative and anaerobic bacteria from the lumen of *Glomus macrocarpus* growing in the rhizospheres of *Ficus bengelensis* and *Acacia* sp. have been reported (Varma *et al.*, 1981). Occurrence of *Klebsiella* sp., *Clostridium* sp. and *Streptococcus lactis* from the lumen of *Glomus microcarpus* extracted from the rhizospheres of certain xerophytes has been documented (Singh and Varma, 1985). Nitrogen-fixing (acetylene-reducing) bacteria associated with the ectomycorrhizae of Douglas-fir (*Pseudotsuga menziesii*) have recently been demonstrated (Li and Hung, 1987). Nitrogen-fixing azospirilla were isolated from within the sporocarps of the ectomycorrhizal fungi like *Hebeloma crustuliniforme*, *Laccaria laccata* and *Rhizopogon vinicolor* (Li and Castellano, 1987).

This study explores the possible association of nitrogen-fixing *Azospirillum* sp. with the spores of VA-mycorrhizal fungi.

Materials and methods

Spores of *Glomus fasciculatum*, *G. intraradices*, *G. scientillans*, *G. mosseae*, *Gigaspora gilmorei* and *Endogone dusii* were isolated from the rhizosphere soil of maize infected with the respective VAM fungi by the standard wet sieving and decanting technique (Gerdeman and Nicholson, 1963). They were rinsed with tap water in 70% alcohol followed by immersion in 1% chloramine -T (J T Baker Chemical Co., Phillipsburg, NJ)* for 1 h shaken frequently, and rinsed in two changes of sterile distilled water (Döbereiner, 1980). About 50–75 surface-sterilized spores of VAM fungi were transferred into 8 ml of nitrogen-free semi-solid sodium malate medium (Döbereiner and Day, 1976) in 20 ml capacity screw-capped tubes and incubated at 30°C for a week. Six replicates were maintained in each case. The formation of white pellicel 2–4 mm below the surface of the medium was noticed with all the cultures.

* Use of trade names does not imply endorsement or approval by the USDA Forest Service.

Acetylene was injected into each tube to 10% of the total gas volume. Tubes without added acetylene served as controls. After 24 h, 0.1 ml gaseous sample from each tube was removed and analyzed for ethylene and acetylene with a Hewlett-Packard 5830A gas chromatograph fitted with 2 m × 2.1 mm, 80–100 mesh Porapak R column with oven temperature at 70°C, and flow rate of the nitrogen carrier gas was adjusted to 40 ml per min.

Acetylene-reducing bacteria from the mixed cultures were purified by repeated streaking on N-free sodium malate medium supplemented with 0.002% yeast extract (Barber and Evans, 1975). To test the nitrogenase activity of isolated bacteria, as measured by acetylene reduction an aqueous suspension of each bacterial isolate was inoculated into screw-capped tubes that contained N-free sodium malate medium supplemented with yeast extract (0.002%). The tubes were incubated under micro-aerophilic conditions (99% N₂ + 1% O₂) at 30°C for 3 days. Acetylene was injected into each tube. The formation of ethylene was determined as previously described.

Bacterial cells in the tubes were harvested and washed with cold 5% trichloro-acetic acid. Cell protein was solubilized with 0.5 N NaOH in a boiling water bath for 10 min (Agarwal and Keister, 1983) and measured by the modified Lowry method (Markwell *et al.*, 1978).

Results and discussion

Bacteria isolated from the surface-sterilized spores of *Glomus fasciculatum*, *G. intraradices*, *G. scientillans*, *G. mosseae*, *Gigaspora gilmorei* and *Endogone dusii* showed characteristic spirillar movement and grew well in nutrient and trypticase soy agar media. The cells are long and slightly curved rods with poly-β-hydroxy-butyrate (PHB) granules inside and showed a distinctive pellicle formation with acetylene reduction activity in N-free sodium malate semi-solid medium under microaerophilic conditions (Table 1). These characteristics are typical of the genus *Azospirillum* (Krieg and Döbereiner, 1984). The *Azospirillum* isolated from *Glomus* species, in general had significantly more nitrogenase activity than that isolated from either *Gigaspora gilmorei* or *Endogone dusii*.

Table 1. Nitrogenase activities of *Azospirillum* sp. isolated from surface-sterilized spores of different vesicular-arbuscular (VA) mycorrhizal fungi¹

VAM spores	Nitrogenase activity (nmoles ethylene produced/ mg protein/h)
<i>Glomus fasciculatum</i>	125b
<i>G. intraradices</i>	119b
<i>G. scientillans</i>	105b
<i>G. mosseae</i>	110b
<i>Gigaspora gilmorei</i>	55a
<i>Endogone dusii</i>	34a

¹ Data are means of six replicates. Treatment means not sharing a common letter differ significantly at 5% level of significance (analysis of variance). The critical difference at 5% level is 22.52.

The implications of nitrogen-fixing azospirilla from the surface-sterilized spores of VAM fungi is not known, although many fungi do utilize large amounts of nitrogen in producing sporocarps and spores. Fixation of atmospheric nitrogen by associated bacteria could perhaps satisfy such demands to certain extent (Cowling and Merrill, 1966). Although the methods adopted to isolate nitrogen-fixing azospirilla from thoroughly surface-sterilized spores of VAM-fungi, there is a possibility of association of few other bacteria which may not be of much ecological significance. However, further studies on the association of bacteria other than nitrogen-fixing azospirilla and their contribution to nitrogen fixation either directly or indirectly needs further investigation.

Acknowledgements

These studies were supported through the Indo-American Science and Technology Initiative, a program of cooperative research between India and the United States. Our sincere thanks are due to Dr. James M. Trappe, Professor of Forest Sciences for encouragement.

References

- Agarwal A K and Keister D L 1983 Appl. Environ. Microbiol. 45, 1592–1601.
- Barber L E and Evans H J 1975 Can. J. Microbiol. 22, 254–260.
- Cowling E B and Merrill W 1966 Can. J. Bot. 44, 1539–1554.
- Döbereiner J 1980 In Methods for Evaluating Biological

- Nitrogen Fixation. Ed. F J Bergersen. pp 535–555. John Wiley and sons, New York.
- Döbereiner J and Day J M 1976 *In* Proc. 1st International Symp. Nitrogen Fixation, Vol. 2. Ed. W E Newton and C J Nyman. pp 518–538. Washington State Univ. Press, Pullman, WA, USA.
- Gerdemann J W and Nicholson T A 1963 *Trans. Br. Mycol. Soc.* 46, 235–244.
- Krieg N R and Döbereiner J 1984 *In* Bergey's Manual of Systematic Bacteriology. Ed. N R Krieg. pp 94–104. Williams and Wilkins, Baltimore, MA, USA.
- Li C Y and Castellano M A 1987 *Trans. Br. Mycol. Soc.* 88(4), 563–565.
- Li C Y and Hung L L 1987 *Plant and Soil* 98, 425–428.
- Markwell M A K *et al.* 1978 *Anal. Biochem.* 87, 206–210.
- Singh K and Varma A K 1985 *Trans. Mycol. Soc. Japan* 26, 511–516.
- Varma A K *et al.* 1981 *Curr. Microbiol.* 6, 207–215.