

## ORIGINAL PAPER

Y. Bashan · L. E. Gonzalez

# Long-term survival of the plant-growth-promoting bacteria *Azospirillum brasilense* and *Pseudomonas fluorescens* in dry alginate inoculant

Received: 1 May 1998 / Received revision: 24 August 1998 / Accepted: 3 September 1998

**Abstract** Two plant-growth-promoting bacteria, *Azospirillum brasilense* Cd and *Pseudomonas fluorescens* 313, immobilized in 1983 in two types of alginate-bead inoculant (with and without skim-milk supplement) and later dried and stored at ambient temperature for 14 years, were recovered in 1996. The population in each type of bead had decreased, yet significant numbers survived ( $10^5$ - $10^6$  cfu/g beads). Population numbers depended on the bead type and the three independent bacterial counting methods: the conventional plate count method, indirect enzyme-linked immunosorbent assay and the limited-enrichment technique. Both bacterial species retained several of their original physiological features. When inoculated onto wheat plants, both species colonized and produced plant-growth effects equal to those of the contemporary strain from a culture collection or to their own 1983 records. This study showed that bacteria can survive in alginate inoculant over long periods.

## Introduction

Today, most marketed bacterial inoculants are peat-based formulations for seed coating or pellets for sowing (Smith 1992), performing satisfactorily primarily with rhizobia (Thompson 1980). During the last two decades several experimental formulations based on polymers have been evaluated (Bashan 1998), and demonstrated to be potential bacterial carriers (Jung et al. 1982). They offer advantages

over peat, encapsulating and protecting the microorganisms against environmental stresses, and releasing them to the soil gradually when the polymers are degraded by soil microbes (Bashan 1998). They can also be stored dry at ambient temperatures, offer consistent batch quality and a better-defined environment for the bacteria, and can be manipulated according to species requirements. These inoculants may be amended with nutrients to improve short-term survival upon inoculation, especially with associated plant growth-promoting bacteria (PGPB) (Bashan and Holguin 1998). However, they are more expensive and are more labor-intensive to produce than peat-based inoculants (Fages 1992). So far, there are no polymer-based inoculants on the market (Ogoshi et al. 1997).

Alginate is the most common polymeric material for encapsulation of microorganisms for commercial use (Cassidy et al. 1996). Macro-alginate beads (1-3 mm) containing bacteria are made by an easy and straightforward multi-step procedure (Bashan 1986a; Digat 1991) that may involve crushing or milling solid sheets of alginate into small particles and sieving them to the required size (Paul et al. 1993).

Several experimental alginate-based preparations were evaluated for agriculture, including the encapsulation of the VA-mycorrhizal fungi (Ganry et al. 1982), ectomycorrhizal fungi (Le Tacon et al. 1985), PGPB (Trevors et al. 1992), and fungi (DeLuca et al. 1990) and bacteria (Aino et al. 1997) used as biocontrol agents against soil-borne pathogens. This technology was also employed to encapsulate the PGPB *A. brasilense* and *P. fluorescens* (Bashan 1986a). *A. brasilense* survived for at least 4 months (Bashan 1986a; Paul et al. 1993) in dry alginate preparations. These preparations were later used successfully to inoculate wheat under field conditions (Bashan et al. 1987). Samples from these earlier studies were the source of the inoculant for the present study.

The aim of this study was to evaluate the capacity of *A. brasilense* and *P. fluorescens* for long-term survival, viability, and growth-promoting capacity after storage in dry alginate beads for 14 years.

Y. Bashan (✉)

Department of Microbiology, The Center for Biological Research of the Northwest (CIB), La Paz, P.O. Box 128, B.C.S. 23000, Mexico

e-mail: bashan@cibnor.mx

Fax: +52-112-54710 or -53625

L. E. Gonzalez

Department of Biology, Pontificia Universidad Javeriana, Apartado Aereo 56710, Santafe de Bogota, Colombia

## Materials and methods

### Bead production

The macrobeads used in this study were produced in 1983 (Bashan 1986a). The dry preparations were stored for 14 years at 10-28 °C in dark, screw-cap bottles containing about 4 g beads and about 5 g dry silica gel. Since the inoculant was conserved for archival purposes, no treatment was given and the bottles remained tightly closed during the entire period.

### Bacteria and plant material

The PGPB *A. brasilense* Cd (DSM 7030) and *P. fluorescens* 313 were the encapsulated bacteria. Wheat plants (*Triticum aestivum* cv. Miriam, Hazera Co., Haifa, Israel), tomato plants (*Lycopersicon esculentum* Mill. cv. M-82-1-8-VF), and cotton plants (*Gossypium barbadense* cv. Pima S-5, Hazera), all known to respond positively to *A. brasilense* inoculation (Bashan et al. 1989; Kapulnik et al. 1985), were used to evaluate the growth-promoting effect of the recovered bacteria.

### Growth-promotion effects of the recovered bacteria

Plant growth-promoting activity was measured under gnotobiotic conditions by the method of Glick et al. (1995) with minor modifications. Seeds were surface-sterilized and inoculated with the respective PGPB as described earlier (Puente and Bashan 1993). Bacteria were isolated after the beads had been dissolved, as described later, and grown using standard methods (Bashan et al. 1993). Five inoculated seeds each were germinated in polyester growth pouches (Northrup King Co. Minneapolis, Minn.), covered with transparent plastic sheets and incubated at 22 ± 2 °C (wheat) or 28 ± 2 °C (tomato, cotton) in a growth chamber (Conviro, model TC 16, Winnipeg, Canada) at a light intensity of 120 μmol m<sup>-2</sup> s<sup>-1</sup> and a photoperiod of 12 h. Root length, root and shoot dry mass, root surface area, formation of root hairs, and root-colonization capacity (obtained by inoculating the recovered strains) were compared with the same parameters of plants inoculated with the same *A. brasilense* strain from a culture collection (stored in liquid N) and with noninoculated seedlings growing under identical conditions. Since a collection strain of the original *P. fluorescens* 313 was no longer available (the original strain, stored by the senior author at -40 °C in 1987 at the Weizmann Institute of Science, Israel, belonged to an Israeli commercial company, no longer existing, and is presumed lost), similar plant promoting effects recorded in unpublished data from 1983 were included as reference.


### Bacterial counts from beads

Bacteria entrapped in the beads were counted after ten bead samples were dissolved in 10 ml potassium phosphate buffer (0.25 M, pH 6.8 ± 0.1) in a test-tube for 16-24 h at 30 ± 2 °C. No bacterial multiplication occurred under these incubations (Bashan 1986a; Levanony et al. 1987). To facilitate solubility, the beads were shaken for 5 min at top speed on a Vortex-Genie-2 mixer (Scientific industries, Bohemia, N.Y.). For verification of results, the *A. brasilense* released from the beads was counted simultaneously by three different methods: a conventional plate count method on nutrient agar (Difco, Detroit, Mich.), indirect enzyme-linked immunosorbent assay (ELISA) (Levanony et al. 1987), and by the limited-enrichment technique (Bashan et al. 1991). In the absence of immunomethods for *P. fluorescens*, its cell numbers were counted only by the plate-count method. Because the solubilization of the beads could not guarantee a single-cell suspension, recovery number should be considered a minimal population level.

### Root-colonization capacity of the recovered bacteria

Bacterial counts from roots were taken after root segments had been homogenized and suspended in 5 ml 0.85% NaCl at 4000 rpm for 30 s at 22 ± 2 °C (Polytron Brinkmann Instruments, NY). The homogenate was serially diluted in the same saline solution and plated on nutrient agar (Difco, Detroit, Ill.), using the plate-count method for *P. fluorescens* and indirect ELISA for *A. brasilense*. The samples were then filtered through Whatman no. 1 filter-paper for the dry-mass determination (60 °C for 16 h).

Bacterial characterization and the effects on plants after inoculation with recovered *A. brasilense* Cd from dry inoculants. These were done using standard methods and are described in footnotes for Table 1 and Fig. 2.



**Fig. 1A, B** Survival of *Azospirillum brasilense* Cd and *Pseudomonas fluorescens* 313 in dry alginate beads (A) and in dry alginate beads supplemented with skim milk (B) after 14 years, as measured by three independent bacterial counting methods: the plate-count method (PCM), indirect enzyme-linked immunosorbent assay (ELISA) and the limited-enrichment method (Enrichment). Bars Represent SE. The original bacterial concentration in the beads produced in 1983 were as follows: in alginate beads,  $7.4 \times 10^6$  *A. brasilense* cfu/g beads and  $6.3 \times 10^6$  *P. fluorescens* cfu/g beads; in alginate beads supplemented with skim-milk,  $9.7 \times 10^9$  *A. brasilense* cfu/g beads and  $8.3 \times 10^9$  *P. fluorescens* cfu/g beads

**Table 1** Typical characteristics of *Azospirillum brasilense* Cd recovered from dried beads as compared to characteristics of the same strain from a culture collection

Characteristic	<i>A. brasilense</i> Cd (recovered)	<i>A. brasilense</i> Cd (collection)
N <sub>2</sub> fixation in culture <sup>a</sup> (nmol C <sub>2</sub> H <sub>4</sub> culture <sup>-1</sup> h <sup>-1</sup> )	1200 ± 150	990 ± 180
Pink color of colony <sup>b</sup> (A <sub>500</sub> )	0.43	0.43
Motility in liquid <sup>c</sup>	+	+
Swarming on agar <sup>d</sup>	+	+
Capacity to form large aggregate <sup>e</sup>	+	+
Root colonization of noncereals (cotton) <sup>f</sup>	5.4 ± 0.8 × 10 <sup>5</sup>	3.8 ± 0.4 × 10 <sup>5</sup>
Root colonization of noncereals (tomato) <sup>f</sup>	6.6 ± 0.7 × 10 <sup>5</sup>	4.7 ± 0.9 × 10 <sup>5</sup>
Increased formation of root hairs and root-hair length <sup>g</sup>	+	+

<sup>a</sup> According to Holguin et al. (1992). <sup>b</sup> According to Eskew et al. (1977). <sup>c</sup> According to Reinhold et al. (1985). <sup>d</sup> According to Hall and Krieg (1983). <sup>e</sup> According to Del Gallo et al. (1989) in the presence of fructose and KNO<sub>3</sub>. <sup>f</sup> According to Bashan et al. (1989), 14 days after inoculation. <sup>g</sup> According to Dubrovsky et al. (1994)

#### Experimental design and statistical analysis


Samples of ten dry beads each, in six replicates, were used for recovering the stored bacteria. Ten replicates were used for the growth-promotion effects and for all bacterial counts. A replicate consisted of five seeds per pouch, and three independent determinations of the bacterial population. Because of the shortage and irreplaceability of dry bead samples, recovery of bacteria from the beads was done only once. Growth-promoting tests were repeated. Results were analyzed by one-way analysis of variance and Student's t-test at  $P < 0.05$  using Statgraphic plus (Statistical

Graphics, Rockville, Maryland) and SigmaPlot (Jandel, San Rafael, Calif.) software.

#### Results

Taking advantage of an experimental dry inoculant preserved in 1983 for archival purposes, we were able to study how two PGPB survived for 14 years and maintain

**Fig. 2A–D** The effect on variables of wheat plant growth after inoculating with recovered (skim-milk beads) and collection based *A. brasilense* Cd and recovered *P. fluorescens* 313. In the absence of *P. fluorescens* from a collection in 1996, the data on this strain's performance were taken from 1983 unpublished notes. Columns denoted by different letters in A, B, C or D differ significantly at  $P < 0.05$  by one-way analysis of variance. Root lengths and dry weights were determined according to Bashan (1986b) and root surface area was measured according to Carley and Watson (1966)



their beneficial plant-growth-promoting characteristics.

The numbers of *A. brasilense* decreased to 10% of the original population after storage in beads for 14 years, as measured by three independent methods, with results not significantly different among the methods (Fig. 1A). A greater decrease occurred in alginate beads supplemented with skim milk, where the population fell to 1/1000 of the original (Fig. 1B). However, since the original population in the skim-milk beads was higher than that in the milk-free beads (Bashan 1986a), its level was still over  $10^6$  cfu/g bead after 14 years, equivalent to the Canadian standard for freshly prepared *Rhizobium* inoculant (Bordeleau and Prevost 1981). The other PGPB, *P. fluorescens*, did not survive as well as *A. brasilense*, yet its population in both types of bead after 14 years was high (Fig. 1).

When eight typical physiological and environmental characteristics of the recovered *A. brasilense* Cd were compared to those of the same strain obtained from the German DSM culture collection, no differences between the strains were recorded (Table 1). The recovered *A. brasilense* Cd significantly affected plant development, similarly to the *A. brasilense* Cd from the collection. The 1996 results of *P. fluorescens* were insignificantly different from the results of the original strain tested in 1983 (Fig. 2). The root

colonization capacity of the recovered *A. brasilense* Cd was superior to that of the original bacteria, whereas the root colonization of *P. fluorescens* in 1996 was inferior to that described in the 1983 records extant on this bacterium (Fig. 3).

## Discussion


The main advantages of alginate inoculants are their nontoxic nature, their degradation in the soil and, above all, their slow release of the entrapped microorganisms into the soil (Bashan 1998). Genetically engineered *P. fluorescens*, encapsulated and later released into soil microcosms, showed significantly longer survival rates than those of unencapsulated cells after 3 months (van Elsas et al. 1992). PGPB are known to survive for months in air-dried and lyophilized inoculants (Bashan 1986a; Paul et al. 1993). For commercialization, a long shelf-life is an advantage for any inoculant (Fages 1990, 1992). Currently, a shelf-life of about a year is mandatory for peat inoculants containing rhizobia (Bashan 1998). A longer shelf-life, with full preservation of strain characteristics, is desirable (Fages 1992; Smith, 1992). However, a very long inoculant survival period is not an industrial requirement.

Long-term survival of inoculants is a matter of commercial secrecy (Bashan 1998). To our knowledge, no information is available in the literature on very long survival periods of plant-growth-promoting *Azospirillum* in synthetic inoculants (Bashan and Holguin 1997a, b). We know of no published results for *P. fluorescens* 313.

The ultimate test for any PGPB is its ability to colonize its target plant roots and to produce growth effects (Bashan and Holguin 1997a; Glick and Bashan 1997). By inoculating a common host plant, such as wheat, and measuring the effects of PGPB on root length, root dry mass, root surface area and shoot dry weight, it was possible to compare and verify that plant responses to the recovered bacteria were similar to those obtained by control, culture-collection strains.

In summary, two PGPB survived for 14 years in dry alginate beads at ambient temperature. Upon recovery, both species retained the capacity to enhance plant growth.

**Acknowledgements** This paper was written in memory of our late first English editor Dr. Roy Bowers and in memory of the late Mr. Avner Bashan from Israel. It was supported by Consejo Nacional de Ciencia y Tecnología (CONACYT), Mexico, contract 26262-B and by Instituto Colombiano para el Desarrollo de la Ciencia y la Tecnología Francisco José de Caldas (COLCIENCIAS), Colombia. We thank Mrs. Hanna Levanony, Department of Plant Sciences, The Weizmann Institute of Science, Israel for excellent technical assistance in preparation of the original beads, Dr. Ellis Glazier for editing the English text, Drs. Marina and Gabor Bethlenfalvai for the English style, and Mr. Edgar Yuan for computer searches of the literature.



**Fig. 3** Root colonization after inoculation of the wheat plants with recovered and collection-based *A. brasilense* Cd and recovered *P. fluorescens* 313. In the absence of *P. fluorescens* from a collection in 1996, the data on this strain's performance were taken from 1983 unpublished notes. Pairs of columns, for each bacterial species separately, denoted by different letters, differ significantly at  $P < 0.05$  by Student's *t*-test

## References

- Aino M, Maekawa Y, Mayama S, Kato H (1997) Biocontrol of bacterial wilt of tomato by producing seedlings colonized with endophytic antagonistic pseudomonads. In: Ogoshi A, Kobayashi K, Homma Y, Kodama F, Kondo N, Akino S (eds) Plant growth-promoting Rhizobacteria - present status and future prospects. Faculty of Agriculture, Hokkaido University, Sapporo, pp 120-123
- Bashan Y (1986a) Alginate beads as synthetic inoculant carriers for the slow release of bacteria that affect plant growth. *Appl Environ Microbiol* 51: 1089-1098
- Bashan Y (1986b) Significance of timing and level of inoculation with rhizosphere bacteria on wheat plants. *Soil Biol Biochem* 18: 297-301
- Bashan Y (1998) Inoculants for plant growth-promoting bacteria in agriculture. *Biotechnol Adv* 16: 729-770
- Bashan Y, Holguin G (1997a) *Azospirillum*-plant relationships: environmental and physiological advances (1990-1996). *Can J Microbiol* 43: 103-121
- Bashan Y, Holguin G (1997b) Short- and medium term avenues for *Azospirillum* inoculation. In: Ogoshi A, Kobayashi K, Homma Y, Kodama F, Kondo N, Akino S (eds) Plant growth-promoting Rhizobacteria - present status and future prospects. Faculty of Agriculture, Hokkaido University, Sapporo, pp 130-149
- Bashan Y, Holguin G (1998) Proposal for the division of plant growth-promoting Rhizobacteria into two classifications: biocontrol-PGPB (plant growth-promoting bacteria) and PGPB. *Soil Biol Biochem* 30: 1225-1228
- Bashan Y, Levanony H, Ziv-Vecht O (1987) The fate of field-inoculated *Azospirillum brasilense* Cd in wheat rhizosphere during the growing season. *Can J Microbiol* 33: 1074-1079
- Bashan Y, Ream Y, Levanony H, Sade A (1989) Nonspecific responses in plant growth, yield, and root colonization of noncereal crop plants to inoculation with *Azospirillum brasilense* Cd. *Can J Bot* 67: 1317-1324
- Bashan Y, Mitiku G, Ziv-Vecht O, Levanony H (1991) Estimation of minimal numbers of *Azospirillum brasilense* using time-limited liquid enrichment combined with enzyme-linked immunosorbent assay. *Soil Biol Biochem* 23: 135-138
- Bashan Y, Holguin G, Lifshitz R (1993) Isolation and characterization of plant growth-promoting rhizobacteria. In: Glick BR, Thompson JE (eds) *Methods in plant molecular biology and biotechnology*. CRC BocaRaton, Fla, pp 331-345
- Bordeleau LM, Prevost D (1981) Quality of commercial legume inoculants in Canada. In: Clarke KW, Stephens JHG (eds) *Proceedings of the 8th North American Rhizobium conference*. University of Manitoba, Winnipeg, pp 562-565
- Carley HE, Watson RD (1966) A new gravimetric method for estimating root-surface areas. *Soil Sci* 102: 289-291
- Cassidy MB, Lee H, Trevors JT (1996) Environmental applications of immobilized microbial cells: a review. *J Ind Microbiol* 16: 79-101
- Del Gallo M, Negi M, Neyra CA (1989) Calcofluor- and lectin-binding exocellular polysaccharides of *Azospirillum brasilense* and *Azospirillum lipoferum*. *J Bacteriol* 171: 3504-3510
- DeLucca AJ, Connick WJ Jr, Fravel DR, Lewis JA, Bland JM (1990) The use of bacterial alginates to prepare biocontrol formulations. *J Ind Microbiol* 6: 129-134
- Digat B (1991) A new encapsulation technology for bacterial inoculants and seed bacterization. In: Keel C, Koller B, Dèfago G (eds) *Plant growth-promoting Rhizobacteria-progress and prospects*. International Organization for Biological and Integrated Control of Noxious Animals and Plants Bulletin, Zurich, pp 383-391
- Dubrovsky JG, Puente ME, Bashan Y (1994) *Arabidopsis thaliana* as a model system for the study of the effect of insulation by *Azospirillum brasilense* Sp-245 on root hairs growth. *Soil Biol Biochem* 26: 1657-1664
- Elsas JD van, Trevors JT, Jain D, Wolters AC, Heijnen CE, Overbeek LS van (1992) Survival of, and colonization by, alginate-encapsulated *Pseudomonas fluorescens* cells following introduction into soil. *Biol Fertil Soils* 14: 14-22
- Eskew DL, Focht DD, Ting IP (1977) Nitrogen fixation, denitrification, and pleomorphic growth in a highly pigmented *Spirillum lipoferum*. *Appl Environ Microbiol* 34: 582-585
- Fages J (1990) An optimized process for manufacturing an *Azospirillum* inoculant for crops. *Appl Microbiol Biotechnol* 32: 473-478
- Fages J (1992) An industrial view of *Azospirillum* inoculants: formulation and application technology. *Symbiosis* 13: 15-26
- Ganry F, Diem HG, Dommergues YR (1982) Effect of inoculation with *Glomus mosseae* on nitrogen fixation by field grown soybeans. *Plant Soil* 68: 321-329
- Glick BR, Bashan Y (1997) Genetic manipulation of plant growth promoting bacteria to enhance biocontrol of phytopathogens. *Biotechnol Adv* 15: 353-378
- Glick BR, Karaturovic DM, Newell PC (1995) A novel procedure for rapid isolation of plant growth promoting pseudomonads. *Can J Microbiol* 41: 533-536
- Hall PG, Krieg NR (1983) Swarming of *Azospirillum brasilense* on solid media. *Can J Microbiol* 29: 1592-1594
- Holguin G, Guzman MA, Bashan Y (1992) Two new nitrogen-fixing bacteria from the rhizosphere of mangrove trees, isolation, identification and in vitro interaction with rhizosphere *Staphylococcus* sp. *FEMS Microbiol Ecol* 101: 207-216
- Jung G, Mugnier J, Diem HG, Dommergues YR (1982) Polymer entrapped *Rhizobium* as an inoculant for legumes. *Plant Soil* 65: 219-231
- Kapulnik Y, Gafny R, Okon Y (1985) Effect of *Azospirillum* spp. inoculation on root development and N<sub>2</sub>O uptake in wheat (*Triticum aestivum* cv. Miriam) in hydroponic systems. *Can J Bot* 63: 627-631
- Le Tacon F, Jung G, Mugnier J, Michelot P, Mauperin C (1985) Efficiency in a forest nursery of an ectomycorrhizal fungus inoculum produced in a fermentor and entrapped in polymeric gels. *Can J Bot* 63: 1664-1668
- Levanony H, Bashan Y, Kahana ZE (1987) Enzyme-linked immunosorbent assay for specific identification and enumeration of *Azospirillum brasilense* Cd. in cereal roots. *Appl Environ Microbiol* 53: 358-364
- Ogoshi A, Kobayashi K, Homma Y, Kodama F, Kondo N, Akino S (eds) (1997) *Plant growth-promoting Rhizobacteria - present status and future prospects*. Faculty of Agriculture, Hokkaido University, Sapporo, p 483
- Paul E, Fages J, Blanc P, Goma G, Pareilleux A (1993) Survival of alginate-entrapped cells of *Azospirillum lipoferum* during dehydration and storage in relation to water properties. *Appl Microbiol Biotechnol* 40: 34-39
- Puente M-E, Bashan Y (1993) Effect of inoculation with *Azospirillum brasilense* strains on the germination and seedlings growth of the giant columnar cactus (*Pachycereus pringley*). *Symbiosis* 15: 49-60
- Reinhold B, Hurek T, Fendrick I (1985) Strain-specific chemotaxis of *Azospirillum* spp. *J Bacteriol* 162: 190-195
- Smith RS (1992) Legume inoculant formulation and application. *Can J Microbiol* 38: 485-492
- Thompson JA (1980) Production and quality control of legume inoculants. In: Begerson FJ (ed) *Methods for evaluating biological nitrogen fixation*. Wiley, New York, pp 489-533
- Trevors JT, Elsas JD van, Lee H, Overbeek LS van (1992) Use of alginate and other carriers for encapsulation of microbial cells for use in soil. *Micron Releases* 1: 61-b9