

HOW THE PLANT GROWTH-PROMOTING BACTERIUM *AZOSPIRILLUM* PROMOTES PLANT GROWTH—A CRITICAL ASSESSMENT

Yoav Bashan^{*,†} and Luz E. de-Bashan^{*,†}

Contents

1. Introduction	78
2. Major Mechanisms	80
2.1. Production of phytohormones	80
2.2. Nitrogen fixation	101
2.3. General improvement of root growth and enhanced uptake of minerals and water	106
2.4. Phosphate solubilization and mobilization and rock weathering	107
2.5. Mitigation of stresses	108
3. Other Proposed Mechanisms	113
3.1. Biological control	113
3.2. Nitric oxide	116
3.3. Nitrite	117
3.4. Signal molecules and enhanced proton extrusion from roots	117
3.5. <i>Azospirillum</i> nitrate reductase	119
3.6. Additive hypothesis	119
4. Concluding Remarks and a Proposal	120
Acknowledgments	122
References	122

* The Bashan Foundation, Corvallis, Oregon, USA

† Environmental Microbiology Group, Northwestern Center for Biological Research (CIBNOR), Colonia Playa Palo de Santa Rita, La Paz, B.C.S., Mexico

This review is dedicated to the memory of Dr. Wolfgang Zimmer (1958–2002) from Fraunhofer-Institute of Atmosphere Research in Garmisch-Partenkirchen, Germany. He intensively studied mechanisms of action by *Azospirillum* in the 1980s and 1990s.

Abstract

During the last 35 years of studies of *Azospirillum*–plant interaction, over 20 proposals were suggested for the mechanism of action by which *Azospirillum* spp., the most intensively studied plant growth-promoting bacteria, enhances plant growth. The proposals include a single phytohormone activity, multiple phytohormones, nitrogen fixation, assortments of small-sized molecules and enzymes, enhanced membrane activity, proliferation of the root system, enhanced water and mineral uptake, mobilization of minerals, mitigation of environmental stressors of plants, and direct and indirect biological control of numerous phytopathogens. By volume, the largest number of published information involves hormonal activities, nitrogen fixation, and root proliferation. After analyzing the accumulated knowledge, it was concluded that this versatile genus possesses a large array of potential mechanisms by which it can effect plant growth. Consequently, this review proposes the “Multiple Mechanisms Theory,” based on the assumption that there is no single mechanism involved in promotion of plant growth by *Azospirillum*, but a combination of a few or many mechanisms in each case of inoculation. These may vary according to the plant species, the *Azospirillum* strain, and environmental conditions when the interaction occurred. The effect can be cumulative, an “additive hypothesis” (proposed before), where the effects of small mechanisms operating at the same time or consecutively create a larger final effect on plant. Additionally, the observed effect on plant growth can be the result of a tandem or a cascade of mechanisms in which one mechanism stimulates another, yielding enhanced plant growth, such as the plausible relations among phytohormones, nitric oxide, membrane activities, and proliferation of roots. Finally, the growth promotion can also be a combination of unrelated mechanisms that operate under environmental or agricultural conditions needed by the crop at particular locations, such as mitigating stress (salt, drought, toxic compounds, adverse environment), and the need for biological control of or reducing pathogenic microflora.

1. INTRODUCTION

Since the rediscovery in the mid-1970s of the genus *Azospirillum* as a plant-associated bacteria by the late Johana Döbereiner and her collaborators in Brazil and its definition as a plant growth-promoting bacteria (PGPB; Döbereiner and Day, 1976), two main characteristics defined the genus; it fixes atmospheric nitrogen and produces phytohormones (Tien *et al.*, 1979). Consequently, these two features were considered, from the onset of plant–bacteria studies, as the cornerstone of the effect of this genus on plant growth and yield. Because *Azospirillum* is the most studied PGPB, excluding rhizobia, and reached commercialization in several countries, including Argentina, Mexico, India, Italy, and France (Díaz-Zorita and

Fernández-Canigia, 2009; Hartmann and Bashan, 2009), considerable knowledge has been accumulated during the last three decades, showing more and different facets of this interaction.

Despite intensive studies of the physiology and molecular biology of this genus, mostly as an easy-to-handle laboratory model of a rhizosphere bacterium, the exact mode of action of the bacteria on plants is not much clearer than it was decades ago (Bashan and Holguin, 1997; Bashan and Levanony, 1990; Bashan *et al.*, 2004). There are three facts that are beyond dispute, each with a reservation: (1) Most *Azospirillum* strains can fix nitrogen but only a fraction of it, if any at all, is transferred to the plant; (2) Many strains, but not all, produce several phytohormones *in vitro* and also a few in association with plants, but transfer of hormones is probably limited and was not always detected and only assumed to occur; (3) A general positive growth response in numerous plant species is evident in the majority of cases of inoculation, but the effect is not always apparent in terms of economic productivity (Díaz-Zorita and Fernández-Canigia, 2009; Okon and Labandera-Gonzalez, 1994). These concerns accelerate the research into alternative mechanisms.

The most apparent outcomes of most inoculations with *Azospirillum* are the major changes in plant root architecture. Inoculation can promote root elongation (Dobbelaere *et al.*, 1999; Levanony and Bashan, 1989), development of lateral and adventitious roots (Creus *et al.*, 2005; Fallik *et al.*, 1994; Molina-Favero *et al.*, 2008), root hairs (Hadas and Okon, 1987; Okon and Kapulnik, 1986), and branching of root hairs (Jain and Patriquin, 1985), some of which occurred in many plant species, consequently significantly increasing and improving their root system. It is generally accepted that these developmental responses in root morphology are triggered by phytohormones, possibly aided by their associated molecules. The fundamental question is: one hormone or several or a fine-tuned combination among several hormones?—all of which were produced by the bacterium, but mainly *in vitro*.

Inoculated plants absorbed more minerals and water, and in many cases, were more vigorous and greener, and showed enhanced plant growth. Several possible mechanisms were suggested to explain these phenomena, some with more experimental data than others. Yet, there is no definite agreement on exactly how the bacteria can effect plant growth. Is this a result of an input by the bacteria and can we manipulate it? Naturally, a multitude of proposed mechanisms claim to be the lead mechanism responsible for the observed effects on plant growth, and specifically, on plant yield, that in most cases, is the desired outcome and the main reason for inoculation. These questions are the driving force in *Azospirillum* research today, because if we have a clearer idea how the bacterium interacts with its host, we may envision ways to improve the interaction.

The aims of this chapter are to critically assess the large amount of knowledge on the possible plant promoting mechanisms of *Azospirillum* (Table 1, Fig. 1) and to present potential avenues for clarifications of this open question or at least some starting points for future research.

2. MAJOR MECHANISMS

2.1. Production of phytohormones

The ability to form plant hormones is a major property of many microorganisms and PGPB in general and specifically, species of *Azospirillum* that stimulate and facilitate plant growth (Tsavkelova *et al.*, 2006). This is believed to be part of the mutualistic relationships developed between plants and their associate bacteria. *Azospirillum* spp. are known for their ability to produce plant hormones, as well as polyamines and amino acids in culture media (Hartmann and Zimmer, 1994; Thuler *et al.*, 2003). Among these hormones, indoles, mainly indole-3-acetic acid (IAA; Spaepen *et al.*, 2007a), and gibberellins (GAs) of several kinds (Bottini *et al.*, 2004) may play a larger role. These phytohormones alter metabolism and morphology of plants, leading to better absorption of minerals and water, consequently larger and healthier plants. In the unicellular microalga *Chlorella vulgaris*, phytohormones lead to larger cell populations (de-Bashan *et al.*, 2008a). Thus far, hormonal effects are the mode of action for the largest volume of experimental data, and it is presented, justifiably or not, as the major (and sometimes as the sole) contribution of *Azospirillum* to plant growth.

The topic of IAA in PGPB in general and *Azospirillum* in particular and specifically the genes involved in synthesis of IAA were intensively, continuously, and excellently reviewed during the last decade (Costacurta and Vanderleyden, 1995; Dobbelaere *et al.*, 2003; Patten and Glick, 1996; Spaepen *et al.*, 2007a,b; Steenhoudt and Vanderleyden, 2000; Vande Brock and Vanderleyden, 1995), and therefore, this review will present only a few key points for and against their proposal.

2.1.1. Indole-3-acetic acid

IAA is a heterocyclic compound containing a carboxymethyl group (acetic acid) that belongs to the auxin phytohormone family. It is the best characterized and the most studied phytohormone and involved in numerous mechanisms in plant physiology. Auxins are responsible for division, extension, and differentiation of plant cells and tissues. Phytohormones of this group increase the rate of xylem and root formation; control processes of vegetative growth, tropism, floescence, and fructification of plants; and also affect photosynthesis, pigment formation, biosynthesis of various metabolites, and resistance to biotic stress factors. In microorganisms, in general, the

Table 1 General evaluation of proposals for mode of action on plants by *Azospirillum*

Proposal	Year proposed and significance with current knowledge ^a	Description	Evidence for	Arguments against and/or lack of evidence for	References (examples) ^b
Phytohormones	1979-present (+++)	<i>Azospirillum</i> can produce <i>in vitro</i> several phytohormones. External application of synthetic hormones or hormones purified from bacterial culture imitated the positive effects of <i>Azospirillum</i> on root development			Costacurta and Vanderleyden (1995), Spaepen <i>et al.</i> (2007a), Steenhoudt and Vanderleyden (2000), Vande Brock and Vanderleyden (1995)
IAA	1979-present (+++)	IAA is produced by the bacterium <i>in vitro</i> in large quantities and is attributed to affect numerous alterations in plant functions yielding eventually growth promotion. Significant knowledge about IAA metabolism and molecular	<ol style="list-style-type: none"> 1. Most strains of <i>Azospirillum</i> produce IAA <i>in vitro</i> 2. In several cases, IAA-attenuated mutants were ineffective compared to their wild-type parental strains 3. Application of IAA mimics <i>Azospirillum</i> inoculation on 	<ol style="list-style-type: none"> 1. Only several cases of direct involvement of IAA in growth promotion of plants are known 2. There are no IAA-deficient mutants 3. Most demonstrated cases are indirect or circumstantial 4. IAA is produced by plant cells and IAA detected in the plants were only indirectly induced, but not 	Baca <i>et al.</i> (1994), Barbieri and Galli (1993), Bothe <i>et al.</i> (1992), de-Bashan <i>et al.</i> (2008a), Dobbelaere <i>et al.</i> (1999), El-Khawas and Adachi (1999), Fallik <i>et al.</i> (1989), Gonzalez and Bashan (2000), Hartmann <i>et al.</i> (1983), Malhotra and Srivastava (2008), Molla <i>et al.</i> (2001), Omay <i>et al.</i> (1993),

(continued)

Table 1 (continued)

Proposal	Year proposed and significance with current knowledge ^a	Description	Evidence for	Arguments against and/or lack of evidence for	References (examples) ^b
		mechanism in the bacterium is known	<p>root morphology and growth promotion of plants and single cell algae</p> <p>4. IAA is involved in numerous functions in the plant cells and therefore might be a part of a cascade employing other mechanisms</p> <p>5. Elevated IAA was detected in inoculated plants</p> <p>6. IAA overproducing mutants showed stronger effect on plants</p>	<p>directly produced by the bacterium</p> <p>5. Direct transfer of bacterial IAA into plant cells and its functional consequences are still lacking</p> <p>6. Production of bacterial IAA in the plant was not demonstrated</p> <p>7. Cases of no evidence of correlation between capacity of IAA biosynthesis and root growth promotion</p> <p>8. Is additional IAA <i>in planta</i> better?</p> <p>9. Studies showing that <i>Azospirillum</i>-IAA biosynthesis alone cannot account for the overall growth promotion observed</p>	<p>Remans <i>et al.</i> (2008), Spaepen <i>et al.</i> (2007b, 2008), Zimmer <i>et al.</i> (1991)</p>

Gibberellins (GA)	1989–present (++)	GA affecting plant development in similar manner like auxins with several differences. GA promotes cell division and elongation and are involved in breaking dormancy	<ol style="list-style-type: none"> 1. GAs are synthesized and metabolized by <i>Azospirillum in vitro</i> 2. GA are produced <i>in planta</i> by <i>Azospirillum</i> 3. Inoculation of GA-deficient mutant dwarf rice mutants with <i>Azospirillum</i>-GA producer reversed dwarfism 	Insufficient evidence for involvement of bacterial GA in promoting growth	Bottini et al. (1989, 2004) , Cassan et al. (2009a,b) , Fulchieri et al. (1993) , Piccoli and Bottini (1994a,b) , Piccoli et al. (1997, 1999) , Perrig et al. (2007)
Cytokinins	1979 (UN)	Cytokinins are involved in cell enlargement and division, shoot and root morphogenesis and senescence	Cytokinins were produced by <i>Azospirillum in vitro</i>	No direct evidence in plants. Insufficient data.	Cacciari et al. (1989) , Horemans et al. (1986) , Strzelczyk et al. (1994) , Tien et al. (1979)
Abscisic acid (ABA)	2007–present (UN)	ABA is involved in response to environmental stress such as heat, water, and salt	<ol style="list-style-type: none"> 1. This compound was found <i>in vitro</i> in several strains 2. Interaction between GA and ABA in water-stress mitigation of plants 	Insufficient evidence for involvement of bacterial ABA in growth promotion	Cohen et al. (2008, 2009) , Perrig et al. (2007)

Table 1 (continued)

Proposal	Year proposed and significance with current knowledge ^a	Description	Evidence for	Arguments against and/or lack of evidence for	References (examples) ^b
Ethylene	2006-present (+)	Ethylene plays a role in breaking dormancy of seeds. Its main effect is in senescence of the plant	<ol style="list-style-type: none"> 1. Ethylene was found in culture filtrate of <i>A. brasilense</i> 2. Growth promotion was associated with low ethylene levels in tomato 3. Insertion gene of ACC-deaminase in <i>Azospirillum</i> improve plant growth 	Too few cases of ethylene involvement in <i>Azospirillum</i> , to compare with other PGPB, where it is a major mechanism	Holguin and Glick (2001, 2003), Perrig <i>et al.</i> (2007), Prigent-Combaret <i>et al.</i> (2008), Ribaudou <i>et al.</i> (2006)
Polyamines: cadaverine, putrescine, spermine, and spermidine	2003-present (UK)	Unclear function. Can act as growth regulating compounds	<ol style="list-style-type: none"> 1. These compounds were found <i>in vitro</i> 2. Application of cadaverine mitigated osmotic stress in rice 	Limited data	Cassan <i>et al.</i> (2009a), Perrig <i>et al.</i> (2007), Thuler <i>et al.</i> (2003)
Enhanced root growth combined with enhanced	1979-present (++++)	Inoculation caused a more developed root system that allows better uptake	<ol style="list-style-type: none"> 1. Enhanced root system is the most common phenotypical effect of inoculation with 	<ol style="list-style-type: none"> 1. Most data is descriptive and does not show whether the improvements are the cause or the 	Bashan <i>et al.</i> (1990), Jain and Patriquin (1984), Kapulnik <i>et al.</i> (1981, 1985b), Lin <i>et al.</i> (1983),

mineral and water uptake	of water and minerals	<p><i>Azospirillum</i> in most plant species</p> <ol style="list-style-type: none"> Enhanced mineral and water uptake by plants follow inoculation 	<p>results of other mechanisms</p> <ol style="list-style-type: none"> The wide range of enzymes related to these phenomena was only slightly studied Despite the large volume of information, relatively few strains were evaluated 	<p>Morgenstern and Okon (1987), Murty and Ladha (1988), Ogut and Er (2006), Sarig <i>et al.</i> (1988, 1992)</p>
Nitrogen fixation 1975-present (++)	Nitrogen fixation is a common feature of most <i>Azospirillum</i> species.	<ol style="list-style-type: none"> Following inoculation significant increase in total N in shoots and grain Many greenhouse and field experiments indicate some contribution of fixed N in the plant Inoculation commonly reduced the level of N fertilization needed for many plant species 	<p>Many studies showed little or minimal contribution of fixed nitrogen in the plant. Some systems showed none</p>	<p>Baldani and Baldani (2005), Bashan <i>et al.</i> (1989b, 2004), Choudhury and Kennedy (2004), Christiansen-Weniger (1992), Garcia de Salamone <i>et al.</i> (1997), Katupitiya <i>et al.</i> (1995a,b), Kennedy and Islam (2001), Kennedy <i>et al.</i> (2004), Mirza <i>et al.</i> (2000), Rodrigues <i>et al.</i> (2008), Saubidet and Barneix (1998), Sriskandarajah <i>et al.</i></p>

Table 1 (continued)

Proposal	Year proposed and significance with current knowledge ^a	Description	Evidence for	Arguments against and/or lack of evidence for	References (examples) ^b
			4. Enhanced nitrogenase activity in inoculated plants		(1993), Van Dommelen <i>et al.</i> (2009)
			5. The contribution of fixed N was apparent in many para-nodule systems		
Nitric oxide (NO)	2005–present (+)	NO is a free radical which participates in metabolic, signaling, defense, and developmental pathways in plants	1. <i>Azospirillum</i> can produce NO <i>in vitro</i> by different pathways 2. NO can modify root architecture	Limited data	Creus <i>et al.</i> (2005), Molina-Favero <i>et al.</i> (2007, 2008)
Nitrite production	1992 (UN)	<i>Azospirillum</i> can produce nitrite as part of its normal metabolism	1. Nitrite participated in plant growth promotion 2. Nitrite can cause sharp decrease in formation of lateral roots	Limited data	Bothe <i>et al.</i> (1992), Zimmer <i>et al.</i> (1988)

Nitrate reductase (NR)	1987 (UN)	An explanation for accumulating nitrogen following <i>Azospirillum</i> inoculation	<ol style="list-style-type: none"> 1. NR activity of wheat leaves was decreased by inoculation with some <i>Azospirillum</i> strains 2. Increase in nitrate assimilation 	Limited data	Boddey and Döbereiner (1988), Ferreira <i>et al.</i> (1987)
Phosphate solubilization and mineral weathering	1998-present (UN)	Solubilization of several rock minerals especially P making them available for the plant	Several strains can solubilize nonsoluble P and other minerals from rocks and stones	Limited data	Carrillo <i>et al.</i> (2002), Chang and Li (1998), Kamnev <i>et al.</i> (1999a, 2002b), Puente <i>et al.</i> (2004a,b, 2006), Rodriguez <i>et al.</i> (2004), Seshadri <i>et al.</i> (2000)
Effect on plant membranes and enhanced proton extrusion	1989-present (UN)	Inoculation induces root cell membranes to release protons	<ol style="list-style-type: none"> 1. Short exposure of roots to <i>A. brasilense</i> significantly enhanced the proton efflux of the root 2. Inoculation significantly reduced the membrane potential in every root part 	<ol style="list-style-type: none"> 1. Signal molecules in bacteria that might affect membranes were not identified 2. Mobilization of ions via the affected membranes was not studied 	Alen'kina <i>et al.</i> (2006), Amooaghaie <i>et al.</i> (2002), Antonyuk <i>et al.</i> (1993, 1995), Bashan (1990, 1991), Bashan and Levanony (1991), Bashan <i>et al.</i> (1989a, 1992), Carrillo <i>et al.</i> (2002), Nikitina <i>et al.</i> (2004)

Table 1 (continued)

Proposal	Year proposed and significance with current knowledge ^a	Description	Evidence for	Arguments against and/or lack of evidence for	References (examples) ^b
			3. Inoculation changed the phospholipid content in plant membranes 4. <i>Azospirillum</i> produces lectins. Some can cause change in growing cells mitosis 5. Wheat germ agglutinin from plants enhanced several metabolic pathways in <i>Azospirillum</i>		
Mitigation of environmental stress	1988-present	Best effects of inoculation occurred when plants are grown under suboptimal conditions			Bashan and Holguin (1997), Bashan and Levanony (1990), Bashan <i>et al.</i> (2004)
Salinity	1997-present (++)	Inoculated plant under saline condition grow better	1. Inoculation improved germination, plant development	Missing information on: 1. Relation between salt tolerance of the bacterium and those of the plant	Bacilio <i>et al.</i> (2004), Barassi <i>et al.</i> (2006), Creus <i>et al.</i> (1997), Hamdia and

			<ol style="list-style-type: none"> 2. Increases in content of water, chlorophyll, essential minerals, proteins, amino acids, enhanced uptake of K and Ca, NR, and nitrogenase 3. Restricted Na uptake 	<ol style="list-style-type: none"> 2. What are the physiological mechanisms involved? 3. How do the bacteria induce these effects? 	<p>El-Komy (1997), Hamdia <i>et al.</i> (2004)</p>
Drought	1988-present (++)	Inoculated plant under drought or osmotic stress are growing better	Inoculation improved plant growth, reduce grain loss, improve water content, increased turgor pressure, positive effect on cell wall elasticity, higher Mg, K, and Ca in grains, and improve fatty acid distribution profile	Too little data about the physiological mechanisms involved	<p>Alvarez <i>et al.</i> (1996), Creus <i>et al.</i> (1998, 2004), El-Komy <i>et al.</i> (2003), Pereyra <i>et al.</i> (2006), Sarig <i>et al.</i> (1990)</p>
Metal toxicity	2000-present (+)	Reduction in toxicity to plants	<ol style="list-style-type: none"> 1. The bacterium tolerates medium levels of metals 	Inoculation does not provide full protection against metal toxicity	<p>Belimov and Dietz (2000), Belimov <i>et al.</i> (2004), Kamnev <i>et al.</i></p>

(continued)

Table 1 (continued)

Proposal	Year proposed and significance with current knowledge ^a	Description	Evidence for	Arguments against and/or lack of evidence for	References (examples) ^b
			2. Inoculation allows plants to grow in metal contaminated soils and in mine tailings		(2005, 2007), Lyubun <i>et al.</i> (2006)
Humic acid toxicity	2003 (UN)	Reduction in toxicity to plants	1. Improved germination and plant growth at elevated humic acids 2. Consumption of humic acid by the bacterium	Inoculation does not provide full protection against toxicity	Bacilio <i>et al.</i> (2003)
pH and tryptophan in aquatic environment	2005-present (+)	Inoculation allows microalgae to grow under unfavorable aquatic conditions	Inoculated microalgae can grow in high pH and toxic levels of tryptophan	Unknown	de-Bashan and Bashan (2008), de-Bashan <i>et al.</i> (2005)
High light intensity	2006-present	Inoculation allows plants to grow under high light intensity	1. Inoculated wheat plants produced photoprotective photosynthetic pigments	Unknown	Bashan <i>et al.</i> (2006), de-Bashan <i>et al.</i> (2008b)

Herbicide	1997 (UN)	Reduction in toxicity to plants	2. Inoculation allowed microalgae to grow under extreme light intensities Cotton plants were partially protected from the herbicide 2-4-D	Unknown	Feng and Kennedy (1997)
Biological control of pathogens	1990-present (++)	Indirect effects on plant growth reducing the deleterious effects of pathogens	1. <i>Azospirillum</i> produces a variety of inhibitory substances 2. Inhibits germination and development of parasitic weeds 3. Can compete with phytopathogens 4. Inhibits development of microfauna and insects 5. Inhibits foliar bacterial diseases and soil-borne fungal pathogens	1. Most studies are descriptive 2. Almost all mechanisms were not studied or are in initial stage	Bashan and de-Bashan (2002a), Dadon <i>et al.</i> (2004), Gonçalves and de Oliveira (1998), Kavitha <i>et al.</i> (2003), Khan and Kounsar (2000), Romero <i>et al.</i> (2003), Sudhakar <i>et al.</i> (2000)

Table 1 (continued)

Proposal	Year proposed and significance with current knowledge ^a	Description	Evidence for	Arguments against and/or lack of evidence for	References (examples) ^b
Additive hypothesis	1990–present	The effects of small mechanisms operating at the same time or consecutively create a larger final effect on plant			Bashan and Dubrovsky (1996) , Bashan and Levanyon (1990)
Multiple mechanisms	2010	A combination of a few or many mechanisms in each case of inoculation			This essay

^a +++, possibly major mechanism; ++, possibly moderate; +, possibly minor; UN, unknown.

^b More comprehensive literature are listed in the text.




Figure 1 Mechanisms by which *Azospirillum* spp. may enhance plant growth and their possible interactions grouped as biological processes. Circles represent processes containing experimental data. Squares represent theories. Size of a circle represents its relative importance according to current data. Solid arrow: mechanism(s) that can fully create the observed growth promotion; dash arrow: mechanism(s) that can only partially explain the observed growth promotion. Simple arrows: proven interactions among different mechanisms; double-line arrow: direct production of molecules or processes by the bacterium cell; ?: unproven as yet, or partially proven pathway.

three known pathways of IAA biosynthesis are related to tryptophan metabolism (amino acid frequently found in plant exudates; Costacurta and Vanderleyden, 1995; Patten and Glick, 1996). Omission of tryptophan from the culture medium decreases the level of IAA synthesis by the culture's microorganisms. Addition of exogenous tryptophan (or, more rarely, tryptamine) may augment auxin biosynthesis by an order of magnitude or even greater. The known routes of IAA biosynthesis includes: (1) IAA formation via indole-3-pyruvic acid (IPyA) and indole-3-acetaldehyde; (2) Conversion of tryptophan into indole-3-acetaldoxyme and indole-3-acetonitrile (IAN); and (3) IAA biosynthesis via indole-3-acetamide formation (IAM; Zakharova *et al.*, 1999). It has been reported that a tryptophan-independent pathway, more common in plants, was also found in azospirilla (Carreño-Lopez *et al.*, 2000; Prinsen *et al.*, 1993). However, the contribution of this pathway to IAA biosynthesis is questionable, and the mechanisms are largely unknown.

Involvement of tryptophan in IAA production by *Azospirillum* has been known for a long time (Reynders and Vlassak, 1979). A key gene *ipdC* encodes for indole pyruvate decarboxylase. This is a key enzyme in the IAA synthesis pathway by *A. brasilense* that mediates conversion of IPyA into indole-3-acetaldehyde; its presence presented conclusive evidence for the IPyA pathway in this bacterium (Costacurta *et al.*, 1994). Zimmer *et al.* (1998) isolated the *ipdC* gene from strain Sp7 of *A. brasilense* and showed tryptophan-dependent stimulation of gene expression in this bacterium. These two findings were later confirmed by IAA production by several strains of *Azospirillum* where production depended on the type of culture media and availability of tryptophan as a precursor (e.g., El-Khawas and Adachi, 1999; Malhotra and Srivastava, 2006, 2008). The pH of the culture medium has a significant effect on the amount of IAA produced (Ona *et al.*, 2003). Release of large amounts of IAA by *Azospirillum* spp. cultures is probably controlled by the stationary phase of the bacteria cells after depletion of the carbon source in the medium used in batch culture. Depletion of the carbon source reduces growth (Ona *et al.*, 2003, 2005). Assessment of possible precursors (indole, anthranilic acid, and tryptophan) for IAA formation in *A. brasilense* Sp245 revealed a high motive force for tryptophan synthesis from chorismic acid and for IAA synthesis from tryptophan (Zakharova *et al.*, 1999). Vitamins may also play a role in the regulation of IAA synthesis in *A. brasilense*. Very low levels of B vitamins, especially pyridoxine and nicotinic acid, increased production of IAA in *A. brasilense* (Zakharova *et al.*, 2000).

To demonstrate direct involvement of IAA produced by *A. brasilense* on plant growth, it would be preferable, if not essential, to use IAA-deficient mutants. It is relatively straightforward to obtain IAA overproducing mutants (Hartmann *et al.*, 1983) but, so far, almost impossible to obtain IAA-deficient mutants. This occurs because of the different pathways that *Azospirillum* spp. has to produce IAA (Spaepen *et al.*, 2007a; Zakharova *et al.*, 1999). For example, in most mutants, the unstable indole pyruvic acid spontaneously breaks down and produces some IAA (Steenhoudt and Vanderleyden, 2000). These IAA-attenuated mutants produce 0.2–10% of the level of IAA produced by the wild type, sometimes even more. Quite a few of these strains were found or constructed and used. A strain of *A. irakense* released about 10 times less IAA into the medium than *A. brasilense* Sp7 (Zimmer *et al.*, 1991). Two mutants of *A. brasilense* produced 2–5% of the IAA produced by the parental strains (Prinsen *et al.*, 1993, Vande Broek *et al.*, 1999). Mutants of *A. brasilense* and *A. lipoferum* that were modified to include the *gfp* (green fluorescent protein) gene produced less than 0.25% IAA of their parental strains (Bacilio *et al.*, 2004; Rodriguez *et al.*, 2006) and mutant of *A. brasilense* Sp6, carrying another Tn5 insertion in the *ipdC* gene, produced less than half the IAA of its parental strain (Barbieri and Galli, 1993). Recently, an *ipdC*-knockout mutant was found to produce

only 10% of the wild-type IAA production level (Spaepen *et al.*, 2007b). Furthermore, when the endogenous promoter of the *ipdC* gene was replaced by either a constitutive or a plant-inducible promoter and both constructs were introduced into the wild-type strain, the introduction of these recombinant *ipdC* constructs improved the growth-promoting effect of *A. brasilense* (Spaepen *et al.*, 2008).

IAA is produced during all stages of culture growth and well after the stationary phase (Malhotra and Srivastava, 2009). This feature makes the bacterium especially qualified for plant growth promotion when the effect last weeks or months after inoculation. Consequently, IAA production by *Azospirillum* sp. was proposed to play a major role in growth promotion and even more auxin-type molecules were detected in *Azospirillum*, such as indole butyric acid (IBA; Fallik *et al.*, 1989), indole lactic acid (Crozier *et al.*, 1988), indole acetamide (Hartmann *et al.*, 1983), indole acetaldehyde (Costacurta *et al.*, 1994), indole ethanol and indole methanol (Crozier *et al.*, 1988), and phenyl acetic acid (Somers *et al.*, 2005). Nonetheless, when compared to the large base of knowledge on IAA production by the bacterium cell, a far smaller volume of indirect and direct evidence regarding the effect of IAA of bacterial origin in plants has been published.

In general, morphological changes in roots, following *Azospirillum* inoculation, were mimicked by applying a combination of plant growth substances, which point to involvement of an auxin produced by *Azospirillum* for root proliferation and consequent plant growth promotion (for reviews, see Bashan and Holguin, 1997; Bashan *et al.*, 2004). Specific evidence for the involvement of auxins in promoting plant growth includes elevated IAA and IBA in *Azospirillum*-inoculated maize plants (Fallik *et al.*, 1989). Addition of filter-sterilized culture supernatants to rice roots grown in hydroponic tanks increased root elongation, root surface area, root dry matter, and development of lateral roots and root hairs, compared with untreated roots. Higher concentrations of the supernatant strongly inhibited root elongation, lateral root development, and caused nodule-like tumors on the roots (El-Khawass and Adachi, 1999). Similarly, a cell-free supernatant of *A. brasilense* Cd applied to soybean plants induced many roots and increased root length (Molla *et al.*, 2001). Inoculation of wheat with wild strains of *A. brasilense* Sp245 and Sp7 led to an exceptional decrease in root length and increase in root hair formation, as is common with such inoculations. The effect on root morphology was further enhanced by adding tryptophan; this could be mimicked by replacing *Azospirillum* cells with IAA (Dobbelaere *et al.*, 1999). Exogenous application of IAA to bean roots resembled responses of these plants to inoculation with *Azospirillum* (Remans *et al.*, 2008). Similarly, application of IAA directly to growing cells of the freshwater microalgae *C. vulgaris* mimicked cell proliferation induced by *Azospirillum* (Gonzalez and Bashan, 2000). More direct evidence for the importance of IAA was provided when several IAA-attenuated mutants

were compared with their parental wild types for their effect on the growth of this microalga, when only the wild types were capable of promoting growth. Yet, adding culture filtrate of these wild types to cultures of IAA-attenuated mutants, incapable of inducing microalgal growth, restored their effect on microalgal growth (de-Bashan *et al.*, 2008a). A mutant of *A. brasilense* with low production of phytohormones, but high N₂-fixation activity, did not enhance root growth over uninoculated controls. In contrast, a mutant with increased phytohormone production significantly affected root morphology. In general, increased plant biomass and N₂-fixation were recorded in strains having increased production of indole compounds (Kundu *et al.*, 1997). Further study of the contribution of auxin biosynthesis by *A. brasilense* in altering root morphology and root proliferation showed that inoculation of wheat seedlings with an *A. brasilense* Sp245 strain, carrying a mutation in the *ipdC* gene, which did not cause shorter roots or stimulate root hair formation, in contrast to inoculation with the wild type (Dobbelaere *et al.*, 1999). The insertion of the heterologous IAM pathway, consisting of the *iaaM* and *iaaH* genes into *A. brasilense* SM increased IAA levels by threefold and the engineered strain showed a superior effect on the lateral branching of sorghum roots, as well as its dry weight when compared with the wild-type strain (Malhotra and Srivastava, 2006).

Several studies showed no evidence of correlation between the capacity for IAA synthesis by *A. brasilense* and the effects on observed root growth promotion (Bothe *et al.*, 1992; Harari *et al.*, 1988; Kapulnik *et al.*, 1985a). Additionally, several studies showed that *Azospirillum*-IAA biosynthesis alone cannot account for the overall growth stimulatory effect observed (for a review, see Spaepen *et al.*, 2007a).

In summary, although evidence of IAA production in *Azospirillum* spp. is the most comprehensive and documented from all hormones or suggested mechanisms, the direct evidence of involvement of this hormone as the *sole* mechanism by which the bacteria affect plant growth is, in our opinion, unproven, although it is very likely that IAA is involved in many of the interactions of this genus. It is feasible though to consider a hormonal effect in very early stages of germination. Most stains of *Azospirillum*, when fermented as an inoculant, are capable of producing IAA and other growth regulators at a concentration sufficient to produce morphological and physiological change in young seed tissues. Such initial "phytohormonal shock" would be the first contact between the bacterial inoculant and the seed and would not necessarily depend on the presence of bacteria. However, the presence of live bacteria may contribute to *in situ* phytohormone production over a longer term. According to this concept, bacterial phytostimulation would be crucial in early developmental stages (germination and initial seedling growth) and will be complementary to other mechanisms operating at later stages of *Azospirillum* interaction with plants (Cassan *et al.*, 2009b), as summarized in Section 3.6.

2.1.2. Gibberellins and abscisic acid

GAs, diterpenoid acids that are synthesized by the terpenoid pathway, are hormones (over 120 types have been found in plants, fungi, and bacteria) that control growth and a wide variety of other plant developmental processes similar to auxins. Primarily, they promote cell division and elongation, but without inhibitory effects presented by some auxins. Additionally, GAs are involved in the natural process of breaking dormancy during seed germination. GAs in the seed embryo signal starch hydrolysis by inducing the synthesis of the enzyme α -amylase in the aleurone cells. This enzyme hydrolyzes starch into glucose; the glucose is used for energy by the seed embryo. GAs cause higher levels of transcription of the gene coding for the α -amylase enzyme to stimulate the enzyme synthesis (Richards *et al.*, 2001). Despite this major role and the fact that *A. brasilense* is known to enhance germination of wheat and soybean seeds (Bacilio *et al.*, 2003; Cassan *et al.*, 2009b), GAs, so far, were not directly linked to this phenomenon. It was only shown that improved seed germination coincides with high GA production in cultures by this bacterium (Cassan *et al.*, 2009b).

Azospirillum has the capacity to synthesize and metabolize GAs *in vitro* (Bottini *et al.*, 1989; Piccoli and Bottini, 1994a,b; Piccoli *et al.*, 1996, 1997) and *in planta* (Bottini *et al.*, 2004; Cassan *et al.*, 2001a,b, and references cited therein). A growth promotion effect of *Azospirillum* spp. on plants has been suggested to be partially caused by the production of GAs by the bacterium as has occurred with other PGPB (for a review, see Bottini *et al.*, 2004). Several studies support this proposal. When a GA-producing strain of *A. lipoferum* was cultured in the presence of glucosyl ester or glucoside of GA A₂₀, both conjugates were hydrolyzed. These *in vitro* results support the hypothesis that growth promotion in plants induced by inoculation with *Azospirillum* results from a combination of GA production and GA-glucoside/glucosyl ester deconjugation by the bacterium (Piccoli *et al.*, 1997). The effect of water potential or concentration of O₂ on growth and GA A₃ (the main GA identified in *Azospirillum*) production in *A. lipoferum* showed that this GA produced by each culture was reduced severely at high water potentials or low O₂ concentrations. At the highest water potential concentration, GA A₃ was reduced by ~50%, despite a 90% reduction in cell numbers. This indicates an increase in the amount of GA A₃ produced per cell with increasing water potential (Piccoli *et al.*, 1999). Involvement of GA A₃ produced by *Azospirillum* spp. in promoting growth of maize was also suggested (Lucangeli and Bottini, 1997). *A. brasilense* Cd and *A. lipoferum* USA 5b promoted elongation of root sheaths with two single genes in GA-deficient dwarf rice mutants, dy and dx, when the inoculated seedlings were supplied with [17, 17-²H₂] GA A₂₀-glucosyl ester. This growth resulted from GA metabolism by the bacteria in the dx mutant and by the rice plant

and microorganism in the dy mutant. In the dy mutant, inoculation by both bacterial strains reversed dwarfism in seedlings incubated with $[17, 17\text{-}^2\text{H}_2]$ GA A₂₀, forming $[17, 17\text{-}^2\text{H}_2]$ GA A₁. It is possible that the bacterial enzyme responsible for these phenomena is 2-oxoglutarate-dependent dioxygenase, similar to those of plants (Cassan *et al.*, 2009a,b).

Initial studies on the effect of *Azospirillum* spp. on plants linked GAs and bacteria-produced abscisic acid (ABA), a common isoprenoid phytohormone usually synthesized in all plant parts. ABA is ubiquitous and produced by higher plants, algae, and fungi (Zeevaart, 1999) and as a by-product of chemically defined cultures of *A. brasilense* Sp245 (Cohen *et al.*, 2008). ABA originated from its role in the abscission of leaves of only a few plant species but its main role in plants is as a response phytohormone to environmental stress, such as decreased soil water potential and heat, water, and salt stresses. ABA produced in roots is then translocated by transpiration in the xylem to the leaves, where it rapidly alters the osmotic potential of stomata guard cells, causing them to shrink and stomata to close. The ABA-induced closure of stomata reduces transpiration, preventing further water loss in times of low water availability (Bartels and Sunkar, 2005). In this stress mitigation process, ABA–GAs were investigated with *Azospirillum* inoculation even though ABA and GAs have antagonistic roles in many processes of plant growth (Achard *et al.*, 2006; Nemhauser *et al.*, 2006).

The effects of *A. lipoferum* in maize plants, in which ABA and GA synthesis were diminished by inhibitors of their own biosynthetic pathways (ABA by fluridone and GA by Ca-prohexadione) and subjected or not to drought stress, were measured. Application of fluridone diminished growth of well-watered plants similar to the effect of drought and *A. lipoferum* inoculation completely reversed this effect. The relative water content of the fluridone-treated and drought-stressed plants was significantly lower, and this effect was completely neutralized by *A. lipoferum*. The results suggest that ABA produced by the bacterium may account, at least partially, for the amelioration of growth parameters in drought-stressed and fluridone-treated plants. Similarly, growth was diminished in plants subjected to drought and treated with Ca-prohexadione, alone or combined with fluridone, even though ABA levels were higher. The results suggest that ABA and GAs participate in alleviating water stress of plants by the presence of *A. lipoferum* (Cohen *et al.*, 2009).

So far, the results indicate that, among the mechanisms involved in water-stress alleviation of plants by *Azospirillum*, is the production of stress-type hormones such as ABA (Cohen *et al.*, 2008) along with growth promoters, such as auxins (Costacurta and Vanderleyden, 1995) and GAs (Bottini *et al.*, 2004). Similar to the studies on IAA, there is far more information about GA metabolism in the bacterium than the effect of bacterial-produced GA in plants where the information about the involvement of ABA is still at an embryonic stage.

2.1.3. Polyamines

The newest compound involved in promoting growth by *Azospirillum* spp. is the polyamine cadaverine synthesized from lysine. Polyamines are low-molecular-weight organic compounds having two or more primary amino groups ($-NH_2$). Polyamines are known to be synthesized in cells via highly regulated pathways, yet, their actual function is not entirely clear. If cellular polyamine synthesis is inhibited, usually cell growth is stopped or severely inhibited. Application of exogenous polyamines restores the growth of these cells. Most eukaryotic cells have a polyamine transporter system on their cell membranes that facilitates internalization of exogenous polyamines. Polyamines serve as growth regulating compounds (Kuznetsov *et al.*, 2006); among them, cadaverine has been correlated with root growth promotion in pine and soybean (Gamarnik and Frydman, 1991; Niemi *et al.*, 2001), response to osmotic stress in turnip (Aziz *et al.*, 1997), and controlling stomata activity in *Vicia faba* beans (Liu *et al.*, 2000). *A. brasilense* strain Az39, which is a widely used as a wheat and maize inoculant in Argentina, is known to produce polyamines such as spermidine and spermine (Perrig *et al.*, 2007), and putrescine (Thuler *et al.*, 2003) in culture, and also produce cadaverine in chemically defined medium supplemented with the precursor L-lysine and in rice plants inoculated with this strain. Application of cadaverine mitigated osmotic stress in rice seedlings, based on improved water status and decreased production of ABA in inoculated seedlings (Cassan *et al.*, 2009a). Cadaverine was proposed as a contributing factor to the whole plant response to *Azospirillum* inoculation, summarized in Section 3.6 (Bashan *et al.*, 2004).

2.1.4. Cytokinins

Cytokinins are a class of purine-type phytohormones that promote cell division, shoot and root morphogenesis, chloroplast maturation, cell enlargement, auxiliary bud release, and senescence. The ratio of auxin to cytokinin is crucial during cell division and differentiation of plant tissues. Auxin is known to regulate the biosynthesis of cytokinin. The adenine-type cytokinins represented by kinetin, zeatin, and 6-benzylaminopurine occur in plants.

Cytokinins are produced in defined culture medium by many rhizosphere bacteria (Barea *et al.*, 1976), including *Azospirillum* (Cacciari *et al.*, 1989; Horemans *et al.*, 1986; Strzelczyk *et al.*, 1994; Tien *et al.*, 1979). Cytokinins from bacteria might affect plant growth positively or negatively. Apart from initial results of plants inoculated with *Azospirillum*, it is questionable if cytokinins, on their own, modified the root morphology observed in many *Azospirillum* inoculation models or if it is the levels of combination with auxin and GAs that induced the observed effect. It is hypothesized (F. Cassán, personal communication) that the contribution of

cytokinins is of some, yet undefined, importance when *Azospirillum* is combined with *Bradyrhizobium* for inoculation of soybeans. Recently, nod factors in soybean were shown as not essential for nodulation and that some strains of *Bradyrhizobium* use purines (cytokinins) as an alternative option for nodulation (Giraud *et al.*, 2007). *Azospirillum* as a potential producer of cytokinins might support this type of nodulation. It is commonly observed that inoculation with *Azospirillum* increased nodulation (Schmidt *et al.*, 1988; Yahalom *et al.*, 1990). Yet, this hypothesis is still an open proposal for future research.

2.1.5. Ethylene

During most phases of plant growth, ethylene production is minimal. Ethylene plays a major role in germination by breaking the dormancy of seeds; however, a high level of ethylene concentration inhibits subsequent root elongation. High levels of ethylene may be synthesized as a response to biological or environmental stresses, causing wilting and senescence (Glick *et al.*, 1999). Controlling ethylene levels, often by lowering them, prevents significant economic losses in agriculture. One of the precursors of ethylene synthesis is the enzyme 1-aminocyclopropane-1-carboxylic acid (ACC) deaminase. ACC-deaminase is a key enzyme, commonly found in many soil microorganisms and PGPBs and capable of degrading ACC. Thus, lowering ethylene levels in plants can be considered as having potential for promoting growth (Glick *et al.*, 1999). Wild strains of *Azospirillum* spp. do not have ACC-deaminase; nevertheless, some strains can produce ethylene (Perrig *et al.*, 2007). A single exception to this role is *A. lipoferum* strain 4B that possesses the ACC-deaminase structural (*acdS*) gene (Prigent-Combaret *et al.*, 2008). This gene of the PGPB *Enterobacter cloacae* UW4 was inserted in *A. brasilense* Cd and Sp245. Roots of canola and tomato seedlings, plants sensitive to ethylene, were significantly longer in plants inoculated with the *A. brasilense* transformants than plants inoculated with nontransformed strains of the same bacterium (Holguin and Glick, 2001). In a further study, they speculated that a construct with the ACC-deaminase gene under control of a constitutive promoter weaker than the *lac* promoter, might impose less metabolic load on *Azospirillum*. The *acdS* gene was cloned under the control of a tetracycline-resistance gene promoter: *A. brasilense* Cd transformants holding *acdS* fused to the Tetr gene promoter showed lower ACC-deaminase activity than transformants with *acdS* controlled by the *lac* promoter. However, *acdS* controlled by the Tetr gene promoter exerted less metabolic load on *A. brasilense* Cd transformants than *acdS* controlled by the *lac* gene, resulting in increased IAA synthesis, growth rate, and survival of tomato leaf surfaces and ability to promote growth of seedlings (Holguin and Glick, 2003). A proposal that growth promotion triggered by inoculation with *A. brasilense* involves a signaling pathway that has ethylene as a central, positive regulator was published. The evidence is

based on higher levels of IAA and ethylene in inoculated plants. Exogenously supplied ethylene mimicked the effect of inoculation, and the addition of an inhibitor of its synthesis or of its physiological activity completely blocked promotion of growth by *A. brasilense* (Ribaudó *et al.*, 2006). Taken together, all of this may show that the involvement of ethylene in promoting growth by *Azospirillum* is probably small.

2.2. Nitrogen fixation

Since nitrogen fixation was the original proposed major mechanism by which *Azospirillum* affected plant growth (Okon *et al.*, 1983), considerable information has been published on this mechanism (for reviews, see Baldani and Baldani, 2005; Bashan and Holguin, 1997; Bashan and Levanony, 1990; Bashan *et al.*, 2004; Choudhury and Kennedy, 2004; Kennedy and Islam, 2001; Kennedy *et al.*, 2004, and references therein). The reason is that, following inoculation, there is a significant increase in the total N in shoots and grains of inoculated plants (Kapulnik *et al.*, 1981 and references in the above reviews). Incorporation of atmospheric nitrogen into the host plant by *Azospirillum* was evaluated initially by the acetylene reduction assay. However, conclusive proof that plants derive some of their N from the atmosphere came from the use of isotopic $^{15}\text{N}_2$ and ^{15}N -dilution techniques. The original seven species of this genus are diazotrophs (Bashan *et al.*, 2004). Most new species, but not all, are defined as nitrogen-fixers, either as free-living bacteria or in association with plants and participate in several transformations in the nitrogen cycle (Doroshenko *et al.*, 2007; Eckert *et al.*, 2001; Mehnaz *et al.*, 2007a,b; Peng *et al.*, 2006).

Subsequently, a very large volume of information on nitrogen-fixation mechanism in the association was published (for a review, see above). Taken together, the evidence collected during the last three decades concerning this mechanism has generated a substantial controversy. On one hand stands the numerous greenhouse and field experiments that repeatedly demonstrate some contribution of fixed nitrogen (measured as transfer of $^{15}\text{N}_2$). This was combined with more common observations that inoculation, commonly and significantly, reduced the required doses of nitrogen fertilization for cultivation of many plant species. Evidence that nitrogen fixation contributes to the N balance of plants is based on the common observation of an increase in nitrogenase activity within inoculated roots, a microbial enzyme that does not exist in plants. This well-documented enzymatic activity in *Azospirillum* is of sufficient magnitude to account for the increase in total N yield of inoculated plants if *all* the fixed N is incorporated into the plants (Kennedy *et al.*, 1997; for earlier studies Bashan and Holguin, 1997 and references therein). On the other hand, many studies show that the contribution of nitrogen fixation by *Azospirillum* to the plant is minimal and ranged, at best, from 5% to 18% of the total N increase in the plant. In many

of those studies, the contribution was smaller than 5% or null. Hence, it is an open debate to this day (Bashan and Holguin, 1997; Bashan and Levanony, 1990; Bashan *et al.*, 2004). These findings almost caused an abandonment of nitrogen-fixation aspects of *Azospirillum*, except for continuing pure genetic studies.

Several confirmatory reports about the contribution of fixed nitrogen by *Azospirillum* to plants, similar in nature to reports of earlier years, illustrate the controversy. The ^{15}N isotope dilution technique indicated that there were significant biological N_2 -fixation contributions to two genotypes of maize that showed similar increases in grain yield when they were inoculated with a mixture of *Azospirillum* strains or fertilized with the equivalent of 100 kg N ha^{-1} . These plant genotypes had a large increase in total N. This suggests that the yield response resulted from increased acquisition of nitrogen, but not from bacterial nitrate reductase (NR); NR^- mutants generally caused plant responses similar to those of the parent strains (Garcia de Salamone *et al.*, 1997). The ability of the bacteria to transfer fixed nitrogen from the atmosphere to wheat plants was tested using a $^{15}\text{N}_2$ -enriched atmosphere. Labeled fixed nitrogen was detected in plant growth media and roots and shoots of wheat grown for 26 days in a $^{15}\text{N}_2$ -enriched atmosphere, but the highest levels of ^{15}N were detected in wheat shoots. Ammonia or nitrate supplied to plants did not repress $^{15}\text{N}_2$ -fixation (Ruppel and Merbach, 1997). Relationships of 12 *A. brasilense* strains with roots of a wheat cultivar were studied. They were compared for responses in root colonization, growth stimulation, and nitrogen supply to the plant. All strains colonized the root surface and interior. Most strains stimulated plant growth, but to different degrees. Some strains increased the total nitrogen in roots and leaves up to 80% over noninoculated plants, while others produced no effect on nitrogen content. Inoculation of five wheat cultivars with the most efficient strain for nitrogen fixation resulted in increased growth and nitrogen content, but the effects varied among the cultivars. These results suggest that a potential exists for *A. brasilense* to supply considerable nitrogen to wheat plants, probably dependent on specific bacteria-cultivar interaction (Saubidet and Barneix, 1998).

Apparently, dismissal of nitrogen fixation as a possible mechanism for promoting plant growth by *Azospirillum* in the 1990s was premature and additional greenhouse studies in the last decade showed significant and direct contribution of nitrogen fixation. Measurement of nitrogen fixation after inoculation with *A. lipoferum* and *A. brasilense* in rice showed that the N derived from the atmosphere were 20.0% (*A. lipoferum*) and 19.9% (*A. brasilense*) in basmati rice and 58.9% (*A. lipoferum*) and 47.1% (*A. brasilense*) in super-basmati rice (Mirza *et al.*, 2000). Using an *in vitro* model (*A. brasilense* and wheat) within 70 h after inoculation, insignificant amounts of newly fixed N were transferred from an ammonia-excreting strain of *A. brasilense* to the shoot tissue of wheat. Adding malate (a preferred

carbon source for *Azospirillum*), transfer of nitrogen to the shoots increased 48-fold, which indicates that 20% of nitrogen in the shoot was derived from nitrogen fixation. Apparently, the inability of the host plant to release sufficient carbon into the rhizosphere is a significant constraint on the development of the *A. brasilense*–wheat association. Perhaps wheat with an increased release of photosynthate to the rhizosphere should be a priority for improving effectiveness of the association (Wood *et al.*, 2001). Inoculation of strains of *A. amazonense* on rice increased grain dry matter and nitrogen accumulation at maturation. Contributions from nitrogen fixation were up to 27% of the contribution to the plant. Promotion of growth by *A. amazonense* for these rice plants was primarily a response to nitrogen fixation (Rodrigues *et al.*, 2008). Finally, winter wheat inoculated with *A. brasilense* having a point mutation in the ammonium binding site of glutamine synthetase showed the importance of its nitrogen contribution to the plant. The glutamine synthetase is one of the main ammonium-assimilating enzymes; mutations in this enzyme generally result in the release of ammonium from the bacterium to the environment. The ammonium-excreting mutant performed better than the wild-type *A. brasilense* strain for wheat growth parameters and yield (Van Dommelen *et al.*, 2009).

An innovative approach to enhance nitrogen fixation to plants by *Azospirillum* was the creation of a specialized site for nitrogen fixation, a para-nodule. This root structure externally resembles a legume nodule and can be induced by adding low concentrations of the auxin herbicide 2,4-D to roots (Tchan *et al.*, 1991). Because *Azospirillum* does not secrete significant amounts of ammonium and sometimes provides the plant only small amounts of nitrogen, spontaneous mutants of *A. brasilense* were selected that excrete substantial amounts of NH_4^+ and the bacteria were established inside para-nodules. When plants were grown on a nitrogen-free medium, these mutants were responsible for significant increases in organic matter (root and shoot dry matter and total plant nitrogen), compared with plants treated with wild-type *Azospirillum* or plants that were not inoculated. Analysis of $^{15}\text{N}_2$ in these plants showed that the mutants were able to transfer more nitrogen to the host plants than the wild-type strain (Christiansen-Weniger and van Veen, 1991). Para-nodules induced in rice seedlings were the preferential sites for colonization by a NH_4^+ -excreting *A. brasilense* mutant. Nitrogenase activity in para-nodules structures inhabited by bacteria significantly increased, compared with untreated control plants (Christiansen-Weniger, 1997). It is probable that within para-nodules, bacterial nitrogenase is less sensitive to increased oxygen tension in the roots, as confirmed by Deaker and Kennedy (2001). Host plants benefit from enhanced nitrogen fixation in their roots with para-nodules because fixed nitrogen is incorporated into the host plant. Host plants probably stimulate nitrogenase activity of endophytic *Azospirillum* spp. by providing a carbon source as energy (Christiansen-Weniger, 1998).

These results show that the Gramineae are capable of establishing an association with diazotrophic bacteria in which ammonium-excreting bacteria provide the host plants with nitrogen. Para-nodule on wheat seedling roots was further developed by the researchers who invented it (Katupitiya *et al.*, 1995b; Sriskandarajah *et al.*, 1993) and specifically and consistently showed that nitrogenase activity in para-nodules was higher than in inoculated roots without para-nodules (Tchan *et al.*, 1991; Yu *et al.*, 1993; Zeman *et al.*, 1992). Similar results were obtained with maize (Saikia *et al.*, 2004, 2007). Para-nodules add a new dimension to research on biological nitrogen fixation, even if extensive developmental and biochemical modification of the para-nodule system is required before effective nitrogen fixation can be achieved. The options are intriguing (Christiansen-Weniger, 1994; Kennedy, 1994; Kennedy and Tchan, 1992).

In the last decade, perhaps as a response to the controversy mentioned earlier, studies have focused on the nitrogen cycle within cells of bacteria and on many details of molecular mechanisms and the genes involved that proliferate as *Azospirillum* was developed as a general model to study nitrogen fixation in nonsymbiotic bacteria. This is not a topic of this review (see e.g., Araujo *et al.*, 2004a,b; Huergo *et al.*, 2006a,b, 2009; Klassen *et al.*, 2005). The full genetic sequences of *A. brasilense* and *A. lipoferum* have been accomplished and they will be accessible at the Genoscope sites (France) and of *A. brasilense* at the Oak Ridge National Laboratory site (USA) (I. Kennedy, personal communication). Meanwhile, the complete nucleotide sequence of the *A. brasilense* *fixA*, *fixB*, *fixC*, and *fixX* genes were reported, as well as several other genes (Sperotto *et al.*, 2004). Mutants of the common *A. brasilense* strains Sp7 and Sp245 (defective in flocculation, differentiation into cyst-like forms, and colonizing of roots) had a higher nitrogenase expression than wild strains in association with wheat. Apparently, the ability of Sp7 and Sp245 mutants to remain in vegetative forms (spirillum and rods) improved their ability to express exceptionally high rates of nitrogenase activity. Restoring cyst formation and a normal colonizing pattern to the spontaneous mutant Sp7S reduced nitrogenase activity to the level of the wild Sp7. This suggests that bacterial cells in the vegetative state provides faster metabolism, which directly affects nitrogen fixation (Pereg-Gerk *et al.*, 2000). *A. brasilense* carrying *gfp* genes expressed pleiotropic physiological effects caused by disruption of the *clpX* gene encoding for heat-shock protein. One of the consequences of inserting the *gfp* gene is a threefold increase in nitrogen fixation (Rodriguez *et al.*, 2006). This phenomenon was confirmed in other *A. brasilense* strains (de Campos *et al.*, 2006). Apparently, higher expression of the *clpX* gene may be involved with creation of the Nif⁻ phenotype of the *A. brasilense* mutants by unknown mechanisms (Castellen *et al.*, 2009). Efficiency of nitrogen fixation and denitrification in *A. lipoferum* can be regulated by varying the concentration

of oxygen, nitrate, and molybdenum. The maximum growth rate in two strains was observed under microaerobic conditions, minimal nitrate, and the maximum concentration of molybdenum. These conditions were also conducive for obtaining maximum efficiency of denitrification (nitrate reduction to molecular N₂; Furina *et al.*, 1999). Microaerobic conditions favor nitrogen fixation. Low dissolved oxygen was also a limiting factor when ammonium concentrations limit growth of *A. lipoferum* (Tsagou *et al.*, 2003). In *A. brasilense*, cytochrome *c* oxidase is required under microaerobic conditions when a high respiration rate is needed. However, under nitrogen-fixing conditions, respiration rates do not seem to be a growth-limiting factor. Evidence for this was provided when a wild-type *A. brasilense* was compared with a *cytN* mutant *A. brasilense*. Under aerobic conditions, growth during the log phase was similar between the two types. Under microaerobic conditions (with NH₄⁺ supplied; no nitrogen fixation), low respiration of *A. brasilense cytN* decreased its growth rate compared with the growth rate of the wild-type *A. brasilense*. Under nitrogen-fixing conditions (without NH₄⁺ supplied), growth and respiration rates of the wild-type bacterium were significantly diminished and the differences in growth and respiration rates between the wild and mutant forms were smaller. Yet, the nitrogen-fixing capacity of the mutant was still approximately 80% of the wild-type (Marchal *et al.*, 1998). Out of 40 thermo-tolerant mutants developed from a mesophilic *A. lipoferum*, only 14 could grow and fix nitrogen at 45 °C. These mutants excrete ammonia only as very old cultures (maximum production after 12 days under stationary conditions; Steenhoudt *et al.*, 2001). Nitrogen fixation by aerobic bacteria is a very energy demanding process, requiring efficient oxidative phosphorylation, since O₂ is toxic to the nitrogenase complex. *Azospirillum* spp. and other well-known nitrogen-fixing soil bacteria have evolved a variety of strategies to deal with and overcome the apparent “O₂ paradox.” The question is whether the specific environmental adaptations of azospirilla are sufficient to allow optimal proliferation and nitrogen fixation in their natural habitat. Could improving O₂-tolerance of the nitrogen-fixing process contribute to the development of more efficient strains for inoculation of plants (Marchal and Vanderleyden, 2000)? This remains a future research objective.

In evaluating the overwhelming data accumulated over the last 35 years on nitrogen fixation by *Azospirillum*, ignoring nitrogen fixation as a mechanism for *Azospirillum*, is premature. In several systems of inoculation, clear demonstration of significant increases of fixed nitrogen for plant growth was demonstrated, while it did not occur in others tests. It is also feasible that in systems where the contribution is small, the quantity of nitrogen provided by the nitrogen-fixing process is accumulative, with other mechanisms to produce the final growth promotion effect (see Section 3.6).

2.3. General improvement of root growth and enhanced uptake of minerals and water

Enhanced root systems, including root hairs, are the most common phenotypic phenomena observed following *Azospirillum* inoculation in most species. Consequently, improved root growth and function leading to improved water and mineral uptake was proposed in the late 1970s. Enhanced mineral uptake was a popular explanation for the inoculation effects in the 1980–1990s (for reviews, see [Bashan and Holguin, 1997](#); [Bashan and Levanony, 1990](#)).

Increased mineral uptake in plants has been suggested due to a general increase in volume of the root system and not to any specific metabolic enhancement of the normal ion uptake mechanism ([Morgenstern and Okon, 1987](#); [Murty and Ladha, 1988](#)) and that this is related to secretion of phytohormones by the bacteria. Other studies suggested a more active involvement in acquisition of minerals. Inoculation may promote availability of ions in the soil by helping the plant scavenge limiting nutrients ([Lin et al., 1983](#)), which may explain the common accumulation of N compounds in the plant without any apparent N₂ fixation. Thus, the plant may absorb N more efficiently from the limited supply in the soil, resulting in a less N fertilization to attain a desired yield. By volume of information this is one of the largest parts of the *Azospirillum* literature, although the physiological, biochemical, and molecular details were left unsearched and only analyses of specific variables was presented. As a result, the available information about this mechanism is largely descriptive.

Several examples, out of many, illustrate the mechanism. In hydroponic systems in greenhouses, inoculation with *A. brasilense* increased the number and length of adventitious roots of *Sorghum bicolor* by 33–40% over non-inoculated controls, such as a higher rate of growth, earlier root appearance, and a greater elongation rate of individual roots ([Sarig et al., 1992](#)). In addition to increasing ([Kapulnik et al., 1981, 1985b](#)) or decreasing ([Kucey, 1988](#)) many root parameters, inoculation affected many foliage parameters. These changes were directly attributed to positive bacterial effects on mineral uptake by the plant. Enhancement in uptake of NO₃⁻, NH₄⁺, PO₄²⁻, K⁺, Rb⁺, and Fe²⁺ and several micronutrients by *Azospirillum* ([Barton et al., 1986](#); [Jain and Patriquin, 1984](#); [Kapulnik et al., 1985a](#); [Lin et al., 1983](#); [Morgenstern and Okon, 1987](#); [Murty and Ladha, 1988](#); [Sarig et al., 1988](#)) was proposed to cause an increase in foliar dry matter and accumulation of minerals in stems and leaves. During the reproductive period, these minerals could have been transferred to the panicles and spikes and result in higher yield and higher mineral content ([Ogut and Er, 2006](#)). Supporting evidence for increased mineral uptake by inoculated roots is provided by enhancement in proton efflux activity of wheat roots inoculated with *Azospirillum* ([Bashan, 1990](#); [Bashan et al., 1989a](#)). It is well known

that proton efflux activity is directly related to the balance of ions in plant roots (described below). Although some studies showed accumulation of N and minerals in the inoculated plants, others showed that enhanced growth of wheat and soybeans was not necessarily because of a general enhancement of mineral uptake (Bashan *et al.*, 1990).

In addition to improved mineral uptake, inoculation improved water status in stressed sorghum plants. Inoculated plants were less stressed, having more water in their foliage, higher leaf water potential, and lower canopy temperature than noninoculated plants. Soil moisture extraction by *Azospirillum*-inoculated plants was greater and water was extracted from deeper layers in the soil. Therefore, increased sorghum yield was primarily attributed to improved utilization of soil moisture (Sarig *et al.*, 1988; see Section 2.5 for more details.)

It is likely that improved mineral and water uptake occur in the *Azospirillum*-plant association. However, the descriptive data presented so far have not shown whether these improvements are the cause or the result of other mechanisms, such as changes in the balance of plant hormones or enhanced proton extrusion. Furthermore, the wide range of enzymatic activities related to these phenomena were poorly studied and no apparent evaluation of *Azospirillum* mutants deficient in induction of mineral and water uptake by plants has been made. Finally, it should be noted that very few strains have been studied and it is doubtful if all *Azospirillum* strains possess these abilities. There is evidence that some strains of *A. brasilense* failed to improve uptake of several ions, but nevertheless improved plant growth (Bashan *et al.*, 1990).

2.4. Phosphate solubilization and mobilization and rock weathering

Despite the reservations listed above, improved mineral uptake by plants was suggested as a major contribution of *Azospirillum* inoculation, therefore, azospirilla weathering of minerals in general and phosphorus in particular were studied. This has received attention because of the related larger field of phosphate solubilization that involves other bacterial genera.

A. halopraeferens, a bacterium that does not use glucose, and consequently does not produce acid, can solubilize insoluble inorganic phosphate *in vitro* by unknown mechanisms (Seshadri *et al.*, 2000). Two strains of *A. brasilense* and one strain of *A. lipoferum* were capable of producing gluconic acid, thereby leading to solubilization of insoluble phosphate in rocks (Puente *et al.*, 2004a; Rodriguez *et al.*, 2004). Sugars, like glucose, are part of the root exudates of pea plants grown in P-deficient substrates and enhanced the capacity of *Azospirillum* spp. to solubilize normally insoluble $\text{Ca}_3(\text{PO}_4)_2$. The relative proportion of glucose in pea exudates decreased under P deficiency, while the content of galactose, ribose, xylose, and fucose increased. *Azospirillum* spp. can metabolize all these sugars. Therefore, the shift in sugars under

P deficiency increased the capability of *Azospirillum* spp. to mobilize phosphate (Deubel *et al.*, 2000). Similarly, inoculation of cardon (a giant cactus) with *A. brasilense* Cd enhanced phosphate solubilization and enhanced plant growth (Carrillo *et al.*, 2002).

These observations can partly be explained by acidification of the nutrient medium by protons and organic acids. *Azospirillum* spp. can produce different organic acids that assist in P solubilization, depending on the sugar in the root exudates. Yet, *Azospirillum* can solubilize P by itself without adding root exudates. For example, three *Azospirillum* strains were isolated from the ectomycorrhizal sporocarps (*Rhizopogon vinicolor*) that colonized Douglas fir trees. *In vitro*, they were able to degrade limestone, marble, and calcium phosphate (Chang and Li, 1998). These observations were confirmed using other strains of *Azospirillum* (Puente *et al.*, 2004a,b, 2006). Uptake by *A. brasilense* cells of the essential elements Mg, Ca, Mn, and Fe and trace elements V, Co, Ni, Cu, Zn, and Pb (which do not essentially suppress growth of bacterial cultures) present in weathered rock fragments and are accumulated by the cells was shown. Zn and Cu were accumulated in the bacterial biomass in relatively significant amounts, but uptake of Co and Ni was much less, and Pb and V were apparently not assimilated by azospirilla. In particular, Cu cations were effectively absorbed by the bacterium and this increased the rate of uptake of other metals; however, the process takes time. Short exposures have only a limited effect on absorption of Cu (Ignatov *et al.*, 2001; Kamnev *et al.*, 1997a). Additionally, these bacteria are capable of producing structural modifications of the magnesium-ammonium orthophosphate molecule when added to the medium (Kamnev *et al.*, 1999a). Fourier transform infrared spectroscopy is a powerful tool for nondestructive identification and characterization of cell components; it was applied to studies of molecular structures in *A. brasilense*, its essential element content (Kamnev *et al.*, 1997b, 2001), heavy metal-induced metabolic changes in the cells (Kamnev *et al.*, 2002), and membrane composition and structure (Kamnev *et al.*, 1999b). These capabilities notwithstanding, it has not been demonstrated so far that these elements, obtained from the environment, were transferred to the plant.

The research field of mineral solubilization and mobilization in *Azospirillum* is potentially useful for studying interactions and survival of the bacteria in the soil. Although the literature treated this proposal as an individual entity, it should be considered as a subfield of enhanced mineral uptake mentioned earlier.

2.5. Mitigation of stresses

From the earliest field experiments with *Azospirillum* in the 1980s, the best effects on plant growth and yield were obtained when the growth conditions were suboptimal. A common explanation for the effects of *Azospirillum* on

plant growth was reduction in environmental stresses by the bacteria, providing the plant a more favorable environment to grow in an otherwise limiting environment. Sometimes inoculation permits plant growth in soils that normally did not allow growth. None of these theories can explain enhanced growth of inoculated plants under favorable plant growth conditions that also occurred and were regularly reported. Environmental stressors varied and included mitigation of drought (Sarig *et al.*, 1990), salinity (Creus *et al.*, 1997), heavy metals (Belimov and Dietz, 2000), toxicity of other substances (de-Bashan and Bashan, 2008), extreme pH (de-Bashan *et al.*, 2005), toxic humic substances (Bacilio *et al.*, 2003), and suboptimal levels of nitrogen (discussed earlier).

2.5.1. Salinity stress

Numerous cultivated soils worldwide are becoming more saline, mainly from the use of marginal irrigation water, from excess fertilization, and various desertification processes. Inoculation with *Azospirillum* sp. under saline stress conditions is therefore commonplace. Prior findings (for a review, see Bashan and Holguin, 1997) showed that common agricultural *Azospirillum* strains tolerated high salinity (~2%). Salt resistance among species increased from *A. amazonense* (lowest) to *A. halopraeferans* (highest), the latter tolerating over 3% NaCl (seawater salinity).

Azospirillum inoculation of maize at NaCl concentrations up to -1.2 MPa significantly increased chlorophyll, K, Ca, soluble saccharides, and protein contents, compared with control maize growing without NaCl (Hamdia and El-Komy, 1997). Alleviation of salt stress in maize involved several changes that probably were related to different operating mechanisms: proline concentration declined significantly, the concentration of most amino acids increased on exposure to NaCl, as well as when inoculated with *Azospirillum*. *Azospirillum* apparently restricted Na⁺ uptake and enhanced the uptake of K⁺ and Ca²⁺. Finally, inoculation stimulated nitrate reductase and nitrogenase activity in shoots and roots (Hamdia *et al.*, 2004). Inoculating wheat seedlings with *A. brasilense* exposed to severe salt (NaCl) or osmotic (polyethylene glycol) stress significantly reversed part of the negative effects; both stresses reduced the relative elongation rate of shoots. Fresh weight, fresh weight/dry weight ratio, water content, and relative water content were higher in shoots from inoculated plants than in stressed controls (Creus *et al.*, 1997). Similarly, under high NaCl concentration, inoculation of wheat with *A. lipoferum* reduced some of the deleterious effects of NaCl (Bacilio *et al.*, 2004). Finally, *Azospirillum*-inoculated lettuce seeds had better germination and vegetative growth than noninoculated controls after being exposed to NaCl (Barassi *et al.*, 2006).

The most fundamental omissions in current knowledge are (1) uncertainty about whether improved salt tolerance of the bacterium is needed to enhance the bacterium's effect on plants or if existing salt tolerance in plants

is adequate to ensure positive growth-promotion by inoculation; (2) what are the mechanisms that are triggered and are responsible for enhanced saline resistance after inoculation; and (3) what is the microbial mechanism that provides resistance in plants.

2.5.2. Water stress

Apparently, inoculation with *Azospirillum* improved growth under water-stress conditions as was initially demonstrated in the 1980s (for a review, see [Bashan and Levanony, 1990](#)). Subjecting inoculated *S. bicolor* plants to osmotic stress in hydroponic systems diminished the adverse effects caused by osmotic stress, such as reduction of leaf senescence ([Sarig et al., 1990](#)). Coleoptile height and fresh and dry weight of wheat seedlings inoculated with *A. brasilense* Sp245 were enhanced, despite the water stress ([Alvarez et al., 1996](#)). Inoculation with *Azospirillum* alleviated the stress on wheat plants grown under drought conditions ([El-Komy et al., 2003](#)). Turgor pressure at low water potential was higher in inoculated seedlings in two wheat cultivars under osmotic stress. This could result from better water uptake as a response to inoculation that, in turn, is reflected by faster shoot growth in inoculated seedlings exposed to these stresses. They showed better water status and effects on cell wall elasticity or apoplastic water ([Creus et al., 1998](#)). To assess the contribution of *A. brasilense* Sp245 during drought when flowers open (anthesis), inoculated wheat seeds were subjected to drought. Even though all the plants underwent osmotic stress, significantly higher water content, relative water content, water potential, apoplastic water fraction, and lower cell wall modulus of elasticity values were obtained in inoculated plants. Grain yield loss to drought in inoculated plants was significantly reduced and significantly higher Mg, K, and Ca in grains were detected. Probably, inoculation improved water status and an additional “elastic adjustment” in plants ([Creus et al., 2004](#)). Recently, inoculation with *A. brasilense* contributed to protection of wheat seedlings under water stress through changes in the fatty acid distribution profiles of phosphatidylcholine and phosphatidylethanolamine, major root phospholipids ([Pereyra et al., 2006](#)). Transformed *A. brasilense* that could produce trehalose, an osmotic-regulating sugar, was more salt resistant than the wild type and significantly enhanced the survival of maize growing under drought stress. It also significantly increased biomass and leaf and root length of the plants ([Rodríguez-Salazar et al., 2009](#)). The limitations in our knowledge regarding the effect of inoculation under saline stress are valid for osmotic stress, as well.

2.5.3. Herbicides

Cotton plants could be partly protected from harmful effects of the herbicide 2,4-D by inoculation with *A. brasilense*. The degrading plasmid of 2,4-D was transferred into *A. brasilense* Sp7. Trans-conjugants degraded

2,4-D in pure culture via cometabolism. However, when the trans-conjugants were inoculated on cotton seeds, the plants were resistant only to low levels of the herbicide, which is not sufficient for protection of cotton. Plants growing in soils with this concentration of herbicide and inoculated with wild-type strains died (Feng and Kennedy, 1997).

2.5.4. Toxic metals

Another possible mechanism for producing a healthier plant is reduction of metal toxicity in contaminated soils and mine tailings that, under normal conditions, almost completely inhibits plant growth. Although the bacterium tolerate only moderate levels of metals and other toxic compounds (see previous reviews Bashan and Holguin, 1997; Bashan and Levanony, 1990; Bashan *et al.*, 2004; also Kamnev *et al.*, 2005, 2007), it apparently contributed mechanisms allowing plants to grow in mine tailings or contaminated soils. Cadmium causes severe inhibition of growth and nutrient uptake in barley. In the presence of CdCl_2 , inoculation with *A. lipoferum* partly decreased Cd toxicity, possibly through the improvement of mineral uptake. Additionally, inoculation slightly enhanced root length and biomass of barley seedling treated with Cd and the amount of nutrients absorbed by the inoculated plants increased significantly. There was only some protection against Cd toxicity, but no uptake of Cd, since Cd content in the inoculated plants was unchanged (Belimov and Dietz, 2000; Belimov *et al.*, 2004). *A. brasilense* Sp245 associated with wheat changes the speciation, bioavailability, and plant uptake of arsenic. Plants inoculated with *Azospirillum* accumulated less arsenic than did uninoculated plants (Lyubun *et al.*, 2006). Inoculation of the wild desert shrub quailbush (*Atriplex lentiformis*) growing in extremely stressed environment with *A. brasilense* strains Sp6 and Cd, such as acidic mine tailings having high metal content, resulted in a significant increase in production of plant biomass (L.E. de-Bashan *et al.*, unpublished data). Similar results were obtained when wild yellow palo verde desert trees (*Parkinsonia microphylla*) were inoculated with *A. brasilense* Cd in rock phosphate tailings (Bashan *et al.*, unpublished data).

2.5.5. Compost and humic substances

Some compost may be toxic to plants because of elevated humic acids or inappropriate preparation. Inoculation of wheat seeds with *A. brasilense* or *A. lipoferum* prior to sowing in soil that was amended with two types of compost improved seed germination and plant development. The bacteria possibly changed or consumed the humic acids because both bacterial species can survive and grow in high humic acid solution as the sole source of carbon; thus, modify the composition of the compost during *in vitro* tests (Bacilio *et al.*, 2003).

2.5.6. pH and toxic substances in aquatic environments

Apart from terrestrial applications, *Azospirillum* is being used as an inoculant in aquatic environments mainly to promote the growth and metabolism of microalgae of the genus *Chlorella* that is used in wastewater treatment (de-Bashan *et al.*, 2004; Gonzalez and Bashan, 2000; Hernandez *et al.*, 2006). Under aquatic conditions, the pH, available dissolved nutrients, and toxic molecules to the microalgae have significant impact on the process of mass production. High pH of the medium interferes with the microalgal cell cycle and decreases microalgal population. Coculturing of the microalgae with *A. brasilense* eliminated this negative effect (de-Bashan *et al.*, 2005). Similarly, high levels of the amino acid tryptophan reduced multiplication of *C. vulgaris* where coculturing with *A. brasilense* significantly reduced the inhibition probably by converting it to IAA that enhances the growth of the microalgae (de-Bashan and Bashan, 2008).

2.5.7. Protection from relative high light intensities

Inoculation of plants sometimes occurs under light intensity that is stressful and has an inhibiting effect on specific crops. Inoculation of wheat seedlings with *A. brasilense* Cd significantly increased the quantity of the photosynthetic pigments chlorophyll *a* and *b*, but also the auxiliary photoprotective pigments violaxanthin, zeaxanthin, antheroxanthin, lutein, neoxanthin, and β -carotene that help the plant to sustain photosynthesis under unfavorable light conditions. This outcome yielded greener plants with no apparent visible stress. The greatest difference in the quantity of all pigments between inoculated and noninoculated plants occurred in the first week of growth (Bashan *et al.*, 2006). Similarly, although the microalgae *C. sorokiniana* is capable of growing at high light intensities, coculturing with *A. brasilense* enhanced this capacity and the microalgae could tolerate extreme light intensities as high as $2500 \mu\text{mol m}^{-2} \text{s}^{-1}$ (de-Bashan *et al.*, 2008b).

Taking all these phenomena together, it appears that a multitude and remotely related or unrelated mechanisms are operating in these complex interactions of *Azospirillum* with plants. All these accumulating findings yielded a recent proposal to include *Azospirillum* in the group of other rhizosphere PGPB that regulated homeostasis of plants under conditions of abiotic stress. This group was designated "Plant Stress-Homeo-regulating Bacteria" (PSHB; Cassan *et al.*, 2009b; Sgroj *et al.*, 2009). These types of bacteria, *Azospirillum* included, may use an assortment of mechanisms, such as biosynthesis of phytohormones, growth regulators, osmoregulator molecules, expression of specific regulatory and metabolic enzymes, and immobilization or catabolism of various toxic molecules for plants to assist plant growth. This proposal forms a part of the initial theory of Section 3.6.

3. OTHER PROPOSED MECHANISMS

3.1. Biological control

Azospirillum is not yet known as a typical biocontrol agent of soil-borne plant pathogens because many strains lack direct suppressive chemicals or hydrolytic enzymes likely to affect plant pathogens. However, reports are accumulating that this mechanism has been overlooked. Some possible mechanisms used by *Azospirillum* to reduce damage from pathogens have been demonstrated as environmental competition and displacement of pathogens, inhibition of seed germination of parasitic weeds, general enhancement of plants to resist pathogen infection, and possible inhibition of fungal growth via production (at least *in vitro*) of microbial toxic substances.

3.1.1. Toxic substances

When iron was withheld, *A. lipoferum* strain M produced catechol-type siderophores under iron starvation that exhibited antimicrobial activity against various bacterial and fungal isolates (Shah *et al.*, 1992). Although some strains from Brazil produce cyanide (HCN) *in vitro* (Gonçalves and de Oliveira, 1998), this feature is uncommon in strains from other geographic locations. Some *Azospirillum* isolates produced bacteriocins that inhibited growth of several indicator bacteria (Tapia-Hernandez *et al.*, 1990). An antimicrobial auxin-like molecule, phenylacetic acid was isolated from an *A. brasilense* culture (Somers *et al.*, 2005). *A. brasilense* cells contain a low molecular-weight compound that inhibits germination and growth of the radicle of Egyptian broomrape seeds (*Orobanche aegyptiaca*), a specific weed parasite of sunflower (Dadon *et al.*, 2004). *Azospirillum* spp. inhibited germination of the parasitic striga weed (witchweed) seeds (*Striga hermonthica*) that infest fields of tropical sorghum, thereby promoting growth of sorghum (Bouillant *et al.*, 1997). *Azospirillum* cells suspended in a synthetic germination stimulant did not inhibit germination of striga weed seeds, but blocked radicle elongation. These radicles had abnormal morphology and contained no vacuolated cells in the root elongation zone. Lipophilic compounds extracted from the medium of bacteria prevented germination of striga seeds (Miché *et al.*, 2000). So far, *Azospirillum* has not been reported to induce any negative effect on healthy plants (Bashan, 1998). If this is the case, these toxic compounds are either *in vitro* artifacts or are induced only in the presence of pathogens.

3.1.2. Competition

The effect of *A. brasilense* on crown gall formation in Dicotyledoneae was studied after inoculation with virulent strains of *Agrobacterium tumefaciens*. When wounded tissues of grapevines and carrot disks were inoculated with live cells of *A. brasilense* strains 94-3 or Sp7, development of the typical

bacterial galls was inhibited and the protective effect of *Azospirillum* lasted over a 24-h period (Bakanchikova *et al.*, 1993). When *A. brasilense* Cd was added to a culture with the pathogenic mangrove rhizosphere bacterium *Staphylococcus* spp., the population of the latter was significantly reduced (Holguin and Bashan, 1996).

To assess displacement of pathogens by inoculation with *Azospirillum*, the tomato leaf pathogen *Pseudomonas syringae* pv. *tomato* (PST, bacterial speck) and *A. brasilense* were inoculated onto tomato plants, as a mixed culture or consecutively. Inoculation of seeds with a mixed culture resulted in reduction of the pathogenic population in the rhizosphere, increased the population of *A. brasilense*, prevented development of PST, and improved plant growth. PST did not survive in the rhizosphere in the presence of *A. brasilense*. Inoculation of leaves with the mixed bacterial culture under mist conditions significantly reduced the population of PST and significantly decreased the severity of the disease. Challenge with PST after *A. brasilense* was established in the leaves further reduced PST and severity of the disease and significantly enhanced plant development. Selective enhancement of the population of *A. brasilense* on leaves occurred by applying malic acid (favorable for *A. brasilense*, but not for PST), decreased PST to almost undetectable levels, almost eliminated disease development, and improved plant growth to the level of uninoculated healthy controls (Bashan and de-Bashan, 2002a). Seeds inoculated with *A. brasilense* Sp7 and later challenged by two foliar bacterial pathogens of tomato (*Clavibacter michiganensis* spp. *michiganensis* [bacterial canker] and *Xanthomonas campestris* pv. *vesicatoria* [XCV, bacterial spot]) delayed leaf- and plant-death compared with untreated controls, but canker severity was not affected. Unfortunately, inoculation with *Azospirillum* increased the severity of XCV on cherry tomatoes (Romero *et al.*, 2003). Several isolated bacterial strains showed antagonism toward the fungus *Aspergillus flavus* that produces aflatoxin (the most potent carcinogenic mycotoxin produced by some fungi), and were capable of degrading the toxin *in vitro*. Since identification of the microorganism was based on morphological characteristics, it is uncertain whether the identification of the strains as *Azospirillum* is valid (Cho *et al.*, 2000). A strain of *A. brasilense* with increased capacity for N₂-fixation was tested *in vitro* against the soil-borne plant pathogens, *Fusarium oxysporum* f. sp. *lycopersici*, *Rhizoctonia solani*, and *Pythium* sp. that infect cucumbers. The bacteria reduced the dry weight of *Fusarium* mycelium by 90–96%, of *Rhizoctonia* by 72–94%, of *Pythium* by 71–95%, and completely eliminated *Sclerotinia* mycelium (Hassouna *et al.*, 1998).

3.1.3. Production of a “healthier plant” by unknown mechanisms

Many examples of possible “biological control” are reported without specifying the mechanisms. It is assumed that inoculation produce healthier plants by deterring pathogenic infections (Tilak *et al.*, 2005). This is a

possibility especially when the effect recorded is against microfauna and insects and not against microorganisms, as is expected.

For example, inoculation with *A. lipoferum* of mung bean (*Vigna radiata*) infected with root-knot nematode (*Meloidogyne incognita*) led to fewer root galls and egg masses per root system. After inoculation with *A. lipoferum*, plants infected with the nematode had significantly greater growth and biomass, probably related to a greater number of functional nodules on roots that had been infected (Khan and Kounsar, 2000). When inoculated with *A. brasilense*, okra (*Abelmoschus esculentus*) showed enhanced plant characteristics and pod yield. At the same time, there was a significant reduction in root-knot nematodes egg masses, eggs per egg mass, and total nematode population (Ramakrishnan *et al.*, 1997). Similar results in sunflower were obtained with a commercial inoculant of *A. brasilense* (Ismail and Hasabo, 2000). Maize that was inoculated with a combination of mycorrhizal fungi, *Glomus fasciculatum*, *Azospirillum* sp., and phosphate-solubilizing bacteria reduced the population of the *Pratylenchus zeae* nematode and induced very high cob yield (Babu *et al.*, 1998). When *A. lipoferum* was inoculated onto wheat plants, it reduced *Heterodera avenae* nematode infection (Bansal *et al.*, 1999). Inoculation of sorghum with *A. brasilense* to control the sorghum shoot fly *Atherigona soccata* that causes dead-heart in sorghum resulted in a 10-fold reduction of the disease and increased grain yield (Kishore, 1998).

A. brasilense was applied as a foliar spray against foliar fungal and bacterial diseases of mulberry, such as powdery mildew caused by *Phyllactinia corylea*, black leaf spot caused by *Pseudocercospora mori*, black leaf rust caused by *Cerotelium fici*, and bacterial leaf blight caused by *P. mori*. Inoculation reduced fungal pathogens and excelled as a treatment against bacterial blight (Sudhakar *et al.*, 2000). The addition of *Rhizobium*, *Azospirillum*, or *Azotobacter* inocula as a combined seed and soil treatment in cultivation of pearl millet (*Pennisetum glaucum*) reduced downy mildew (*Sclerospora graminicola*) in the leaves (Gupta and Singh, 1999). Inoculation with arbuscular mycorrhizal (AM) fungi and *Azospirillum* spp. suppressed damping-off disease in chili (*Capsicum* sp.) caused by *Pythium aphanidermatum* (Kavitha *et al.*, 2003). Combinations of several ineffective management tactics (spraying Cu and streptomycin combined with *Azospirillum* seed inoculation and seed disinfections, individually ineffective against PST, significantly reduced occurrence and severity caused by PST and also improved plant growth. Additionally, the combined treatment significantly reduced the amount of chemical pesticides required to protect tomato plants from PST (Bashan and de-Bashan, 2002b). The mechanisms by which this happens in all the described cases remain unknown.

So far, *Azospirillum* is not commonly reported to induce systemic resistance in plants. However, inoculation of rice plant with the endophyte *Azospirillum* sp. B510 induced disease resistance against diseases caused by

the virulent rice blast fungus *Magnaporthe oryzae* and bacterial pathogen *Xanthomonas oryzae*, apparently by activating a novel type of resistance mechanism independent of salicylic acid-mediated defense that does not signal accumulation or expression of pathogenesis-related genes (Yasuda *et al.*, 2009).

At this time, these reports do not provide conclusive evidence that *Azospirillum* is a true biological control agent, although significant biological control activity can be attributed to this genus.

3.2. Nitric oxide

Nitric oxide (NO) is a volatile, lipophilic free radical which participates in metabolic, signaling, defense, and developmental pathways in plants (Cohen *et al.*, 2010; Lamattina and Polacco, 2007; Lamattina *et al.*, 2003). As its major role, NO participates in the IAA signaling pathways. This participation leads to lateral and adventitious root formation where the exact role of NO is as an intermediary in IAA-induced root development (Correa-Aragunde *et al.*, 2004, 2006; Pagnussat *et al.*, 2002, 2003).

One wild-type *A. brasilense* Sp245 can produce NO *in vitro*, under anoxic and oxic (or aerobic) conditions (Creus *et al.*, 2005). The latter can be achieved by possible different pathways, such as aerobic denitrification and heterotrophic nitrification. NO is produced during the middle and late logarithmic phases of growth (Molina-Favero *et al.*, 2007, 2008). An NO-dependent promoting activity in *A. brasilense* Sp245 induces morphological changes in tomato roots regardless of the full bacterial capacity for IAA synthesis. An IAA-attenuated mutant of this strain, producing up to 10% of the IAA level compared with the wild-type strain (Dobbelaere *et al.*, 1999) had the same physiological characteristics and slightly less effect on root development. When the NO was removed, using a chemical NO scavenger, both types of root formation were inhibited. This demonstrates that NO-mediated *Azospirillum* induced branching of roots. These results provide further evidence of an NO-dependent promoting activity of tomato root branching, regardless of the bacterium's capacity for synthesizing IAA (Molina-Favero *et al.*, 2008), a phenomenon that occurs in other inoculation systems lacking IAA activity (see above). It is commonly argued that denitrification in agriculture is considered, in general, and specifically in plant inoculations, as an undesirable feature of PGPB because it reduces availability of N (Zimmer *et al.*, 1984) for the plant. Yet, the capacity of *A. brasilense* to reduce nitrate aerobically to NO, which in turn, could promote growth of tomato roots is a point for reconsideration.

Several studies demonstrate the continuous relation between NO and IAA on root development (Huang *et al.*, 2007; Lombardo *et al.*, 2006; Tewari *et al.*, 2007). It is possible that a connection, not proven so far, exists in *Azospirillum*-plant systems. However, the way that IAA and NO

are acting together, if acting, on plant cells triggering the branching of roots is still an open question for research. Nonetheless, the relationship between NO and *A. brasilense* showed that, in addition to the well-established connection between NO production and defense responses to pathogenic microorganisms (Modolo *et al.*, 2005; Zeidler *et al.*, 2004), it seems that NO metabolism also plays a role in the positive close association of PGPB with roots.

3.3. Nitrite

Nitrite (NO_2^-), either directly added or excreted by *A. brasilense* during nitrate respiration may participate in growth promotion effects. It causes a sharp increase in the formation of lateral roots (Zimmer *et al.*, 1988). Nitrite is produced under anaerobic or microaerobic conditions by the dissimilatory nitrate reduction pathway, in addition to NO and nitrous oxide (N_2O ; Hartmann and Zimmer, 1994). Nitrite could have promoting effects when reacting with ascorbate (Bothe *et al.*, 1992; Zimmer *et al.*, 1988). This avenue has not been investigated further.

3.4. Signal molecules and enhanced proton extrusion from roots

Whatever the operating mechanism, *Azospirillum* affects plant cell metabolism from outside the cell (without entering the intact plant cells) and this suggests that these bacteria are capable of excreting and transmitting a signal (s) that crosses the plant cell wall and is recognized by the plant membranes. This interaction can initiate a chain of events resulting in altered metabolism of the inoculated plant and proliferation of roots. Since plant membranes are extremely sensitive to any change, their response may serve as a precise indicator of *Azospirillum* activity at the cellular level. Improving plant growth by affecting proton and organic acid extrusion (proton pump) mechanisms in plants by inoculation with *Azospirillum* spp. was proposed two decades ago.

A proton pump is an integral membrane protein that is capable of moving protons (H^+) across the membrane of a cell, mitochondrion, or other subcellular compartment. In cell respiration, the pumps move protons from the space enclosed by the two membranes within the organelle and release the protons into the intermembrane space. The confined protons create a gradient in both pH and electrical charge across the plasma membrane that acts as a reservoir of stored energy for the cell. For plants to react to their constantly changing environments and simultaneously maintain optimal metabolic conditions, the expression, activity, and interplay of the pumps generating these H^+ gradients have to be tightly regulated (Gaxiola *et al.*, 2007; Schumacher, 2006). Additional functions, such as opening and

closing of stomata, cell growth, and intracellular pH homeostasis, have been proposed (Duby and Boutry, 2009). Short exposure of wheat roots to live *A. brasilense* Cd significantly enhanced the proton efflux of the root at 5 h after inoculation. Bacteria in the logarithmic phase are required for this enhancement, which is a triggering nature (Bashan, 1990; Bashan *et al.*, 1989a). Inoculation of soybean seedlings with the same *Azospirillum* strain significantly reduced the membrane potential in every root part and this was greatest in the root elongation zone (Bashan, 1991; Bashan and Levanony, 1991). Inoculation of soybeans and cowpeas with this strain increased proton efflux from their roots and changed the phospholipid content in membranes of cowpeas (Bashan *et al.*, 1992). Although the nature of the released signal molecule is still unknown, *Azospirillum* probably targets plant membranes on plant roots. This phenomenon also occurs in cardon cactus. Lowering the pH of the rhizosphere increases the availability of phosphorus and iron to plants, especially in arid lands with high calcium content and soil pH (Carrillo *et al.*, 2002). A confirmatory study of the proton extrusion phenomenon in wheat showed that inoculation enhanced proton efflux and elongation of the roots. Although the evidence is circumstantial, perhaps these two phenomena are related. This effect was directly dependent on the bacterial strain–plant combination, suggesting that compatible strains are necessary to induce this activity (Amooaghaie *et al.*, 2002). This kind of investigation has not been pursued in recent years.

It is possible that a receptor in *A. brasilense* is involved in the binding of wheat germ agglutinin (WGA; one of the most studied plant lectins; Antonyuk *et al.*, 1993). This binding induced changes in the cellular metabolism of *A. brasilense* Sp245 and promoted nitrogen fixation, excretion of ammonium ions, and synthesis of IAA (Antonyuk and Evseeva, 2006; Antonyuk *et al.*, 1993, 1995). WGA changed the relative proportion of acidic phospholipids of the membrane. It is possible that acidic phospholipids participate in trans-membrane communication. WGA may function as a signal molecule in the *Azospirillum*–plant association (Antonyuk *et al.*, 1995). Some *Azospirillum* strains are known to produce several lectins *in vitro* (Castellanos *et al.*, 1998). Two cell-surface lectins isolated from *A. brasilense* Sp7 and from a mutant (defective in hem-agglutinating activity), *A. brasilense* Sp7.2.3, affected activities of α -glucosidase, β -glucosidase, and β -galactosidase in the membrane and apoplast fractions of roots of wheat seedlings (Alen'kina *et al.*, 2006). Other lectins induced changes in the mitotic state of growing onion plant cells (Nikitina *et al.*, 2004).

In general, effects on proton extrusion merit further investigation because the changes in the metabolism of the roots may induce enhanced mineral and water uptake even without proliferation of roots that are induced by phytohormones. This may provide further support to the theory of enhanced mineral uptake in cases when hormonal activity is not detected.

3.5. *Azospirillum* nitrate reductase

An alternative to nitrogen fixation as an explanation for N accumulation after inoculation of wheat plants by *Azospirillum* is the bacterial NR theory. NR activity in wheat leaves was decreased by inoculation with some *Azospirillum* strains. Inoculation with NR⁻ mutants resulted in a small response, concomitant with an increase in leaf NR, compared with inoculation with the parental NR⁺ strain (Ferreira *et al.*, 1987). Inoculation of field-grown plants with *A. brasilense* Sp245 and its NR⁻ mutant confirmed that the mutant was significantly less effective in increasing yield than the parental strain (Boddey and Döbereiner, 1988). This phenomenon indicates that the effect of some *Azospirillum* strains on wheat plants is not solely via nitrogen fixation (both the parental and the mutant strains have this ability), but rather results from an increase in assimilating nitrate. The parental strain aided reduction of nitrate in the roots and thus decreased translocation of nitrate to the leaves, while inoculation with the NR⁻ mutant caused direct translocation and reduction of nitrate in the foliage. This theory might explain, in part, the observation of increased N accumulation in shoots because the unaffected ability to fix nitrogen may also contribute N to the plants in addition to NR activity. It also might be a part of a larger theory of enhanced mineral uptake by *Azospirillum* inoculation (described earlier). This line of research has not been pursued further.

3.6. Additive hypothesis

Several recent studies on modes of action in *Azospirillum* gave new momentum to the additive hypothesis that was suggested 20 years ago. The hypothesis considers multiple mechanisms rather than one mechanism participating in the association of *Azospirillum* with plants. These mechanisms operate simultaneously or in succession, the contribution of an individual mechanism being less significant when evaluated separately. The sum of activities under appropriate environmental conditions results in the observed changes in plant growth (Bashan and Levanony, 1990). For example are the cases where nitrogen fixation contributes less than 5% of the observed effect of *Azospirillum* on the plant. As such low levels, it is not sufficient and does not fully explain increases in yield. When combined with other small mechanisms, this may be a significant contribution. With a general mechanism unknown, or more likely, does not exist after 30 years of intensive research, it would be more practical to look at the effects of *Azospirillum* spp. on the whole plant as an outcome of multiple mechanisms rather than a single mechanism operating at the organ, tissue, cellular, or molecular levels. Support for this notion is provided by an analysis of literature of many of the known cases of the effect of inoculation on the root to shoot (S/R) ratio that shows that the general effect of *Azospirillum*

spp. on the entire plant was largely overlooked. From the changes the bacteria produce in the S/R ratio, it appears that it also participates in the partitioning of carbon compounds within the plant, a phenomenon that is well recognized as multiparametric. The analysis provides supportive experimental data (although collected from many diverse studies) that indicate that the mode of action of *Azospirillum* spp. is probably composed of multiple mechanisms (Bashan and Dubrovsky, 1996). Additional supportive experimental data are provided by recent studies on polyamines (Cassan *et al.*, 2009a) and nitrogen fixation (Van Dommelen *et al.*, 2009) that were presented earlier.

4. CONCLUDING REMARKS AND A PROPOSAL

Today, the prevailing explanation for the effect of *Azospirillum* on plants is the production of an assortment of phytohormones, mainly IAA, altering the metabolism and morphology of the roots, yielding better mineral and water absorption, hence, higher yields. The contribution of nitrogen fixation is more controversial and, despite the increasing large volume of literature on other possible mechanisms, these are largely ignored by reviews on the topic of plant growth promotion, mostly evaluating PGPB in general.

In a comprehensive analysis of the knowledge about physiology, metabolic pathways, and molecular biology mechanisms of *Azospirillum* and their possible mode of action, it is apparent that phytohormones, especially IAA working in synchronization with other phytohormones produced by the bacterium, play a major role in various aspects of metabolism for growth. However, to attribute extremely complex phenomena for nonspecific causes of growth promotion in numerous plant species inoculated with many strains of *Azospirillum* having great differences in physiological traits, to one or a few substance (s) produced in abundance, mainly *in vitro*, is an oversimplification. Yet, it is, useful research tool for probing the mode of action of these bacteria.

There was, and still is, a disproportion between the large amount of knowledge on the bacterium cell and less knowledge about its interaction with the plants. In many aspects of interaction, such as mitigation of stresses or biological control, our knowledge about the mode of action is close to nil. Unfortunately but frequently, the knowledge about bacterial metabolisms *per se* is extrapolated to explain possible effects on plants without providing solid evidence that such activity do exist *in planta*.

Mutants that are defective in several traits are used in this field of research, but are employed on a smaller scale than in the related fields such as biological control of plant pathogens. For a more accurate determination of the role of phytohormones in promoting growth in general, and

IAA in particular, there is a need to obtain a mutant that is totally deficient in IAA production, but otherwise identical to the parent strain. Although several IAA-attenuated mutants were constructed, this goal has not yet been achieved. The same is true for other phytohormones. Additionally, to clearly state whether hormones are the main mechanism for promoting growth, we need to demonstrate that other proposed mechanisms have a minor role. Yet, there is much evidence to the contrary. These include the importance of nitrogen fixation under specific circumstances, including the postpara-nodule colonization (presented earlier) and new data collected under greenhouse and field conditions (Rodrigues *et al.*, 2008; Van Dommelen *et al.*, 2009). The overall accumulated evidence that nitrogen fixation plays a role in the association reconfirms that dismissal of nitrogen fixation, as a mechanism for plant growth reported in several reviews in recent years, is premature, and that nitrogen fixation should be reconsidered as a plausible comechanism. Additionally, the importance of signal molecules in initiating the cascade of events that induce a plant response, should be considered, perhaps in relation to root membranes (the main subcellular units responsible for mineral uptake detected in numerous associations of plant with *Azospirillum*). Many cases of mitigation of environmental stresses, possibly by mechanisms not envisioned so far or by a combination of several proposed mechanisms, as well as the possibly of limited biological control of plant pathogens, deserve critical evaluation and reconsideration.

The multitude of options for enhancing plant growth by inoculation with *Azospirillum* led us to propose the “Multiple Mechanisms Theory,” based on the assumption that there is no single mechanism involved in promoting plant growth with *Azospirillum*, but rather a combination of a few or many mechanisms in each specific case of inoculation. The mechanisms may vary with the plant species, the strain of *Azospirillum*, and environmental conditions prevailing during the interaction. The effect can be cumulative, as proposed earlier by the “additive hypothesis” (Bashan and Levany, 1990), where the effects of small mechanisms, operating at the same time or consecutively, create a larger final effect on the plant. The effect on plant growth can also be a result from tandem or cascading mechanisms in which one mechanism stimulates the other, which finally yields enhanced plant growth (such as the plausible relations among plant hormones, NO, membrane activities, and proliferation of root). Finally, promoting growth can be the result of a combination of unrelated mechanisms that operate according to environmental or agricultural conditions in a certain location. These include stress mitigation (salt, drought, toxic compounds) and biological control of pathogenic microflora. This inclusive kind of theory may close the gaps between competing theories and might lead to new insights about overlapping and cooperation among seemingly different mechanisms that affect plant growth than have been studied so far.

ACKNOWLEDGMENTS

Yoav Bashan participated in this essay in memory of the late Avner Bashan of Israel. We thank the following scientists for critical reading and insightful suggestions during manuscript preparation: Fabricio Cassan, University of Rio Cuarto, Argentina (phytohormones), Ivan Kennedy, Nitrogen Fixation Center, University of Sydney, Australia (N_2 -fixation), Cecilia Creus, University de Mar del Plata, Argentina and Michael Cohen, Sonoma State University, California (NO). Juan-Pablo Hernandez at CIBNOR, Mexico prepared the drawing. This review was mainly supported by The Bashan Foundation USA and partly by Consejo Nacional de Ciencia y Tecnología (CONACYT Investigacion Cientifica Basica 2005 50560-Z), and Secretaria de Medio Ambiente y Recursos Naturales of Mexico (SEMAR-NAT 23510).

REFERENCES

- Achard, P., Cheng, H., De Grauwe, L., Decat, J., Schoutteten, H., Moritz, T., Van Der Straeten, D., Peng, J., and Harberd, N. P. (2006). Integration of plant responses to environmentally activated phytohormones signals. *Science* **311**, 91–94.
- Alen'kina, S. A., Payusova, O. A., and Nikitina, V. E. (2006). Effect of *Azospirillum* lectins on the activities of wheat-root hydrolytic enzymes. *Plant Soil* **283**, 147–151.
- Alvarez, M. I., Sueldo, R. J., and Barassi, C. A. (1996). Effect of *Azospirillum* on coleoptile growth in wheat seedlings under water tress. *Cereal Res. Commun.* **24**, 101–107.
- Amooghaie, R., Mostajeran, A., and Emtiazi, G. (2002). The effect of compatible and incompatible *Azospirillum brasilense* strains on proton efflux of intact wheat roots–*Azospirillum* and proton efflux of wheat root. *Plant Soil* **243**, 155–160.
- Antonyuk, L. P., and Evseeva, N. V. (2006). Wheat lectin as a factor in plant–microbial communication and a stress response protein. *Microbiology (Moscow)* **75**, 470–475.
- Antonyuk, L. P., Fomina, O. R., Galkin, M. A., and Ignatov, V. V. (1993). The effect of wheat germ agglutinin on dinitrogen fixation, glutamine synthetase activity and ammonia excretion in *Azospirillum brasilense* Sp 245. *FEMS Microbiol. Lett.* **110**, 285–289.
- Antonyuk, L. P., Fomina, O. R., Kalinina, A., Semenov, S., Nesmeyanova, M., and Ignatov, V. (1995). Wheat lectin possibly serves as a signal molecule in the *Azospirillum*–wheat association. *NATO ASI Ser. Ser. G* **37**, 319–324.
- Araujo, M. S., Baura, V. A., Souza, E. M., Benelli, E. M., Rigo, L. U., Steffens, M. B. R., Pedrosa, F. O., and Chubatsu, L. S. (2004a). In vitro uridylylation of the *Azospirillum brasilense* N-signal transducing GlnZ protein. *Protein Expr. Purif.* **33**, 19–24.
- Araujo, L. M., Monteiro, R. A., Souza, E. M., Steffens, M. B. R., Rigo, L. U., Pedrosa, F. O., and Chubatsu, L. S. (2004b). GlnB is specifically required for *Azospirillum brasilense* NifA activity in *Escherichia coli*. *Res. Microbiol.* **155**, 491–495.
- Aziz, A., Martin-Tanguy, J., and Larher, F. (1997). Plasticity of polyamine metabolism associated with high osmotic stress in rape leaf discs and with ethylene treatment. *Plant Growth Regul.* **21**, 153–163.
- Babu, R. S., Sankaranarayanan, C., and Jothi, G. (1998). Management of *Pratylenchus zea* on maize by biofertilizers and VAM. *Indian J. Nematol.* **28**, 77–80.
- Baca, B. E., Soto-Urzuu, L., Xochihua-Corona, Y. G., and Cuervo-Garcia, A. (1994). Characterization of two aromatic amino acid aminotransferases and production of indoleacetic acid in *Azospirillum* strains. *Soil Biol. Biochem.* **26**, 57–63.
- Bacilio, M., Vazquez, P., and Bashan, Y. (2003). Alleviation of noxious effects of cattle ranch composts on wheat seed germination by inoculation with *Azospirillum* spp. *Biol. Fertil. Soils* **38**, 261–266.

- Bacilio, M., Rodriguez, H., Moreno, M., Hernandez, J.-P., and Bashan, Y. (2004). Mitigation of salt stress in wheat seedlings by a *gfp*-tagged *Azospirillum lipoferum*. *Biol. Fertil. Soils* **40**, 188–193.
- Bakanchikova, T. I., Lobanok, E. V., Pavlova Ivanova, L. K., Red'kina, T. V., Nagapetyan, Z. A., and Majsuryan, A. N. (1993). Inhibition of tumor formation process in dicotyledonous plants by *Azospirillum brasilense* strains. *Mikrobiologiya (Russian Federation)* **62**, 515–523, (in Russian).
- Baldani, J. I., and Baldani, V. L. D. (2005). History on the biological nitrogen fixation research in graminaceous plants: Special emphasis on the Brazilian experience. *An. Acad. Bras. Cienc.* **77**, 549–579.
- Bansal, R. K., Dahiya, R. S., Lakshminarayana, K., Suneja, S., Anand, R. C., and Narula, N. (1999). Effect of rhizospheric bacteria on plant growth of wheat infected with *Heterodera avenae*. *Nematologia Mediterranea* **27**, 311–314.
- Barassi, C. A., Ayrault, G., Creus, C. M., Sueldo, R. J., and Sobrero, M. T. (2006). Seed inoculation with *Azospirillum* mitigates NaCl effects on lettuce. *Sci. Hortic.* **109**, 8–14.
- Barbieri, P., and Galli, E. (1993). Effect on wheat root development of inoculation with an *Azospirillum brasilense* mutant with altered indole-3-acetic acid production. *Res. Microbiol.* **144**, 69–75.
- Barea, J. M., Navarro, M., and Montoya, E. (1976). Production of plant growth regulators by rhizosphere phosphate-solubilizing bacteria. *J. Appl. Bacteriol.* **40**, 129–134.
- Bartels, D., and Sunkar, R. (2005). Drought and salt tolerance in plants. *Crit. Rev. Plant Sci.* **24**, 23–58.
- Barton, L. L., Johnson, G. V., and Orbock Miller, S. (1986). The effect of *Azospirillum brasilense* on iron absorption and translocation by sorghum. *J. Plant Nutr.* **9**, 557–565.
- Bashan, Y. (1990). Short exposure to *Azospirillum brasilense* Cd inoculation enhanced proton efflux in intact wheat roots. *Can. J. Microbiol.* **36**, 419–425.
- Bashan, Y. (1991). Changes in membrane potential of intact soybean root elongation zone cells induced by *Azospirillum brasilense*. *Can. J. Microbiol.* **37**, 958–963.
- Bashan, Y. (1998). Inoculants of plant growth-promoting bacteria for use in agriculture. *Biotechnol. Adv.* **16**, 729–770.
- Bashan, Y., and de-Bashan, L. E. (2002a). Protection of tomato seedlings against infection by *Pseudomonas syringae* pv *tomato* by using the plant growth-promoting bacterium *Azospirillum brasilense*. *Appl. Environ. Microbiol.* **68**, 2637–2643.
- Bashan, Y., and de-Bashan, L. E. (2002b). Reduction of bacterial speck (*Pseudomonas syringae* pv *tomato*) of tomato by combined treatments of plant growth-promoting bacterium, *Azospirillum brasilense*, streptomycin sulfate, and chemo-thermal seed treatment. *Eur. J. Plant Pathol.* **108**, 821–829.
- Bashan, Y., and Dubrovsky, J. G. (1996). *Azospirillum* spp. participation in dry matter partitioning in grasses at the whole plant level. *Biol. Fertil. Soils* **23**, 435–440.
- Bashan, Y., and Holguin, G. (1997). *Azospirillum*–plant relationships: Environmental and physiological advances (1990–1996). *Can. J. Microbiol.* **43**, 103–121.
- Bashan, Y., and Levanony, H. (1990). Current status of *Azospirillum* inoculation technology: *Azospirillum* as a challenge for agriculture. *Can. J. Microbiol.* **36**, 591–608.
- Bashan, Y., and Levanony, H. (1991). Alterations in membrane potential and in proton efflux in plant roots induced by *Azospirillum brasilense*. *Plant Soil* **137**, 99–103.
- Bashan, Y., Levanony, H., and Mitiku, G. (1989a). Changes in proton efflux of intact wheat roots induced by *Azospirillum brasilense* Cd. *Can. J. Microbiol.* **35**, 691–697.
- Bashan, Y., Singh, M., and Levanony, H. (1989b). Contribution of *Azospirillum brasilense* Cd to growth of tomato seedlings is not through nitrogen fixation. *Can. J. Bot.* **67**, 2429–2434.
- Bashan, Y., Harrison, S. K., and Whitmoyer, R. E. (1990). Enhanced growth of wheat and soybean plants inoculated with *Azospirillum brasilense* is not necessarily due to general enhancement of mineral uptake. *Appl. Environ. Microbiol.* **56**, 769–775.

- Bashan, Y., Alcaraz-Melendez, L., and Toledo, G. (1992). Responses of soybean and cowpea root membranes to inoculation with *Azospirillum brasilense*. *Symbiosis* **13**, 217–228.
- Bashan, Y., Holguin, G., and de-Bashan, L. E. (2004). *Azospirillum*–plant relationships: Physiological, molecular, agricultural, and environmental advances (1997–2003). *Can. J. Microbiol.* **50**, 521–577.
- Bashan, Y., Bustillos, J. J., Leyva, L. A., Hernandez, J.-P., and Bacilio, M. (2006). Increase in auxiliary photoprotective photosynthetic pigments in wheat seedlings induced by *Azospirillum brasilense*. *Biol. Fertil. Soils* **42**, 279–285.
- Belimov, A., and Dietz, K.-J. (2000). Effect of associative bacteria on element composition of barley seedlings grown in solution culture at toxic cadmium concentrations. *Microbiol. Res.* **155**, 113–121.
- Belimov, A. A., Kunakova, A. M., Safronova, V. I., Stepanok, V. V., Yudkin, L. Y., Alekseev, Y. V., and Kozhemyakov, A. P. (2004). Employment of rhizobacteria for the inoculation of barley plants cultivated in soil contaminated with lead and cadmium. *Microbiology (Moscow)* **73**, 99–106.
- Boddey, R. M., and Döbereiner, J. (1988). Nitrogen fixation associated with grasses and cereals: Recent results and perspectives for future research. *Plant Soil* **108**, 53–65.
- Bothe, H., Körsgen, H., Lehmacher, T., and Hundeshagen, B. (1992). Differential effects of *Azospirillum*, auxin and combined nitrogen on the growth of the roots of wheat. *Symbiosis* **13**, 167–179.
- Bottini, R., Fulchieri, M., Pearce, D., and Pharis, R. P. (1989). Identification of gibberellins A₁, A₃ and iso-A₃ in cultures of *Azospirillum lipoferum*. *Plant Physiol.* **90**, 45–47.
- Bottini, R., Cassan, F., and Piccoli, P. (2004). Gibberellin production by bacteria and its involvement in plant growth promotion and yield increase. *Appl. Microbiol. Biotechnol.* **65**, 497–503.
- Bouillant, M. L., Miché, L., Ouedraogo, O., Alexandre, G., Jacoud, C., Sallé, G., and Bally, R. (1997). Inhibition of *Striga* seed germination associated with sorghum growth promotion by soil bacteria. *C.R. Acad. Sci. Paris-Sciences de la vie* **320**, 159–162.
- Cacciari, I., Lippi, D., Pietrosanti, T., and Pietrosanti, W. (1989). Phytohormone-like substances produced by single and mixed diazotrophic cultures of *Azospirillum* spp. and *Arthrobacter*. *Plant Soil* **115**, 151–153.
- Carreño-Lopez, R., Campos-Reales, N., Elmerich, C., and Baca, B. E. (2000). Physiological evidence for differently regulated tryptophan-dependent pathways for indole-3-acetic acid synthesis in *Azospirillum brasilense*. *Mol. Gen. Genet.* **264**, 521–530.
- Carrillo, A. E., Li, C. Y., and Bashan, Y. (2002). Increased acidification in the rhizosphere of cactus seedlings induced by *Azospirillum brasilense*. *Naturwissenschaften* **89**, 428–432.
- Cassan, F., Bottini, R., Schneider, G., and Piccoli, P. (2001a). *Azospirillum brasilense* and *Azospirillum lipoferum* hydrolyze conjugates of GA₂₀ and metabolize the resultant aglycones to GA₁ in seedlings of rice dwarf mutants. *Plant Physiol.* **125**, 2053–2058.
- Cassan, F. D., Lucangeli, C. D., Bottini, R., and Piccoli, P. N. (2001b). *Azospirillum* spp. metabolize [17, 17-²H₂] gibberellin A₂₀ to [17, 17-²H₂] gibberellin A₁ in vivo in dy rice mutant seedlings. *Plant Cell Physiol.* **42**, 763–767.
- Cassan, F., Maiale, S., Masciarelli, O., Vidal, A., Luna, V., and Ruiz, O. (2009a). Cadaverine production by *Azospirillum brasilense* and its possible role in plant growth promotion and osmotic stress mitigation. *Eur. J. Soil Biol.* **45**, 12–19.
- Cassan, F., Perrig, D., Sgroy, V., Masciarelli, O., Penna, C., and Luna, V. (2009b). *Azospirillum brasilense* Az39 and *Bradyrhizobium japonicum* E109, inoculated singly or in combination, promote seed germination and early seedling growth in corn (*Zea mays* L.) and soybean (*Glycine max* L.). *Eur. J. Soil Biol.* **45**, 28–35.
- Castellanos, T., Ascencio, F., and Bashan, Y. (1998). Cell-surface lectins of *Azospirillum* spp. *Curr. Microbiol.* **36**, 241–244.

- Castellen, P., Wassem, R., Monteiro, R. A., Magalhães Cruz, L., Steffens, M. B. R., Chubatsu, L. S., de Souza, E. M., and Pedrosa, F. O. (2009). Structural organization of the *glnBA* region of the *Azospirillum brasilense* genome. *Eur. J. Soil Biol.* **45**, 100–105.
- Chang, T. T., and Li, C. Y. (1998). Weathering of limestone, marble, and calcium phosphate by ectomycorrhizal fungal and associated microorganisms. *Taiwan J. For. Sci.* **13**, 85–90.
- Cho, J. I., Hong, K. W., and Kang, K. J. (2000). Control of aflatoxin production of *Aspergillus flavus* by inhibitory action of antagonistic bacteria. *J. Microbiol. Biotechnol.* **10**, 154–160.
- Choudhury, A. T. M. A., and Kennedy, I. R. (2004). Prospects and potentials for systems of biological nitrogen fixation in sustainable rice production. *Biol. Fertil. Soils* **39**, 219–227.
- Christiansen-Weniger, C. (1992). N₂-fixation by ammonium-excreting *Azospirillum brasilense* in auxin-induced tumours of wheat (*Triticum aestivum* L.). *Biol. Fertil. Soils* **12**, 100–106.
- Christiansen-Weniger, C. (1994). Para-nodule induction in maize with indoleacetic acid (IAA) and its infection with ammonium-excreting *Azospirillum brasilense*. In "Nitrogen-Fixation with Non-Legumes" (N. A. Hegazi, M. Fayez, and M. Monib, Eds.), pp. 525–536. American University of Cairo Press, Cairo, Egypt.
- Christiansen-Weniger, C. (1997). Ammonium-excreting *Azospirillum brasilense* C3:gusA inhabiting induced tumors along stem and roots of rice. *Soil Biol. Biochem.* **29**, 943–950.
- Christiansen-Weniger, C. (1998). Endophytic establishment of diazotrophic bacteria in auxin-induced tumors of cereal crops. *Crit. Rev. Plant Sci.* **17**, 55–76.
- Christiansen-Weniger, C., and van Veen, J. A. (1991). NH₄⁺-excreting *Azospirillum brasilense* mutants enhance the nitrogen supply of a wheat host. *Appl. Environ. Microbiol.* **57**, 3006–3012.
- Cohen, A. C., Bottini, R., and Piccoli, P. N. (2008). *Azospirillum brasilense* Sp 245 produces ABA in chemically-defined culture medium and increases ABA content in arabidopsis plants. *Plant Growth Regul.* **54**, 97–103.
- Cohen, A. C., Travaglia, C. N., Bottini, R., and Piccoli, P. N. (2009). Participation of abscisic acid and gibberellins produced by endophytic *Azospirillum* in the alleviation of drought effects in maize. *Botany* **87**, 455–462.
- Cohen, M. F., Lamattina, L., and Yamasaki, H. (2010). Nitric oxide signaling by plant-associated bacteria. In "Nitric Oxide in Plant Physiology" (S. Hayat, M. Mori, J. Pichtel, and A. Ahmad, Eds.), pp. 161–172. Wiley-VCH, Weinheim, Germany.
- Correa-Aragunde, N., Graziano, M., and Lamattina, L. (2004). Nitric oxide plays a central role in determining lateral root development in tomato. *Planta* **218**, 900–905.
- Correa-Aragunde, N., Graziano, M., Chevalier, C., and Lamattina, L. (2006). Nitric oxide modulates the expression of cell cycle regulatory genes during lateral root formation in tomato. *J. Exp. Bot.* **57**, 581–588.
- Costacurta, A., and Vanderleyden, J. (1995). Synthesis of phytohormones by plant-associated bacteria. *Crit. Rev. Microbiol.* **21**, 1–18.
- Costacurta, A., Keijers, V., and Vanderleyden, J. (1994). Molecular cloning and sequence analysis of an *Azospirillum brasilense* indole-3-pyruvate decarboxylase gene. *Mol. Gen. Genet.* **243**, 463–472.
- Creus, C. M., Sueldo, R. J., and Barassi, C. A. (1997). Shoot growth and water status in *Azospirillum*-inoculated wheat seedlings grown under osmotic and salt stresses. *Plant Physiol. Biochem.* **35**, 939–944.
- Creus, C. M., Sueldo, R. J., and Barassi, C. A. (1998). Water relations in *Azospirillum*-inoculated wheat seedlings under osmotic stress. *Can. J. Bot.* **76**, 238–244.
- Creus, C. M., Sueldo, R. J., and Barassi, C. A. (2004). Water relations and yield in *Azospirillum*-inoculated wheat exposed to drought in the field. *Can. J. Bot.* **82**, 273–281.

- Creus, C. M., Graziano, M., Casanovas, E. M., Pereyra, M. A., Simontacchi, M., Puntarulo, S., Barassi, C. A., and Lamattina, L. (2005). Nitric oxide is involved in the *Azospirillum brasilense*-induced lateral root formation in tomato. *Planta* **221**, 297–303.
- Crozier, A., Arruda, P., Jasmin, J. M., Monteiro, A. M., and Sandberg, G. (1988). Analysis of indole-3-acetic acid and related indoles in culture medium from *Azospirillum lipoferum* and *Azospirillum brasilense*. *Appl. Environ. Microbiol.* **54**, 2833–2837.
- Dadon, T., Bar Nun, N., and Mayer, A. M. (2004). A factor from *Azospirillum brasilense* inhibits germination and radicle growth of *Orobanche aegyptiaca*. *Isr. J. Plant Sci.* **52**, 83–86.
- de Campos, S. B., Roesch, L. F. W., Zanettini, M. H. B., and Passaglia, L. M. P. (2006). Relationship between in vitro enhanced nitrogenase activity of an *Azospirillum brasilense* Sp7 mutant and its growth-promoting activities *in situ*. *Curr. Microbiol.* **53**, 43–47.
- Deaker, R., and Kennedy, I. R. (2001). Improved potential for nitrogen fixation in *Azospirillum brasilense* Sp7-S associated with wheat: *nifH* expression as a function of oxygen pressure. *Acta Biotechnol.* **21**, 3–17.
- de-Bashan, L. E., and Bashan, Y. (2008). Joint immobilization of plant growth-promoting bacteria and green microalgae in alginate beads as an experimental model for studying plant–bacterium interactions. *Appl. Environ. Microbiol.* **74**, 6797–6802.
- de-Bashan, L. E., Hernandez, J.-P., Morey, T., and Bashan, Y. (2004). Microalgae growth-promoting bacteria as “helpers” for microalgae: A novel approach for removing ammonium and phosphorus from municipal wastewater. *Water Res.* **38**, 466–474.
- de-Bashan, L. E., Antoun, H., and Bashan, Y. (2005). Cultivation factors and population size control uptake of nitrogen by the microalgae *Chlorella vulgaris* when interacting with the microalgae growth-promoting bacterium *Azospirillum brasilense*. *FEMS Microbiol. Ecol.* **54**, 197–203.
- de-Bashan, L. E., Antoun, H., and Bashan, Y. (2008a). Involvement of indole-3-acetic-acid produced by the growth-promoting bacterium *Azospirillum* spp. in promoting growth of *Chlorella vulgaris*. *J. Phycol.* **44**, 938–947.
- de-Bashan, L. E., Trejo, A., Huss, V. A. R., Hernandez, J.-P., and Bashan, Y. (2008b). *Chlorella sorokiniana* UTEX 2805, a heat and intense, sunlight-tolerant microalga with potential for removing ammonium from wastewater. *Bioresour. Technol.* **99**, 4980–4989.
- Deubel, A., Gransee, A., and Merbach, W. (2000). Transformation of organic rhizodepositions by rhizosphere bacteria and its influence on the availability of tertiary calcium phosphate. *J. Plant Nutr. Soil Sci.* **163**, 387–392.
- Diaz-Zorita, M., and Fernández-Canigia, M. V. (2009). Field performance of a liquid formulation of *Azospirillum brasilense* on dryland wheat productivity. *Eur. J. Soil Biol.* **45**, 3–11.
- Dobbelaere, S., Croonenborghs, A., Thys, A., Vande Broek, A., and Vanderleyden, J. (1999). Phytostimulatory effect of *Azospirillum brasilense* wild type and mutant strains altered in IAA production on wheat. *Plant Soil* **212**, 155–164.
- Dobbelaere, A., Vanderleyden, J., and Okon, Y. (2003). Plant growth-promoting effects of diazotrophs in the rhizosphere. *Crit. Rev. Plant Sci.* **2**, 107–149.
- Döbereiner, J., and Day, L. (1976). Associative symbiosis in tropical grasses: Characterization of microorganisms and dinitrogen fixing sites. In “Proceedings First International Symposium on Nitrogen Fixation” (W. E. Newton and C. J. Nyman, Eds.), pp. 518–538. Washington State University Press, Pullman.
- Doroshenko, E. V., Boulygina, E. S., Spiridonova, E. M., Tourova, T. P., and Kravchenko, I. K. (2007). Isolation and characterization of nitrogen-fixing bacteria of the genus *Azospirillum* from the soil of a *Sphagnum* peat bog. *Microbiology (Moscow)* **76**, 93–101.
- Duby, G., and Boutry, M. (2009). The plant plasma membrane proton pump ATPase: A highly regulated P-type ATPase with multiple physiological roles. *Pflügers Archiv. Eur. J. Physiol.* **457**, 645–655.

- Eckert, B., Weber, O. B., Kirchhof, G., Halbritter, A., Stoffels, M., and Hartmann, A. (2001). *Azospirillum dobereineriae* sp. nov., a nitrogen-fixing bacterium associated with the C₄-grass *Miscanthus*. *Int. J. Syst. Evol. Microbiol.* **51**, 17–26.
- El-Khawass, H., and Adachi, K. (1999). Identification and quantification of auxins in culture media of *Azospirillum* and *Klebsiella* and their effect on rice roots. *Biol. Fertil. Soils* **28**, 377–381.
- El-Komy, H. M., Hamdia, M. A., and El-Baki, G. K. A. (2003). Nitrate reductase in wheat plants grown under water stress and inoculated with *Azospirillum* spp. *Biol. Plantarum* **46**, 281–287.
- Fallik, E., Okon, Y., Epstein, E., Goldman, A., and Fischer, M. (1989). Identification and quantification of IAA and IBA in *Azospirillum brasilense*-inoculated maize roots. *Soil Biol. Biochem.* **21**, 147–153.
- Fallik, E., Sarig, S., and Okon, Y. (1994). Morphology and physiology of plant roots associated with *Azospirillum*. In “*Azospirillum*/Plant Associations” (Y. Okon, Ed.), pp. 77–86. CRC Press, Boca Raton, FL, USA.
- Feng, L., and Kennedy, I. R. (1997). Biodegradation and plant protection from the herbicide 2, 4-D by plant-microbial associations in cotton production systems. *Biotechnol. Bioengineering* **54**, 513–519.
- Ferreira, M. C. B., Fernandes, M. S., and Döbereiner, J. (1987). Role of *Azospirillum brasilense* nitrate reductase in nitrate assimilation by wheat plants. *Biol. Fertil. Soils* **4**, 47–53.
- Fulchieri, M., Lucangeli, C., and Bottini, R. (1993). Inoculation with *Azospirillum lipoferum* affects growth and gibberellin status on corn seedling roots. *Plant Cell Physiol.* **34**, 1305–1309.
- Furina, E. K., Bonartseva, G. A., and Lvov, N. P. (1999). Effect of various concentrations of oxygen, molybdenum, and nitrate on nitrogen fixation and denitrification in *Azospirillum lipoferum*. *Appl. Biochem. Microbiol.* **35**, 44–47.
- Gamarnik, A., and Frydman, R. (1991). Cadaverine an essential diamine for the normal root development of germinating soybean (*Glycine max*) seeds. *Plant Physiol.* **97**, 778–785.
- Garcia de Salamone, I. E., Döbereiner, J., Urquiaga, S., and Boddey, R. M. (1997). Biological nitrogen fixation in *Azospirillum* strain-maize genotype associations as evaluated by the ¹⁵N isotope dilution technique. *Biol. Fertil. Soils* **23**, 249–256.
- Gaxiola, R. A., Palmaren, M. G., and Schumacher, K. (2007). Plant proton pumps. *FEBS Lett.* **581**, 2204–2214.
- Giraud, E., et al. (2007). Legumes symbioses. Absence of genes in nod in photosynthetic bradyrhizobia. *Science* **316**, 1307–1312, (+33 authors).
- Glick, B. R., Patten, C. L., Holguin, G., and Penrose, D. M. (1999). Biochemical and genetic mechanisms used by plant growth promoting bacteria. Imperial College Press, London, UK.
- Gonçalves, A. F. S., and de Oliveira, R. G. B. (1998). Cyanide production by Brazilian strains of *Azospirillum*. *Rev. Microbiol.* **29**, 36–39.
- Gonzalez, L. E., and Bashan, Y. (2000). Increased growth of the microalga *Chlorella vulgaris* when coimmobilized and cocultured in alginate beads with the plant-growth-promoting bacterium *Azospirillum brasilense*. *Appl. Environ. Microbiol.* **66**, 1527–1531.
- Gupta, G. K., and Singh, D. (1999). Role of edaphic factors in the development of downy mildew (*Sclerospora graminicola*) in pearl millet. *J. Agric. Sci.* **133**, 61–68.
- Hadas, R., and Okon, Y. (1987). Effect of *Azospirillum brasilense* inoculation on root morphology and respiration in tomato seedlings. *Biol. Fertil. Soils* **5**, 241–247.
- Hamdia, M. A., and El-Komy, H. M. (1997). Effect of salinity, gibberellic acid and *Azospirillum* inoculation on growth and nitrogen uptake of *Zea mays*. *Biol. Plant.* **40**, 109–120.
- Hamdia, A. B. E., Shaddad, M. A. K., and Doaa, M. M. (2004). Mechanisms of salt tolerance and interactive effects of *Azospirillum brasilense* inoculation on maize cultivars grown under salt stress conditions. *Plant Growth Regul.* **44**, 165–174.

- Harari, A., Kigel, J., and Okon, Y. (1988). Involvement of IAA in the interaction between *Azospirillum brasilense* and *Panicum miliaceum* roots. *Plant Soil* **110**, 275–282.
- Hartmann, A., and Bashan, Y. (2009). Ecology and application of *Azospirillum* and other plant growth-promoting bacteria (PGPB)—Special issue. *Eur. J. Soil Biol.* **45**, 1–2.
- Hartmann, A., and Zimmer, W. (1994). Physiology of *Azospirillum*. In “*Azospirillum*/Plant Association” (Y. Okon, Ed.), pp. 15–39. CRC Press, Boca Raton, FL, USA.
- Hartmann, A., Singh, M., and Klingmüller, W. (1983). Isolation and characterization of *Azospirillum* mutants excreting high amounts of indoleacetic acid. *Can. J. Microbiol.* **29**, 916–923.
- Hassouna, M. G., El-Saedy, M. A. M., and Saleh, H. M. A. (1998). Biocontrol of soil-borne plant pathogens attacking cucumber (*Cucumis sativus*) by rhizobacteria in a semiarid environment. *Arid Soil Res. Rehab.* **12**, 345–357.
- Hernandez, J.-P., de-Bashan, L. E., and Bashan, Y. (2006). Starvation enhances phosphorus removal from wastewater by the microalga *Chlorella* spp. co-immobilized with *Azospirillum brasilense*. *Enzyme Microb. Technol.* **38**, 190–198.
- Holguin, G., and Bashan, Y. (1996). Nitrogen-fixing by *Azospirillum brasilense* Cd is promoted when co-cultured with a mangrove rhizosphere bacterium (*Staphylococcus* sp.). *Soil Biol. Biochem.* **28**, 1651–1660.
- Holguin, G., and Glick, B. R. (2001). Expression of the ACC deaminase gene from *Enterobacter cloacae* UW4 in *Azospirillum brasilense*. *Microb. Ecol.* **41**, 281–288.
- Holguin, G., and Glick, B. R. (2003). Transformation of *Azospirillum brasilense* Cd with an ACC deaminase gene (*adcS*) from *Enterobacter cloacae* UW4 fused to the Tetr gene promoter improves its fitness and plant growth promoting ability. *Microb. Ecol.* **46**, 122–133.
- Horemans, S., De Koninck, K., Neuray, J., Hermans, R., and Vlassak, K. (1986). Production of plant growth substances by *Azospirillum* sp. and other rhizosphere bacteria. *Symbiosis* **2**, 341–346.
- Huang, A. X., She, X. P., Huang, C., and Song, T. S. (2007). The dynamic distribution of NO and NADPH-diaphorase activity during IBA induced adventitious root formation. *Physiol. Plant.* **130**, 240–249.
- Huergo, L., Chubatsu, L., Souza, E., Pedrosa, F., Steffens, M., and Merrick, M. (2006a). Interactions between PII proteins and the nitrogenase regulatory enzymes DraT and DraG in *Azospirillum brasilense*. *FEBS Lett.* **580**, 5232–5236.
- Huergo, L. F., Souza, E. M., Araujo, M. S., Pedrosa, F. O., Chubatsu, L. S., Steffens, M. B. R., and Merrick, M. (2006b). ADP-ribosylation of dinitrogenase reductase in *Azospirillum brasilense* is regulated by AmtB-dependent membrane sequestration of DraG. *Mol. Microbiol.* **59**, 326–337.
- Huergo, L. F., Merrick, M., Monteiro, R. A., Chubatsu, L. S., Steffens, M. B. R., Pedrosa, F. O., and Souza, E. M. (2009). *In vitro* interactions between the P_{II} proteins and the nitrogenase regulatory enzymes Dinitrogenase Reductase ADP-ribosyltransferase (DraT) and Dinitrogenase Reductase-activating Glycohydrolase (DraG) in *Azospirillum brasilense*. *J. Biol. Chem.* **284**, 6674–6682.
- Ignatov, O. V., Kamnev, A. A., Markina, L. N., Antonyuk, L. P., Colina, M., and Ignatov, V. V. (2001). Electrooptical properties of cells of the soil nitrogen-fixing bacterium *Azospirillum brasilense*: Effects of copper ions. *Appl. Biochem. Microbiol.* **37**, 219–223.
- Ismail, A. E., and Hasabo, S. A. (2000). Evaluation of some new Egyptian commercial biofertilizers, plant nutrients and a biocide against *Meloidogyne incognita* root knot nematode infecting sunflower. *Pak. J. Nematol.* **18**, 39–49.
- Jain, D. K., and Patriquin, D. G. (1984). Root hair deformation, bacterial attachment, and plant growth in wheat–*Azospirillum* associations. *Appl. Environ. Microbiol.* **48**, 1208–1213.

- Jain, D. K., and Patriquin, D. G. (1985). Characterization of a substance produced by *Azospirillum* which causes branching of wheat root hairs. *Can. J. Microbiol.* **31**, 206–210.
- Kamnev, A. A., Renou-Gonnord, M. F., Antonyuk, L. P., Colina, M., Chernyshev, A. V., Frolov, I., and Ignatov, V. V. (1997a). Spectroscopic characterization of the uptake of essential and xenobiotic metal cations in cells of the soil bacterium *Azospirillum brasilense*. *Biochem. Mol. Biol. Int.* **41**, 123–130.
- Kamnev, A. A., Ristic, M., Antonyuk, L. P., Chernyshev, A. V., and Ignatov, V. V. (1997b). Fourier transform infrared spectroscopic study of intact cells of the nitrogen-fixing bacterium *Azospirillum brasilense*. *J. Mol. Struct.* **408**, 201–205.
- Kamnev, A. A., Antonyuk, L. P., Colina, M., Chernyshev, A. V., and Ignatov, V. V. (1999a). Investigation of a microbially produced structural modification of magnesium-ammonium orthophosphate. *Monatsh. Chem.* **130**, 1431–1442.
- Kamnev, A. A., Antonyuk, L. P., Matora, L. Y., Serebrennikova, O. B., Sumaroka, M. V., Colina, M., Renou Gonnord, M. F., and Ignatov, V. V. (1999b). Spectroscopic characterization of cell membranes and their constituents of the plant-associated soil bacterium *Azospirillum brasilense*. *J. Mol. Struct.* **481**, 387–393.
- Kamnev, A. A., Tarantilis, P. A., Antonyuk, L. P., Bespalova, L. A., Polissiou, M. G., Colina, M., Gardiner, P. H. E., and Ignatov, V. V. (2001). Fourier transform Raman spectroscopic characterization of cells of the plant-associated soil bacterium *Azospirillum brasilense* Sp7. *J. Mol. Struct.* **536**, 199–207.
- Kamnev, A. A., Antonyuk, L. P., Tugarova, A. V., Tarantilis, P. A., Polissiou, M. G., and Gardiner, P. H. E. (2002). Fourier transform infrared spectroscopic characterization of heavy metal-induced metabolic changes in the plant-associated soil bacterium *Azospirillum brasilense* Sp7. *J. Mol. Struct.* **610**, 127–131.
- Kamnev, A. A., Tugarova, A. V., Antonyuk, L. P., Tarantilis, P. A., Polissiou, M. G., and Gardiner, P. H. E. (2005). Effects of heavy metals on plant-associated rhizobacteria: Comparison of endophytic and non-endophytic strains of *Azospirillum brasilense*. *J. Trace Elem. Med. Biol.* **19**, 91–95.
- Kamnev, A. A., Tugarova, A. V., and Antonyuk, L. P. (2007). Endophytic and epiphytic strains of *Azospirillum brasilense* respond differently to heavy metal stress. *Microbiology (Moscow)* **76**, 809–811.
- Kapulnik, Y., Kigel, J., Okon, Y., Nur, I., and Henis, Y. (1981). Effect of *Azospirillum* inoculation on some growth parameters and N content of wheat, sorghum and panicum. *Plant Soil* **61**, 65–70.
- Kapulnik, Y., Gafni, R., and Okon, Y. (1985a). Effect of *Azospirillum* spp. inoculation on root development and NO₃– uptake in wheat (*Triticum aestivum* cv. Miriam) in hydroponic systems. *Can. J. Bot.* **63**, 627–631.
- Kapulnik, Y., Okon, Y., and Henis, Y. (1985b). Changes in root morphology of wheat caused by *Azospirillum* inoculation. *Can. J. Microbiol.* **31**, 881–887.
- Katupitiya, S., Millet, J., Vesk, M., Viccars, L., Zeman, A., Lidong, Z., Elmerich, C., and Kennedy, I. R. (1995a). A mutant of *Azospirillum brasilense* Sp7 impaired in flocculation with a modified colonization pattern and superior nitrogen fixation in association with wheat. *Appl. Environ. Microbiol.* **61**, 1987–1995.
- Katupitiya, S., New, P. B., Elmerich, C., and Kennedy, I. R. (1995b). Improved N₂ fixation in 2, 4-D treated wheat roots associated with *Azospirillum lipoferum*: Studies of colonization using reporting genes. *Soil Biol. Biochem.* **27**, 447–452.
- Kavitha, K., Meenakumari, K. S., and Sivaprasad, P. (2003). Effect of dual inoculation of native arbuscular mycorrhizal fungi and *Azospirillum* on suppression of damping off in chilli. *Indian Phytopathol.* **56**, 112–113.
- Kennedy, I. R. (1994). Auxin-induced N₂-fixing associations between *Azospirillum brasilense* and wheat. In “Nitrogen-Fixation with Nonlegumes” (N. A. Hegazi, M. Fayed, and M. Monib, Eds.), pp. 513–523. American University of Cairo Press, Cairo, Egypt.

- Kennedy, I. R., and Islam, N. (2001). The current and potential contribution of asymbiotic nitrogen fixation to nitrogen requirements on farms: A review. *Aust. J. Exp. Agric.* **41**, 447–457.
- Kennedy, I. R., and Tchan, Y. T. (1992). Biological nitrogen-fixation in non-leguminous field crops: Recent advances. *Plant Soil* **141**, 93–118.
- Kennedy, I. R., Pereg-Gerk, L. L., Wood, C., Deaker, R., Gilchrist, K., and Katupitiya, S. (1997). Biological nitrogen fixation in nonleguminous field crops: Facilitating the evolution between *Azospirillum* and wheat. *Plant Soil* **194**, 65–79.
- Kennedy, I. R., Choudhury, A. T. M. A., and Kecskes, M. L. (2004). Non-symbiotic bacterial diazotrophs in crop-farming systems: can their potential for plant growth promotion be better exploited? *Soil Biol. Biochem.* **36**, 1229–1244.
- Khan, M. R., and Kounsar, K. (2000). Effect of seed treatment with certain bacteria and fungi on the growth of mungbean and reproduction of *Meloidogyne incognita*. *Nematol. Mediterr.* **28**, 221–226.
- Kishore, P. (1998). Response of sorghum variety Pusa chari -121 to carrier based inoculants (*Azotobacter* and *Azospirillum*), fermented residue and shootfly (*Atherigona soccata* Rondani) under field conditions. *J. Entomol. Res.* **22**, 101–105.
- Klassen, G., Souza, E. M., Yates, M. G., Rigo, L. U., Costa, R. M., Inaba, J., and Pedrosa, F. O. (2005). Nitrogenase switch-off by ammonium ions in *Azospirillum brasilense* requires the GlnB nitrogen signal-transducing protein. *Appl. Environ. Microbiol.* **71**, 5637–5641.
- Kucey, R. M. N. (1988). Alteration of size of wheat root systems and nitrogen fixation by associative nitrogen-fixing bacteria measured under field conditions. *Can. J. Microbiol.* **34**, 735–739.
- Kundu, B. S., Sangwan, P., Sharma, P. K., and Nandwal, A. S. (1997). Response of pearl millet to phytohormones produced by *Azospirillum brasilense*. *Indian J. Plant Physiol.* **2**, 101–104.
- Kuznetsov, V., Radyukina, N., and Shevyakova, N. (2006). Polyamines and stress: Biological role, metabolism, and regulation. *Russ. J. Plant Physiol.* **53**, 583–604.
- Lamattina, L., and Polacco, J. (2007). Nitric Oxide in Plant Growth, Development and Stress Physiology. *Plant Cell Monographs*. Springer-Verlag, Berlin Heidelberg, Germany.
- Lamattina, L., García-Mata, C., Graziano, M., and Pagnussat, G. (2003). Nitric oxide: The versatility of an extensive signal molecule. *Annu. Rev. Plant Biol.* **54**, 109–136.
- Levanony, H., and Bashan, Y. (1989). Enhancement of cell division in wheat root tips and growth of root elongation zone induced by *Azospirillum brasilense* Cd. *Can. J. Bot.* **67**, 2213–2216.
- Lin, W., Okon, Y., and Hardy, R. W. F. (1983). Enhanced mineral uptake by *Zea mays* and *Sorghum bicolor* roots inoculated with *Azospirillum brasilense*. *Appl. Environ. Microbiol.* **45**, 1775–1779.
- Liu, K., Fu, H., Bei, Q., and Luan, S. (2000). Inward potassium channel in guard cells as a target for polyamine regulation of stomatal movements. *Plant Physiol.* **124**, 1315–1325.
- Lombardo, M. C., Graziano, M., Polacco, J. C., and Lamattina, L. (2006). Nitric oxide functions as a positive regulator of root hair development. *Plant Signal. Behav.* **1**, 28–33.
- Lucangeli, C., and Bottini, R. (1997). Effects of *Azospirillum* spp. on endogenous gibberellin content and growth of maize (*Zea mays* L.) treated with uniconazole. *Symbiosis* **23**, 63–72.
- Lyubun, Y. V., Fritzsche, A., Chernyshova, M. P., Dudel, E. G., and Fedorov, E. E. (2006). Arsenic transformation by *Azospirillum brasilense* sp245 in association with wheat (*Triticum aestivum* L.) roots. *Plant Soil* **286**, 219–227.
- Malhotra, M., and Srivastava, S. (2006). Targeted engineering of *Azospirillum brasilense* SM with indole acetamide pathway for indoleacetic acid over-expression. *Can. J. Microbiol.* **52**, 1078–1084.

- Malhotra, M., and Srivastava, S. (2008). An *ipdC* gene knock-out of *Azospirillum brasilense* strain SM and its implications on indole-3-acetic acid biosynthesis and plant growth promotion. *Antonie van Leeuwenhoek J. Gen.* **93**, 425–433.
- Malhotra, M., and Srivastava, S. (2009). Stress-responsive indole-3-acetic acid biosynthesis by *Azospirillum brasilense* SM and its ability to modulate plant growth. *Eur. J. Soil Biol.* **45**, 73–80.
- Marchal, K., and Vanderleyden, J. (2000). The “oxygen paradox” of dinitrogen-fixing bacteria. *Biol. Fertil. Soils* **30**, 363–373.
- Marchal, K., Sun, J., Keijers, V., Haaker, H., and Vanderleyden, J. (1998). A cytochrome *cbb3* (cytochrome *c*) terminal oxidase in *Azospirillum brasilense* Sp7 supports microaerobic growth. *J. Bacteriol.* **180**, 5689–5696.
- Mehnaz, S., Weselowski, B., and Lazarovits, G. (2007a). *Azospirillum canadiense* sp. nov., a nitrogen-fixing bacterium isolated from corn rhizosphere. *Int. J. Syst. Evol. Microbiol.* **57**, 620–624.
- Mehnaz, S., Weselowski, B., and Lazarovits, G. (2007b). *Azospirillum zea* sp. nov., a diazotrophic bacterium isolated from rhizosphere soil of *Zea mays*. *Int. J. Syst. Evol. Microbiol.* **57**, 2805–2809.
- Miché, L., Bouillant, M. L., Rohr, R., Sallé, G., and Bally, R. (2000). Physiological and cytological studies on the inhibition of *Striga* seed germination by the plant growth-promoting bacterium *Azospirillum brasilense*. *Eur. J. Plant Pathol.* **106**, 347–351.
- Mirza, M. S., Rasul, G., Mehnaz, S., Ladha, J. K., So, R. B., Ali, S., and Malik, K. A. (2000). Beneficial effects of inoculated nitrogen-fixing bacteria on rice. In “The Quest for Nitrogen Fixation in Rice” (J. K. Ladha and P. M. Reddy, Eds.), pp. 191–204. International Rice Research Institute, Los Banos, Philippines.
- Modolo, L. V., Augusto, O., Almeida, I. M., Magalhaes, J. R., and Salgado, I. (2005). Nitrite as the major source of nitric oxide production by *Arabidopsis thaliana* in response to *Pseudomonas syringae*. *FEBS Lett.* **579**, 3814–3820.
- Molina-Favero, C., Creus, C. M., Lanteri, M. L., Correa-Aragunde, N., Lombardo, M. C., Barassi, C. A., and Lamattina, L. (2007). Nitric oxide and plant growth promoting rhizobacteria: Common features influencing root growth and development. *Adv. Bot. Res.* **46**, 1–33.
- Molina-Favero, C., Creus, C. M., Simontacchi, M., Puntarulo, S., and Lamattina, L. (2008). Aerobic nitric oxide production by *Azospirillum brasilense* Sp245 and its influence on root architecture in tomato. *Mol. Plant Microbe Interact.* **21**, 1001–1009.
- Molla, A. H., Shamsuddin, Z. H., and Saud, H. M. (2001). Mechanism of root growth and promotion of nodulation in vegetable soybean by *Azospirillum brasilense*. *Commun. Soil Sci. Plant Anal.* **32**, 2177–2187.
- Morgenstern, E., and Okon, Y. (1987). Promotion of plant growth and NO_3^- and Rb^+ uptake in *Sorghum bicolor* X *Sorghum sudanense* inoculated with *Azospirillum brasilense*—Cd. *Arid Soil Res. Rehabil.* **1**, 211–217.
- Murty, M. G., and Ladha, J. K. (1988). Influence of *Azospirillum* inoculation on the mineral uptake and growth of rice under hydroponic conditions. *Plant Soil* **108**, 281–285.
- Nemhauser, J. L., Hong, F., and Chory, J. (2006). Different plant hormones regulate similar processes through largely nonoverlapping transcriptional responses. *Cell* **126**, 467–475.
- Niemi, K., Haggman, H., and Sarjala, T. (2001). Effects of exogenous diamines on the interaction between ectomycorrhizal fungi and adventitious root formation in Scots pine in vitro. *Tree Physiol.* **22**, 373–381.
- Nikitina, V. E., Bogomolova, N. V., Ponomareva, E. G., and Sokolov, O. I. (2004). Effect of azospirilla lectins on germination capacity of seeds. *Biol. Bull. (Moscow)* **31**, 354–357.
- Ogut, M., and Er, F. (2006). Micronutrient composition of field-grown dry bean and wheat inoculated with *Azospirillum* and *Trichoderma*. *J. Plant Nutr. Soil Sci.* **169**, 699–703.

- Okon, Y., and Kapulnik, Y. (1986). Development and function of *Azospirillum*—Inoculated roots. *Plant Soil* **90**, 3–16.
- Okon, Y., and Labandera-Gonzalez, C. A. (1994). Agronomic applications of *Azospirillum*: An evaluation of 20 years of worldwide field inoculation. *Soil Biol. Biochem.* **26**, 1591–1601.
- Okon, Y., Heytler, P. G., and Hardy, R. W. F. (1983). N₂ fixation by *Azospirillum brasilense* and its incorporation into host *Setaria italica*. *Appl. Environ. Microbiol.* **46**, 694–697.
- Omay, S. H., Schmidt, W. A., Martin, P., and Bangerth, F. (1993). Indoleacetic acid production by the rhizosphere bacterium *Azospirillum brasilense* Cd under in vitro conditions. *Can. J. Microbiol.* **39**, 187–192.
- Ona, O., Smets, I., Gysegom, P., Bernaerts, K., Impe, J. V., Prinsen, E., and Vanderleyden, J. (2003). The effect of pH on indole-3-acetic acid (IAA) biosynthesis of *Azospirillum brasilense* sp7. *Symbiosis* **35**, 199–208.
- Ona, O., van Impe, J., Prinsen, E., and Vanderleyden, J. (2005). Growth and indole-3-acetic acid biosynthesis of *Azospirillum brasilense* Sp245 is environmentally controlled. *FEMS Microbiol. Lett.* **246**, 125–132.
- Pagnussat, G. C., Simontacchi, M., Puntarulo, S., and Lamattina, L. (2002). Nitric oxide is required for root organogenesis. *Plant Physiol.* **129**, 954–956.
- Pagnussat, G. C., Lanteri, M. L., and Lamattina, L. (2003). Nitric oxide and cyclic GMP are messengers in the indole acetic acid-induced adventitious rooting process. *Plant Physiol.* **132**, 1241–1248.
- Patten, C. L., and Glick, B. R. (1996). Bacterial biosynthesis of indole-3-acetic acid. *Can. J. Microbiol.* **42**, 207–220.
- Peng, G., Wang, H., Zhang, G., Hou, W., Liu, Y., Wang, E. T., and Tan, Z. (2006). *Azospirillum melinis* sp. nov., a group of diazotrophs isolated from tropical molasses grass. *Int. J. Syst. Evol. Microbiol.* **56**, 1263–1271.
- Pereg-Gerk, L., Gilchrist, K., and Kennedy, I. R. (2000). Mutants with enhanced nitrogenase activity in hydroponic *Azospirillum brasilense*-wheat associations. *Appl. Environ. Microbiol.* **66**, 2175–2184.
- Pereyra, M. A., Zalazar, C. A., and Barassi, C. A. (2006). Root phospholipids in *Azospirillum*-inoculated wheat seedlings exposed to water stress. *Plant Physiol. Biochem.* **44**, 873–879.
- Perrig, D., Boiero, M. L., Masciarelli, O. A., Penna, C., Ruiz, O. A., Cassan, F. D., and Luna, M. V. (2007). Plant-growth-promoting compounds produced by two agronomically important strains of *Azospirillum brasilense*, and implications for inoculant formulation. *Appl. Microbiol. Biotechnol.* **75**, 1143–1150.
- Piccoli, P., and Bottini, R. (1994a). Metabolism of 17, 17-[²H₂]-gibberellin A₂₀ to 17, 17-[²H₂]-gibberellin A₁ by *Azospirillum lipoferum* cultures. *AgriScientia Argentina* **11**, 13–15.
- Piccoli, P., and Bottini, R. (1994b). Effects of C/N ratio, N content, pH, and incubation time on growth and gibberellin production by *Azospirillum lipoferum*. *Symbiosis* **17**, 229–236.
- Piccoli, P., Masciarelli, O., and Bottini, R. (1996). Metabolism of 17, 17-[²H₂]-gibberellins A₄, A₉, and A₂₀ by *Azospirillum lipoferum* in chemically-defined culture medium. *Symbiosis* **21**, 263–274.
- Piccoli, P., Lucangeli, C. D., Schneider, G., and Bottini, R. (1997). Hydrolysis of (17, 17-²H₂) gibberellin A₂₀-glucoside and (17, 17-²H₂) gibberellin A₂₀-glucosyl ester by *Azospirillum lipoferum* cultured in a nitrogen-free biotin-based chemically-defined medium. *Plant Growth Regul.* **23**, 179–182.
- Piccoli, P., Masciarelli, O., and Bottini, R. (1999). Gibberellin production by *Azospirillum lipoferum* cultured in chemically defined medium as affected by oxygen availability and water status. *Symbiosis* **27**, 135–145.
- Prigent-Combaret, C., Blaha, D., Pothier, J. F., Vial, L., Poirier, M.-A., Wisniewski-Dyé, F., and Moëgne-Loccoz, Y. (2008). Physical organization and phylogenetic analysis of *acdR* as leucine-responsive regulator of the 1-aminocyclopropane-1-carboxylate deaminase gene *acdS* in phytobeneficial *Azospirillum lipoferum* 4B and other *Proteobacteria*. *FEMS Microbiol. Ecol.* **65**, 202–219.

- Prinsen, E., Costacurta, A., Michiels, K., Vanderleyden, J., and Van Onckelen, H. (1993). *Azospirillum brasilense* indole-3-acetic acid biosynthesis: Evidence for a non-tryptophan dependent pathway. *Mol. Plant Microbe Inter.* **6**, 609–615.
- Puente, M. E., Bashan, Y., Li, C. Y., and Lebsky, V. K. (2004a). Microbial populations and activities in the rhizoplane of rock-weathering desert plants I. Root colonization and weathering of igneous rocks. *Plant Biol.* **6**, 629–642.
- Puente, M. E., Li, C. Y., and Bashan, Y. (2004b). Microbial populations and activities in the rhizoplane of rock-weathering desert plants II. Growth promotion of cactus seedlings. *Plant Biol.* **6**, 643–650.
- Puente, M. E., Rodriguez-Jaramillo, M. C., Li, C. Y., and Bashan, Y. (2006). Image analysis for quantification of bacterial rock weathering. *J. Microbiol. Methods* **64**, 275–286.
- Ramakrishnan, S., Gunasekaran, C. R., and Vadivelu, S. (1997). Effect of bio-fertilizers *Azolla* and *Azospirillum* on root-knot nematode, *Meloidogyne incognita* and plant growth of okra. *Indian J. Nematol.* **26**, 127–130.
- Remans, R., Beebe, S., Blair, M., Manrique, G., Tovar, E., Rao, I., Croonenborghs, A., Torres-Gutierrez, R., El-Howeity, M., Michiels, J., and Vanderleyden, J. (2008). Physiological and genetic analysis of root responsiveness to auxin-producing plant growth-promoting bacteria in common bean (*Phaseolus vulgaris* L.). *Plant Soil* **302**, 149–161.
- Reynders, L., and Vlassak, K. (1979). Conversion of tryptophan to indoleacetic acid by *Azospirillum brasilense*. *Soil Biol. Biochem.* **11**, 547–548.
- Ribaudo, C. M., Krumpholz, E. M., Cassan, F. D., Bottini, R., Cantore, M. L., and Cura, J. A. (2006). *Azospirillum* sp. promotes root hair development in tomato plants through a mechanism that involves ethylene. *J. Plant Growth Regul.* **25**, 175–185.
- Richards, D. E., King, K. E., Ait-ali, T., and Harberd, N. P. (2001). How gibberellin regulates plant growth and development: A molecular genetic analysis of gibberellin signaling. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **52**, 67–88.
- Rodrigues, E. P., Rodrigues, L. S., de Oliveira, A. L. M., Baldani, V. L. D., Teixeira, K. R. D., Urquiaga, S., and Reis, V. M. (2008). *Azospirillum amazonense* inoculation: Effects on growth, yield and N₂-fixation of rice (*Oryza sativa* L.). *Plant Soil* **302**, 249–261.
- Rodriguez, H., Gonzalez, T., Goire, I., and Bashan, Y. (2004). Gluconic acid production and phosphate solubilization by the plant growth-promoting bacterium *Azospirillum* spp. *Naturwissenschaften* **91**, 552–555.
- Rodriguez, H., Mendoza, A., Cruz, M. A., Holguin, G., Glick, B. R., and Bashan, Y. (2006). Pleiotropic physiological effects in the plant growth-promoting bacterium *Azospirillum brasilense* following chromosomal labeling in the *clpX* gene. *FEMS Microbiol. Ecol.* **57**, 217–225.
- Rodriguez-Salazar, J., Suárez, R., Caballero-Mellado, J., and Iturriaga, G. (2009). Trehalose accumulation in *Azospirillum brasilense* improves drought tolerance and biomass in maize plants. *FEMS Microbiol. Lett.* **296**, 52–59.
- Romero, A. M., Correa, O. S., Moccia, S., and Rivas, J. G. (2003). Effect of *Azospirillum*-mediated plant growth promotion on the development of bacterial diseases on fresh-market and cherry tomato. *J. Appl. Microbiol.* **95**, 832–838.
- Ruppel, S., and Merbach, W. (1997). Effect of ammonium and nitrate on ¹⁵N₂-fixation of *Azospirillum* spp., and *Pantoea agglomerans* in association with wheat plants. *Microbiol. Res.* **152**, 377–383.
- Saikia, S. P., Srivastava, G. C., and Jain, V. (2004). Nodule-like structures induced on the roots of maize seedlings by the addition of synthetic auxin 2, 4-D and its effects on growth and yield. *Cereal Res. Commun.* **32**, 83–89.
- Saikia, S. P., Jain, V., Khetarpal, S., and Aravind, S. (2007). Dinitrogen fixation activity of *Azospirillum brasilense* in maize (*Zea mays*). *Curr. Sci.* **93**, 1296–1300.

- Sarig, S., Blum, A., and Okon, Y. (1988). Improvement of the water status and yield of field-grown grain sorghum (*Sorghum bicolor*) by inoculation with *Azospirillum brasilense*. *J. Agric. Sci.* **110**, 271–277.
- Sarig, S., Okon, Y., and Blum, A. (1990). Promotion of leaf area development and yield in Sorghum bicolor inoculated with *Azospirillum brasilense*. *Symbiosis* **9**, 235–245.
- Sarig, S., Okon, Y., and Blum, A. (1992). Effect of *Azospirillum brasilense* inoculation on growth dynamics and hydraulic conductivity of *Sorghum bicolor* roots. *J. Plant. Nutr.* **15**, 805–819.
- Saubidet, M. I., and Barneix, A. J. (1998). Growth stimulation and nitrogen supply to wheat plants inoculated with *Azospirillum brasilense*. *J. Plant Nutr.* **21**, 2565–2577.
- Schmidt, W., Martin, P., Omay, H., and Bangerth, F. (1988). Influence of *Azospirillum brasilense* on nodulation of legumes. In “*Azospirillum* IV, Genetics, Physiology, Ecology” (W. Klingmüller, Ed.), pp. 92–100. Springer Verlag Heidelberg, Germany.
- Schumacher, K. (2006). Endomembrane proton pumps: Connecting membrane and vesicle transport. *Curr. Opin. Plant Biol.* **9**, 595–600.
- Seshadri, S., Muthukumuramasamy, R., Lakshiminarasami, C., and Ignacimuthu, S. (2000). Solubilization of inorganic phosphates by *Azospirillum halopraeferans*. *Curr. Sci.* **79**, 565–567.
- Sgroy, V., Cassán, F., Masciarelli, O., Del Papa, M. F., Lagares, A., and Luna, V. (2009). Isolation and characterization of endophytic plant growth-promoting (PGPB) or stress homeostasis-regulating (PSHB) bacteria associated to the halophyte *Prosopis strombulifera*. *Appl. Microbiol. Biotechnol.* **85**, 371–381.
- Shah, S., Karkhanis, V., and Desai, A. (1992). Isolation and characterization of siderophore, with antimicrobial activity, from *Azospirillum lipoferum*. *Curr. Microbiol.* **25**, 347–351.
- Somers, E., Ptacek, D., Gysegom, P., Srinivasan, M., and Vanderleyden, J. (2005). *Azospirillum brasilense* produces the auxin-like phenylacetic acid by using the key enzyme for indole-3-acetic acid biosynthesis. *Appl. Environ. Microbiol.* **71**, 1803–1810.
- Spaepen, S., Vanderleyden, J., and Remans, R. (2007a). Indole-3-acetic acid in microbial and microorganism–plant signaling. *FEMS Microbiol. Rev.* **31**, 425–448.
- Spaepen, S., Versees, W., Gocke, D., Pohl, M., Steyaert, J., and Vanderleyden, J. (2007b). Characterization of phenylpyruvate decarboxylase, involved in auxin production of *Azospirillum brasilense*. *J. Bacteriol.* **189**, 7626–7633.
- Spaepen, S., Dobbelaere, S., Croonenborghs, A., and Vanderleyden, J. (2008). Effects of *Azospirillum brasilense* indole-3-acetic acid production on inoculated wheat plants. *Plant Soil* **312**, 15–23.
- Sperotto, R. A., Gross, J., Vedoy, C., Passaglia, L. M. P., and Schrank, I. S. (2004). The electron transfer flavoprotein *fixABCX* gene products from *Azospirillum brasilense* show a NifA-dependent promoter regulation. *Curr. Microbiol.* **49**, 267–273.
- Sriskandarajah, S., Kennedy, I. R., Yu, D., and Tchan, Y. T. (1993). Effects of plant growth regulators on acetylene-reducing associations between *Azospirillum brasilense* and wheat. *Plant Soil* **153**, 165–178.
- Steenhoudt, O., and Vanderleyden, J. (2000). *Azospirillum*, a freeliving nitrogen-fixing bacterium closely associated with grasses: Genetic, biochemical and ecological aspects. *FEMS Microbiol. Rev.* **24**, 487–506.
- Steenhoudt, O., Ping, Z., Vande Broek, A., and Vanderleyden, J. (2001). A spontaneous chlorate-resistant mutant of *Azospirillum brasilense* Sp245 displays defects in nitrate reduction and plant root colonization. *Biol. Fertil. Soils* **33**, 317–322.
- Strzelczyk, E., Kampert, M., and Li, C. Y. (1994). Cytokinin-like substances and ethylene production by *Azospirillum* in media with different carbon sources. *Microbiol. Res.* **149**, 55–60.
- Sudhakar, P., Chattopadhyay, G. N., Gangwar, S. K., and Ghosh, J. K. (2000). Effect of foliar application of *Azotobacter*, *Azospirillum* and *Beijerinckia* on leaf yield and quality of mulberry (*Morus alba*). *J. Agric. Sci.* **134**, 227–234.

- Tapia-Hernandez, A., Mascarua-Esparza, M., and Caballero-Mellado, J. (1990). Production of bacteriocins and siderophore-like activity by *Azospirillum brasilense*. *Microbios* **64**, 73–83.
- Tchan, Y. T., Zeman, A. M. M., and Kennedy, I. R. (1991). Nitrogen fixation in paradonules of wheat roots by introduced free-living diazotrophs. *Plant Soil* **137**, 43–47.
- Tewari, R. K., Hahn, E. J., and Paek, K. Y. (2007). Function of nitric oxide and superoxide anion in the adventitious root development and antioxidant defence in *Panax ginseng*. *Plant Cell Rep.* **27**, 563–573.
- Thuler, D. S., Floh, E. I. S., Handro, W., and Barbosa, H. R. (2003). Plant growth regulators and amino acids released by *Azospirillum* sp in chemically defined media. *Let. Appl. Microbiol.* **37**, 174–178.
- Tien, T. M., Gaskins, M. H., and Hubell, D. H. (1979). Plant growth substances produced by *Azospirillum brasilense* and their effect on the growth of pearl millet (*Pennisetum americanum* L). *Appl. Environ. Microbiol.* **37**, 1016–1024.
- Tilak, K. V. B. R., Ranganayaki, N., Pal, K. K., De, R., Saxena, A. K., Nautiyal, C. S., Mittal, S., Tripathi, A. K., and Johri, B. N. (2005). Diversity of plant growth and soil health supporting bacteria. *Curr. Sci.* **89**, 136–150.
- Tsagou, V., Kefalogianni, I., Sini, K., and Aggeli, G. (2003). Metabolic activities in *Azospirillum lipoferum* grown in the presence of NH_4^+ . *Appl. Microbiol. Biotechnol.* **62**, 574–578.
- Tsavelkova, E. A., Klimova, S. Y., Cherdyntseva, T. A., and Netrusov, A. I. (2006). Microbial producers of plant growth stimulators and their practical use: A review. *Appl. Biochem. Microbiol.* **42**, 117–126.
- Van Dommelen, A., Croonenborghs, A., Spaepen, S., and Vanderleyden, J. (2009). Wheat growth promotion through inoculation with an ammonium-excreting mutant of *Azospirillum brasilense*. *Biol. Fertil. Soils* **45**, 549–553.
- Vande Brock, A., and Vanderleyden, J. (1995). Genetics of the *Azospirillum*–plant root association–review. *Crit. Rev. Plant Sci.* **14**, 445–466.
- Vande Broek, A., Lambrecht, M., Eggermont, K., and Vanderleyden, J. (1999). Auxins upregulate expression of the indole-3-pyruvate decarboxylase gene in *Azospirillum brasilense*. *J. Bacteriol.* **181**, 1338–1342.
- Wood, C. C., Islam, N., Ritchie, R. J., and Kennedy, I. R. (2001). A simplified model for assessing critical parameters during associative $^{15}\text{N}_2$ fixation between *Azospirillum* and wheat. *Aust. J. Plant Physiol.* **28**, 969–974.
- Yahalom, E., Okon, Y., and Dovrat, A. (1990). Possible mode of action of *Azospirillum brasilense* strain Cd on the root morphology and nodule formation in burr medic (*Medicago polymorpha*). *Can. J. Microbiol.* **36**, 10–14.
- Yasuda, M., Isawa, T., Shinozaki, S., Minamisawa, K., and Nakashita, H. (2009). Effects of colonization of a bacterial endophyte, *Azospirillum* sp. B510, on disease resistance in rice. *Biosci. Biotechnol. Biochem.* **73**, 2595–2599.
- Yu, D., Kennedy, I. R., and Tchan, Y. T. (1993). Verification of nitrogenase activity (C_2H_2 reduction) in *Azospirillum* populated, 2, 4-dichlorophenoxyacetic acid induced, root structures of wheat. *Aust. J. Plant. Physiol.* **20**, 187–195.
- Zakharova, E. A., Shcherbakov, A. A., Brudnik, V. V., Skripko, N. G., Bulkhin, N. S., and Ignatov, V. V. (1999). Biosynthesis of indole-3-acetic acid in *Azospirillum brasilense*: Insights from quantum chemistry. *Eur. J. Biochem.* **259**, 572–576.
- Zakharova, E. A., Iosipenko, A. D., and Ignatov, V. V. (2000). Effect of water-soluble vitamins on the production of indole-3-acetic acid by *Azospirillum brasilense*. *Microbiol. Res.* **155**, 209–214.
- Zeevaart, J. A. D. (1999). Abscisic acid metabolism and its regulation. In “Biochemistry and Molecular Biology of Plant Hormones” (P. P. J. Hooykaas, M. A. Hall, and K. R. Libbenga, Eds.), pp. 189–207. Elsevier Science, Amsterdam.

- Zeidler, D., Zahringer, U., Gerber, I., Dubery, I., Hartung, T., Bors, W., Hutzler, P., and Durner, J. (2004). Innate immunity in *Arabidopsis thaliana*: Lipopolysaccharides activate nitric oxide synthase (NOS) and induce defense genes. *Proc. Natl. Acad. Sci. USA* **101**, 15811–15816.
- Zeman, A. M. M., Tchan, Y. T., Elmerich, C., and Kennedy, I. R. (1992). Nitrogenase activity in wheat seedlings bearing paranodules induced by 2, 4-dichlorophenoxyacetic acid (2, 4-D) and inoculated with *Azospirillum*. *Res. Microbiol.* **143**, 847–855.
- Zimmer, W., Penteadó-Stephan, M., and Bothe, H. (1984). Denitrification by *Azospirillum brasilense* Sp 7. *Arch. Microbiol.* **138**, 206–211.
- Zimmer, W., Roeben, K., and Bothe, H. (1988). An alternative explanation for plant growth promotion by bacteria of the genus *Azospirillum*. *Planta* **176**, 333–342.
- Zimmer, W., Aparicio, C., and Elmerich, C. (1991). Relationship between tryptophan biosynthesis and indole-3-acetic acid production in *Azospirillum*: Identification and sequencing of *trpGDC* cluster. *Mol. Gen. Genet.* **229**, 41–51.
- Zimmer, W., Wesche, M., and Timmermans, L. (1998). Identification and isolation of the indole-3-pyruvic decarboxylase gene from *Azospirillum brasilense* Sp7: Sequencing and functional analysis of the gene locus. *Curr. Microbiol.* **36**, 327–331.