



NEWS AND VIEWS

PROPOSAL FOR THE DIVISION OF PLANT GROWTH- PROMOTING RHIZOBACTERIA INTO TWO CLASSIFICATIONS: BIOCONTROL-PGPB (PLANT GROWTH-PROMOTING BACTERIA) AND PGPB

YOAV BASHAN* and GINA HOLGUIN

Department of Microbiology, The Center for Biological Research of the Northwest (CIB), P.O. Box
128, La Paz, B.C.S., 230(X) Mexico

(Accepted 6 July 1997)

Bacteria that can increase plant growth and productivity have been known for over a century (Brown, 1974). Prominent among these organisms are species of the genus *Rhizobium* (Dejordjevic *et al.*, 1987) whose potential and practical use in agriculture is beyond doubt (O'Gara *et al.*, 1995). Although other bacterial species associated with plants, but, without creating symbiosis like *Rhizobium*, were discovered early in this century (Beijerinck, 1925), the fact that many of them were able to promote plant growth was not widely recognized. The breakthrough in this field occurred in the mid 1970s with (i) the discovery that some bacteria, mainly pseudomonads, are capable of controlling soil-borne pathogens to indirectly enhance plant growth (Kloepper and Schroth, 1978, Kloepper *et al.*, 1980), and (ii) the re-discovery (Döbereiner and Day, 1976) of *Azospirillum spp.* (Beijerinck, 1925) which directly affect plant metabolism and consequent growth. These findings established two research approaches to bacterial inoculation (i) the biocontrol of plant pathogens (Kloepper *et al.*, 1989), and (ii) *Azospirillum*-related organisms (Bashan and Levanony, 1990). Most bacteria in both groups are root associated, free living in soil or the phyllosphere and unlike *Rhizobium* or *Agrobacterium* form no apparent structural association with plant roots (Kloepper *et al.*, 1989; Bashan and Levanony, 1990). Curiously, the two fields have developed almost independently of one another, yielding only few cross-citations (Glick, 1995; Institute for Scientific Information (Philadelphia) citation search, 1996 done by the

authors) in papers with subjects in common such as migration towards the plant, survival in soil and effects on plant growth. The primary barriers between these two fields have been of both taxonomic and functional nature.

When Kloepper and Schroth (1978) and Kloepper *et al.* (1980) coined the term Plant Growth-Promoting *Rhizobacteria* (PGPR), it was originally used to describe the biocontrol group. This term is used appropriately by many scientists, and is almost universally accepted to describe this research field. The other bacterial group has no particular name and is vaguely referred to by different terms depending on the research group.

During the last decade, close to 4000 publications have appeared in the field of Plant Growth Promoting Bacteria (Life Science data base; Cambridge Scientific abstracts). Many of these studies characterized new bacterial strains which had not belonged to any previously defined group, thus contributing to the confusion in classification and terminology. The numerous terms currently being used to describe plant growth-enhancing bacteria (Table 1 and Appendix) foster incorrect and confusing communication between scientists and unnecessary hardship in extracting data from databases. We believe that an accurate, accepted terminology is fundamental for biological precision, scientific cooperation, and understanding between various researchers.

We propose two new terms for general scientific use; "biocontrol plant growth-promoting bacteria" (Biocontrol-PGPB) and "plant growth-promoting bacteria" (PGPB). These terms would seem to encompass all the plant beneficial bacteria according to their particular role as understood today.

*Author for correspondence.

Table 1. Currently used terms for Plant Growth-Promoting Bacteria (PGPB) and their limitations

Term	Where used?	Major drawbacks	Selected reference
PGPR (plant growth-promoting rhizobacteria; rhizosphere bacteria)	This term covers all bacteria of rhizosphere origin that promote plant growth. It has been used solely in the field of bacterial biocontrol. These bacteria suppress disease development by either producing pathogen-inhibiting substances or by increasing plant resistance.	(1) The term does not cover non-rhizosphere interactions like biocontrol in the phyllosphere. (2) This term was coined when this field was relatively small (in number of publications). At that time, its broad reference was an advantage. Today, the term is too general and non specific.	Brown <i>et al.</i> , 1990; Keel <i>et al.</i> , 1991; Casida and Lukezic, 1992; Bashan <i>et al.</i> , 1993; Chanway and Holl, 1994a; Tuzun and Kloepper, 1994
Plant-beneficial bacteria (PBB)	This term covers all, experimentally proven, bacteria exhibiting positive effects on plants. The common term for non-biocontrol type of bacteria.	(1) This term does not appropriately differentiate between bacteria that are beneficial to plants (i.e., increase plant growth) and bacteria that may have one beneficial characteristic (e.g. nitrogen fixation, phosphate solubilization or plant-hormone production) among other non-beneficial characteristics. For example, can the nitrogen-fixing, tumor-producing <i>Agrobacterium tumefaciens</i> be considered beneficial bacteria? (2) Term is too broad and does not precisely delineate the nature of the bacteria.	Kanvinde and Sastry, 1990; Bashan and Holguin, 1994; Chanway and Holl, 1994b.
Direct and Indirect PGPB	"Indirect" refers to biocontrol PGPB and "direct" refers to all other PGPB. While these two classifiers distinguish the nature of bacterial effects, they are rarely found in the literature.	The term "direct" excludes non-biocontrol bacteria that indirectly affect plant growth by their metabolic products like phosphate solubilization, sulfur oxidation, nitrogen fixation by cyanobacteria or ethylene production.	Gaind and Gaur, 1991; Grayston and Germida, 1991; Glick <i>et al.</i> , 1994; Ryder <i>et al.</i> , 1994; Toledo <i>et al.</i> , 1995

A classification scheme for plant enhancing bacteria should: (i) accurately describe its primary nature: capacity to increase plant growth or yield. (ii) cover as many of the bacteria as possible that are known at the time. (iii) allow for accumulating data afterwards with only minor modification, (iv) be clear and encompassing, (v) describe each group of bacteria by their unique features and not by their lack of features, and (vi) if more than one term is used, they should not be contradictory.

We therefore propose that the common descriptor Plant Growth-Promoting Rhizobacteria (Kloepper *et al.*, 1980) be updated to accommodate new knowledge in the field. We believe that this term best describes the nature of many beneficial bacteria and is already widely accepted by the scientific community. Since many beneficial bacteria are not rhizosphere bacteria, we propose replacing "rhizobacteria" with "bacteria", creating the modified term Plant Growth-promoting Bacteria (PGPB). When this term is used to describe bacteria that suppress a plant pathogen (by either producing inhibitory substances or by increasing the natural resistance of the plant), the word "Biocontrol" will combine to produce the new term Biocontrol PGPB. The term PGPB is applicable where the bacteria affect plants by means other than the suppression of other microorganisms. They can do this through their own metabolism [phosphate solubilization, hormone production, N₂-fixation (Dejordjevic *et al.*, 1987; Leinhos, 1994; Strzelczyk *et al.*, 1994), directly affect the plant metabolism [enhance

water and mineral uptake (Bashan *et al.*, 1990), improve root development (Okon and Kapulnik, 1986), enhance plant enzyme activity (Ferreira *et al.*, 1987)] or affect the plant by "helping" another beneficial microorganism to function better [e.g. *Azospirillum* increasing modulation of legumes by rhizobia (Yahalom *et al.*, 1987), or enhancing mycorrhizal phosphate solubilization (Barea *et al.*, 1983) or mycorrhizal infection (Li *et al.*, 1992; Garbaye, 1994)].

These terms could be used regardless of the bacterial genus or species. However, in some cases different strains of the same species might be biocontrol PGPB or PGPB. In rare cases, these different functional features could be found in the same strain (Oberhänsli *et al.*, 1991). The new terms proposed include, to the best of our knowledge, all the plant growth-enhancing bacteria known today, including rhizobia. Adoption of these terms would allow scientists to communicate more accurately on a global basis, and will standardize this field of biology. At the same time, the old terms should be abandoned.

Acknowledgements---This essay was written in memory of the late Mr Avner Bashan from Israel. It was supported by Consejo Nacional de Ciencia y Tecnologia contract # 26262-B (CONACyT), Mexico. We thank Dr Bernard Click, University of Waterloo, Canada. Dr Ching Li, USDA Oregon, U.S.A.. Dr Chris Chanway, University of British Columbia, Canada for many helpful and encouraging comments during manuscript preparation, the three anonymous referees for constructive comments, Dr Roy Bowers and Dr Ellis Glazier for crystallizing the English and Mr Joseph Gabon, Ohio State University Library for archival computer search.

REFERENCES

- Barea J. M., Bonis A. F. and Olivares A. (1983) Interactions between *Azospirillum* and VA mycorrhiza and their effects on growth and nutrition of maize and ryegrass. *Soil Biology & Biochemistry* **15**, 705-709.
- Bashan Y. (1993) Potential use of *Azospirillum* as biofertilizer. *Turrialba* **43**, 286-291.
- Bashan Y., Harrison S. K. and Whitmoyer R. E. (1990) Enhanced growth of wheat and soybean plants inoculated with *Azospirillum brasilense* is not necessarily due to general enhancement of mineral uptake. *Applied and Environmental Microbiology* **56**, 769-775.
- Bashan Y. and Holguin G. (1994) Root-to-root travel of the beneficial bacterium *Azospirillum brasilense*. *Applied and Environmental Microbiology* **60**, 2120-2131.
- Bashan Y. (1995) *Azospirillum* and other non-biocontrol PGPR: do they have a place in the agricultural future of developed countries? Key-note lecture in *Biotechnology of Rhizosphere Enhancing Microorganisms*. Vancouver, Canada.
- Bashan Y., Holguin G. and Lifshitz R. (1993) Isolation and characterization of plant growth-promoting rhizobacteria. In *Methods in Plant Molecular Biology and Biotechnology*, eds B. R. Glick and J. E. Thompson, pp. 331-345. CRC Press, USA.
- Bashan Y. and Levanony H. (1990) Current status of *Azospirillum* inoculation technology: *Azospirillum* as a challenge for agriculture. *Canadian Journal of Microbiology* **36**, 591-608.
- Bazin M. J., Markham P., Scott E. M. and Lynch J. M. (1990) Population dynamics and rhizosphere interactions. In *The Rhizosphere*, ed. J. M. Lynch, pp. 991-1037. John Wiley, Chichester.
- Beijerinck M. W. (1925) Über ein *Spirillum* welches freien Stickstoff binden kann? *Zentralblatt für Bakteriologie, Parasitenkunde, Infektionskrankheiten und Hygiene, Naturwissenschaftliche Abteilung* **2**, 63, 353-359.
- Brown G., Khan Z. and Lifshitz R. (1990) Plant growth promoting rhizobacteria: strain identification by restriction fragment length polymorphism. *Canadian Journal of Microbiology* **36**, 242-248.
- Brown M. E. (1974) Seed and root bacterization. *Annual Review of Phytopathology* **12**, 181-197.
- Casida L. E. Jr. and Lukezic F. L. (1992) Control of leaf spot diseases of alfalfa and tomato with applications of the bacterial predator *Pseudomonas* strain 679-2. *Plant Disease* **76**, 1217-1220.
- Chanway C. P. and Holt F. B. (1994a) Growth of outplanted lodgepole pine seedlings one year after inoculation with plant growth-promoting rhizobacteria. *Forest Science* **40**, 238-246.
- Chanway C. P. and Holt F. B. (1994b) Ecological growth response specificity of two Douglas-fir ecotypes inoculated with coexistent beneficial rhizosphere bacteria. *Canadian Journal of Botany* **72**, 582-586.
- de Freitas J. R. and Germida J. J. (1990) A root tissue culture system to study winter wheat-rhizobacteria interactions. *Applied Microbiology and Biotechnology* **33**, 589-595.
- Dejordjevic M. A., Gabriel D. W. and Rolfe B. G. (1987) *Rhizobium*-the refined parasite of legumes. *Annual Review of Phytopathology* **25**, 145-168.
- Del Gallo M. and Fendrik I. (ed) (1992) *Azospirillum and Related Microorganisms V. Genetics, Physiology, Ecology*. Balaban Publishers, Philadelphia.
- Döbereiner J. and Day J. M. (1976) Associative symbioses in tropical grasses: characterization of microorganisms and dinitrogen fixing sites. In *Proceedings of the First International Symposium on Nitrogen Fixation*, eds W. E. Newton and C. J. Nyman, Vol. 2, pp. 518-538. Washington State University Press, Pullmann.
- Dupler M. and Baker R. (1984) Survival of *Pseudomonas pulida*, a biological control agent, in soil. *Phytopathology* **74**, 195-200.
- Fendrik L., Del Gallo M., Vanderleyden J. and de Zamaroczy M. (ed) (1995) *Azospirillum VI and Related Microorganisms, Genetics-Physiology-Ecology*. NATO Advance Science Institutes Series. Vol. G37. Springer-Verlag, Berlin.
- Ferreira M. C. B., Fernandes M. S. and Döbereiner J. (1987) Role of *Azospirillum brasilense* nitrate reductase in nitrate assimilation by wheat plants. *Biology and Fertility of Soils* **4**, 47-53.
- Gaind S. and Gaur A. C. (1991) Thermotolerant phosphate solubilizing microorganisms and their interaction with mung bean. *Plant and Soil* **133**, 141-149.
- Garbaye J. (1994) Helper bacteria: a new dimension to the mycorrhizal symbiosis. *New Phytologist* **128**, 197-210.
- Glick B. R. (1995) The enhancement of plant growth by free-living bacteria. *Canadian Journal of Microbiology* **41**, 109-117.
- Glick B. R., Jacobson C. B., Schwarze M. M. K. and Pasternak J. J. (1994) 1-aminocyclopropane-1-carboxylic acid deaminase mutants of the plant growth-promoting rhizobacterium *Pseudomonas putida* GR12-2 do not stimulate canola root elongation. *Canadian Journal of Microbiology* **40**, 911-915.
- Grayston S. J. and Germida J. J. (1991) Sulfur-oxidizing bacteria as plant growth promoting rhizobacteria for canola. *Canadian Journal of Microbiology* **37**, 521-529.
- James E. K., Reis V. M., Olivares F. L., Baldani J. I. and Döbereiner J. (1994) Infection of sugar cane by the nitrogen-fixing bacterium *Acetohacter diazotrophicus*. *Journal of Experimental Botany* **45**, 757-766.
- Kanvinde U. and Sastry G. R. R. (1990) *Agrobacterium tumefaciens* is a diazotrophic bacterium. *Applied and Environmental Microbiology* **56**, 2087-2092.
- Keel C., Koller B. and Défago G. (ed) (1991) *Plant Growth-Promoting Rhizobacteria-Progress and Prospects*. International Organization for Biological and Integrated Control of Noxious Animals and Plants, Bulletin 1991' XIV/8, Zurich, Switzerland.
- Kloepper J. W., Lifshitz R. and Zablutowicz R. M. (1989) Free-living bacterial inocula for enhancing crop productivity. *Trends in Biotechnology* **7**, 39-43.
- Kloepper J. W., Leong J., Teintze M. and Schroth M. N. (1980) Enhanced plant growth by siderophores produced by plant growth-promoting rhizobacteria. *Nature* **286**, 885-886.
- Kloepper J. W., Schippers B. and Bakker P. A. H. M. (1992) Proposed elimination of the term endorhizosphere. *Phytopathology* **82**, 726-727.
- Kloepper J. W. and Schroth M. N. (1978) Plant growth promoting rhizobacteria on radishes. In *Proceedings of the 4th International Conference on Plant Pathogenic Bacteria*, ed. Station de Pathologie Végétale et Phytobactériologie. Vol. 2, pp. 879-882. Angers, France.
- Ladha J. K. and So R. B. (1994) Numerical taxonomy of photosynthetic rhizobia nodulating *Aeschynomene* species. *International Journal of Systematic Bacteriology* **44**, 62-73.
- Leinhos V. (1994) Effects of pH and glucose on auxin production of phosphate-solubilizing rhizobacteria in vitro. *Microbiological Research* **149**, 135-138.
- Li C. Y., Massicote H. B. and Moore L. V. H. (1992) Nitrogen-fixing *Bacillus* sp. associated with Douglas-fir tuberculate ectomycorrhizae. *Plant and Soil* **140**, 35-40.
- Lindow S. E. (1987) Competitive exclusion of epiphytic bacteria by *Ice* *Pseudomonas syringae* mutants. *Applied and Environmental Microbiology* **53**, 2520-2527.
- Lynch J. M. (1982) Interactions between bacteria and plants in the root environment. In *Bacteria and Plants*,

- gds M. E. Rhodes-Roberts and F. A. Skinner, pp. 1-23. Academic Press, New York.
- McInroy J. A. and Kloepper J. W. (1991) Analysis of population densities and identification of endophytic bacteria of maize and cotton in the field. In *Plant Growth-Promoting Rhizobacteria-Progress and Prospects*, gds C. Keel, B. Koller and G. Défago, pp. 328-331. International Organization for Biological and Integrated Control of Noxious Animals and Plants, Bulletin 1991 J XIV/8, Zürich, Switzerland.
- Misaghi I. J. and Donndelinger C. R. (1990) Endophytic bacteria in symptom-free cotton plants. *Phytopathology* **80**, 808-811.
- Oberhänsli T., Défago G. and Haas D. (1991) Indole-3-acetic acid (IAA) synthesis in the biocontrol strain CHAD of *Pseudomonas-fluorescens*: role of tryptophan side chain oxidase. *Journal of General Microbiology* **137**, 2273-2279.
- O'Gara F., Moenne-Loccoz Y., Dowling D. N. and Nuti M. P. (1995) Recent developments in biotechnology and the potential impact of nitrogen-fixation in the field. In *Nitrogen Fixation: Fundamentals and Applications*, gds I. A. Tikhonovich, N. A. Provorov, V. I. Romanov, W. E. Newton, pp. 623-628. Kluwer Academic Publishers, Dordrecht.
- Okon Y. and Kapulnik Y. (1986) Development and function of *Azospirillum*-inoculated roots. *Plant and Soil* **90**, 3-16.
- Puente M.-E. and Bashan Y. (1993) Effect of inoculation with *Azospirillum brasilense* strains on the germination and seedlings growth of the giant columnar cactus (*Pachycereus pringlei*). *Symbiosis* **15**, 49-60.
- Ryder M. H., Stephens P. M. and Bowen G. D. (1994). Preface. In *Improving Plant Productivity with Rhizosphere Bacteria*. CSIRO Division of Soils, Glen Osmond.
- Strzelczyk E., Kampert M. and Li C. Y. (1994) Cytokinin-like substances and ethylene production by *Azospirillum* in media with different carbon sources. *Microbiological Research* **149**, 55-60.
- Tang W.-H. (1994) Yield-increasing bacteria (YIB) and biocontrol of sheath blight of rice. In *Improving Plant Productivity with Rhizosphere Bacteria*, gds M. H. Ryder, P. M. Stephens and G. D. Bowen, pp. 267-273. CSIRO Division of Soils, Glen Osmond.
- Toledo G., Bashan Y. and Soeldner A. (1995) Cyanobacteria and black mangroves in Northwestern Mexico: colonization, and diurnal and seasonal nitrogen fixation on aerial roots. *Canadian Journal of Microbiology* **41**, 999-1011.
- Tuzun S. and Kloepper J. (1994) Induced systemic resistance by plant growth-promoting rhizobacteria. In *Improving Plant Productivity with Rhizosphere Bacteria*, gds M. H. Ryder, P. M. Stephens and G. D. Bowen, pp. 104-109. CSIRO Division of Soils, Glen Osmond.
- van Peer R., Punte H. L. M., de Weger L. A. and Schippers B. (1990) Characterization of root surface and endorhizosphere pseudomonads in relation to their colonization on roots. *Applied and Environmental Microbiology* **56**, 2462-2470.
- Yahalom E., Okon Y. and Dovrat A. (1987) *Azospirillum* effects on susceptibility to *Rhizobium* nodulation and on nitrogen fixation of several forage legumes. *Canadian Journal of Microbiology* **33**, 510-514.
- You C. and Zhou F. (1989) Non-nodular endorhizospheric nitrogen fixation in wetland rice. *Canadian Journal of Microbiology* **35**, 403-408.

APPENDIX

Terms In Use Describing Habitat

Root associative bacteria. The original term for the *Azospirillum*-like organisms. This term covers all bacteria which live in the rhizosphere and rhizoplane. Major drawbacks: (1) Not appropriate for endophytic organisms such as *Acetobacter diazotrophicus*, rhizobia living in stem and leave nodules, phyllosphere bacteria and many other endophytic bacteria whether root residents or introduced biocontrol agents. (2) The term is too general to be meaningfully related to the nature of the bacteria. (Döbereiner and Day, 1976; Lindow, 1987; Misaghi and Donndelinger, 1990; McInroy and Kloepper, 1991; James *et al.*, 1994; Ladha and So, 1994).

Rhizosphere bacteria (rhizobacteria). Like the term "root associative bacteria", it covers all bacteria living in the rhizosphere. These include numerous species and strains, either pathogenic or saprophytic. Major drawback: the term refers to the location from which the bacteria were isolated rather than the effect these bacteria have on plants (Bazin *et al.*, 1990; de Freitas and Germida, 1990; Chanway and Holl, 1994b).

Endorhizosphere bacteria. This term is supposed to describe bacteria living in the "inner" parts of the rhizosphere (Lynch, 1982; You and Zhou, 1989; van Peer *et al.*, 1990). Major drawback: a semantically incorrect term that should not be used for the arguments presented by Kloepper *et al.* (1992).

Special Use Terms (Sub-division Of PGPB)

Biofertilizer. The term is favored by the inoculation industry. It is believed that the term "biofertilizer" facilitates the registration process of new bacterial products in industrialized countries. In developing countries, it is also easier to sell a product with a name that reminds the farmer of "fertilizer" rather than "bacteria", which connotes human and animal diseases. Major drawback: Despite its popularity, this term is inappropriate because it excludes all bacterial strains which do not contribute to plant growth via mineral uptake and microbial extraction of minerals from rocks like Biocontrol-PGPB (Bashan, 1993).

Yield increasing bacteria. The term was originated in China and is exclusively being used there. Major drawback: the term is incomplete. It wrongly assumes that all PGPB increase plant yield. This is not true in plants which have no defined yield like cactus. There, the entire plant simply grows more vigorously (Puente and Bashan, 1993; Tang, 1994).

***Azospirillum*-related microorganisms.** Currently in use in the non-biocontrol field. Major drawbacks: it has a limited scope (referring only to *Azospirillum*-type bacteria) and does not properly define which are the "related microorganisms" and by what criteria a bacterium should be considered "*Azospirillum*-related" (Del Gallo and Fendrik, 1992; Fendrik *et al.*, 1995).

Non-biocontrol PGPR. The term has been used to define non-biocontrol PGPB. Major drawback: it is inappropriate to define an organism by what it does not do rather than by what it does (Bashan, 1995)

Other Used Term

Biocontrol Agent. Used solely in the biocontrol field. Major drawback: the term does not differentiate between bacteria, fungi, and other organisms that have biological control capacity over pathogens and thus, it is non-specific (Dupler and Baker, 1984).