

ABSENCE OF A CORRELATION BETWEEN NITRATE REDUCTASE AND SYMBIOTIC NITROGEN FIXATION EFFICIENCY IN RHIZOBIUM MELILOTI

H. ANTOUN¹, L. M. BORDELEAU¹, D. PRÉVOST¹,
and R. A. LACHANCE²

¹Research Station, Agriculture Canada, 2560 Hochelaga Boulevard, Ste-Foy, Quebec G1V 2J6; and ²Department of Phytology, Faculty of Agriculture and Food Science, Laval University, Que. G1K 7P4. Contribution no. 144, received 30 May 1979, accepted 31 Aug. 1979.

ANTOUN, H., BORDELEAU, L. M., PRÉVOST, D. AND LACHANCE, R. A. 1980. Absence of a correlation between nitrate reductase and symbiotic nitrogen fixation efficiency in *Rhizobium meliloti*. *Can. J. Plant Sci.* **60**: 209-212.

Specific activities of the assimilatory and "regulatory" types of nitrate reductase were studied in 41 strains of *Rhizobium meliloti* having different symbiotic nitrogen fixation activities. Both nitrate reductase enzymes were present in very effective and ineffective strains and no significant correlation was found between the specific activities of the two enzymes and the dry matter yields of alfalfa obtained with the 41 strains. Measurements of the specific activities of the two nitrate reductase enzymes in the vegetative bacteria cannot be used as a rapid physiological test for the selection of very effective strains of *R. meliloti*.

Les activités spécifiques des nitrates réductases assimilatoires et "régulateurs" ont été étudiées chez 41 souches de *Rhizobium meliloti* ayant des activités différentes de fixation symbiotique d'azote. Les deux nitrates réductases ont été retrouvées chez les souches très efficaces ainsi que chez les souches inefficaces et aucune corrélation significative n'a été trouvée entre les activités spécifiques et le rendement de la matière sèche de la luzerne obtenu avec les 41 souches à l'essai. La détermination des activités spécifiques de ces deux nitrates réductases chez les bactéries végétatives ne peut donc pas être utilisée comme un test physiologique rapide pour la sélection de souches très efficaces de *R. meliloti*.

The presence of a significant correlation between nitrate reductase and nitrogenase enzyme activities was suggested in *Rhizobium japonicum* (Cheniae and Evans 1960) and *Rhizobium meliloti* (Kondorosi et al. 1973; Sik et al. 1976). Evans and Russel (1971) also suggested that the two enzymes share a common structural protein component. A mutation in the suggested common component should have a pleiotropic effect on both nitrate reductase and nitrogenase. By using three ineffective mutants of *Rhizobium meliloti*, Sik and Barabás (1977) showed that the assimilatory type of nitrate reductase (produced when nitrate is the only

nitrogen source in the culture medium) could be induced in the effective wild-type bacteria as well as in its ineffective mutants, and they concluded that the connection between the two reducing enzymes is not of a structural nature. However, a special type of nitrate reductase labelled "regulatory" was detected only in the effective wild-type bacteria, cultured under aeration on amino acids both as nitrogen and carbon sources. A possible regulatory connection between the "regulatory" type of nitrate reductase and nitrogenase was proposed (Barabás and Sik 1979; Sik and Barabás 1977). If this hypothesis, based on observations made with a few mutants, could be applied to the

Can. J. Plant Sci. **60**: 209-212 (Jan. 1980)

symbiotic nitrogen fixation systems of a relatively larger population of *R. meliloti*, then a positive significant correlation is to be expected between the specific activity of the "regulatory" nitrate reductase and the symbiotic nitrogen fixation efficiency, and a rapid physiological test for the selection of very effective strains will be established. Because the selection of very effective strains of *Rhizobium* is a time-consuming process (Bordeleau et al. 1977), there is an increasing need for rapid and practical methods of strain selection (Bordeleau and Antoun 1978).

In the present work, the specific activities of the assimilatory and "regulatory" types of nitrate reductase were studied in 41 strains of *R. meliloti* having different symbiotic nitrogen fixation activities.

MATERIALS AND METHODS

Bacteria

The 41 strains of *Rhizobium meliloti* used in this study and their symbiotic nitrogen fixation efficiency were described in a previous report (Bordeleau et al. 1977).

Assimilatory Nitrate Reductase

This enzyme is formed in bacteria growing with nitrate as the only nitrogen source (Sik and Barabás 1977). Bacteria were cultured, at 26.6°C during 7 days on a rotary shaker, in a liquid medium containing per liter of distilled water: mannitol, 10.0 g; KNO₃, 1.0 g; NaCl, 0.1 g; K₂HPO₄, 0.5 g; MgSO₄ · 7H₂O, 0.2 g; yeast extract (Difco), 0.1 g; pH, 7.0. Bacterial cells from 10 mL of the suspension were washed in saline and digested at 100°C in 1 mL of 1 N NaOH for 10–20 min. After neutralization with 1 N HCl, the protein content was measured by the Bio-Rad protein assay (Bio-Rad Laboratories, Richmond, Calif.) based on the method of Bradford (1976). Residual nitrate in the liquid supernatant was measured by the method of Cataldo et al. (1975) and the nitrite formed was determined as described by Bremner (1965). The specific enzyme activity was expressed as the amount of nitrate reduced or nitrite detected per microgram of protein.

"Regulatory" Nitrate Reductase

This type of enzyme is detectable in bacteria

grown on amino acids both as nitrogen and carbon sources, without added nitrate, and it was suggested that it is induced by endogenous formation of nitrate (Sik and Barabás 1977). Bacteria were cultured for 48 h at 26.6°C on a rotary shaker, in YTB medium (Orosz and Sik 1970) which contained per liter of distilled water: tryptone (Difco), 10.0 g; yeast extract (Difco), 1.0 g; NaCl, 5.0 g; CaCl₂ · 2H₂O, 0.15 g; MgSO₄ · 7H₂O, 0.25 g; pH, 7.0. Bacterial cells were washed in saline and resuspended in fresh YTB medium and incubated for 2.5 h at 26.6°C on a rotary shaker. Cells from 5 mL of this suspension were collected and resuspended in a buffered saline solution (NaCl, 7.0 g; KH₂PO₄, 0.34 g; K₂HPO₄, 1.21 g; pH, 7.2). To 1-mL samples, 10 mmoles nitrate was added and after incubation for 10 or 20 min at 30°C without aeration, nitrite was determined as described by Nicholas and Nason (1957). The specific enzyme activity was expressed as the amount of nitrite produced per microgram of protein per minute.

RESULTS AND DISCUSSION

All the tested strains of *R. meliloti* reduced nitrate except the effective strain 3DOa2oa. This strain failed to utilize nitrate and showed slight growth on yeast extract added as vitamin source in the culture medium (Table 1). The highest nitrate reduction was performed by the effective strain 23A. The nitrite detected in the culture medium measures the nitrate reductase activity but also reflects the nitrite reductase activity. In fact, one might expect to detect more nitrite if a strain has a high nitrate reductase activity and/or a low nitrite reductase activity. The very effective strain A₃ reduced nitrate but no nitrite was detected, indicating a high activity of nitrite reductase. The highest amount of nitrite was detected with the effective strain A₅, indicating a high activity of nitrate reductase and also a lower activity of nitrite reductase if compared to strain 23A, for example (Table 1). The two effective strains S₈ and S₉ had assimilatory nitrate reductase activity but did not show any "regulatory" nitrate reductase activity. The highest "regulatory" nitrate reductase activity was recorded with the effective strain S₁₉ (Table 1). All very effective and

Table 1. Assimilatory and regulatory nitrate reductase (NR) activities in *Rhizobium meliloti*

Strains	Assimilatory NR		Regulatory NR	
	nanog N-NO ₃ reduced per μg protein	nanog N-NO ₂ detected per μg protein	picog N-NO ₂ (μg protein) ⁻¹ min ⁻¹	Efficiency†
S ₁	572.2	44.95	31.50	E
S ₂	617.0	1.91	14.00	E
S ₃	274.2	0.68	10.78	E
S ₄	271.4	1.32	18.06	E
S ₅	329.2	0.36	14.42	E
S ₆	41.6	1.83	18.90	E
S ₇	217.2	0.38	9.38	E
S ₈	230.3	0.39	0	E
S ₉	94.3	6.98	0	E
S ₁₁	256.3	6.35	11.62	E
S ₁₂	300.0	0.47	10.78	E
S ₁₄	248.5	2.66	57.12	VE
S ₁₅	323.6	9.28	22.68	E
S ₁₆	372.8	3.49	111.72	E
S ₁₉	368.3	4.27	145.88	E
S ₂₀	302.1	0.86	61.74	E
S ₂₁	429.1	12.37	5.88	E
S ₂₂	390.0	1.66	73.64	E
V ₁	272.5	4.95	43.82	E
V ₂	368.5	79.74	58.38	E
V ₃	219.8	4.80	51.10	VE
V ₄	496.6	10.66	8.82	E
D ₁	477.7	11.98	73.22	I
D ₂	1168.1	154.54	52.36	E
D ₃	381.7	10.37	3.64	E
A ₂	355.9	2.88	50.82	VE
A ₃	408.7	0	1.40	VE
A ₄	360.8	16.98	1.54	VE
A ₅	1292.1	168.44	20.16	E
I ₁	32.0	4.23	27.44	I
I ₂	518.4	5.26	33.74	I
I ₃	382.1	4.44	21.56	E
I ₄	291.1	2.08	27.16	I
R ₁	507.0	4.06	21.70	E
E ₂	319.1	1.96	5.74	E
23A	1325.2	84.51	32.48	E
3DOa2oa	0	0	8.40	E
3DOa8	479.8	15.96	4.06	VE
54032	510.7	5.80	29.12	E
54033	251.4	4.55	10.64	E
Alfalfa D	580.3	12.26	75.88	I

†Based on yields obtained in the second cutting (Bordeleau et al. 1977). VE, very effective; E, effective; I, ineffective.

ineffective strains had assimilatory and "regulatory" nitrate reductase activities. This indicates that no relationship exists between the nitrate reductase activities and the symbiotic nitrogen fixation activity. In fact, statistical analysis did not show any

significant coefficient of correlation between the nitrate reductase activities and the dry matter yields of alfalfa (Table 2) which reflects the symbiotic nitrogen fixation activity of the tested strains (Bordeleau et al. 1977). The presence of assimilatory nitrate

Table 2. Coefficients of correlation (r) between nitrate reductase (NR) activity of *Rhizobium meliloti* and its nitrogenase activity with alfalfa

	Nitrogenase activity†		
	1	2	1+2
Assimilatory NR			
NO ₃ reduced	-0.14NS‡	0.03NS	-0.01NS
NO ₂ detected	-0.04NS	-0.05NS	-0.03NS
“Regulatory” NR	-0.08NS	-0.14NS	-0.13NS

†Nitrogenase activity expressed as alfalfa dry matter yield obtained at first cutting (1) and second cutting (2).

‡NS, not significant.

reductase in the very effective and ineffective strains (Table 1) confirms the previous observations made on ineffective mutants by Sik and Barabás (1977) that there is no structural connection between nitrate reductase and nitrogenase.

Sik and Barabás (1977) detected the “regulatory” nitrate reductase only in the effective wild type *R. meliloti*, and because this enzyme was absent in the ineffective mutants they suggested the presence of a common regulation factor governing the activities of this type of nitrate reductase and nitrogenase. In this study, the “regulatory” nitrate reductase was detected in both very effective and ineffective strains. This indicates that there is no common direct regulation factor governing the activities of the “regulatory” nitrate reductase and the symbiotic nitrogen fixation activities in the 41 tested strains. From these results, we can also conclude that the specific activities of the assimilatory and “regulatory” types of nitrate reductase do not offer useful physiological tools for rapid selection of very effective strains of *R. meliloti*.

BARABÁS, I. and SIK, T. 1979. Phenotypic reversion of nitrogenase in pleiotropic mutants of *Rhizobium meliloti*. *Can. J. Microbiol.* **25**: 298–301.

BORDELEAU, L. M., ANTOUN, H. and LACHANCE, R. A. 1977. Effets des souches de *Rhizobium meliloti* et des coupes successives de la luzerne (*Medicago sativa*) sur la fixation symbiotique d'azote. *Can. J. Plant Sci.* **57**: 433–439.

BORDELEAU, L. M. and ANTOUN, H. 1978. Une méthode rapide de présélection des souches de *Rhizobium meliloti*. *Can. J. Plant Sci.* **58**: 1125–1126.

BRADFORD, M. M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* **36**: 207–212.

BREMNER, J. M. 1965. Inorganic forms of nitrogen: Nitrite by colorimetric methods. Pages 1219–1224 in C. A. Black, ed. *Methods of soil analysis*. American Society of Agronomy, Madison, Wisc.

CATALDO, D. A., HAROON, M., SCHRAEDER, L. E. and YOUNGS, V. L. 1975. Rapid colorimetric determination of nitrate in plant tissue by nitration of salicylic acid. *Commun. Soil Sci. Plant Anal.* **6**: 71–80.

CHENIAE, G. and EVANS, H. J. 1960. Physiological studies on nodule-nitrate reductase. *Plant Physiol.* **35**: 454–462.

EVANS, H. J. and RUSSEL, S. A. 1971. Physiological chemistry of symbiotic nitrogen fixation by legumes. Pages 191–244 in J. R. Postgate, ed. *The chemistry and biochemistry of nitrogen fixation*. Plenum Press, London, New York.

KONDOROSI, A., BARABÁS, I., SVÁB, Z., OROSZ, L. and SIK, I. 1973. Evidence for common genetic determinants of nitrogenase and nitrate reductase in *Rhizobium meliloti*. *Nat. New Biol.* **246**: 153–154.

NICHOLAS, D. J. and NASON, A. 1957. Determination of nitrate and nitrite. Pages 981–984 in S. P. Colowick and N. O. Kaplan, eds. *Methods in enzymology*. Vol. III. Academic Press, New York, N. Y.

OROSZ, L. and SIK, T. 1970. Genetic mapping of *Rhizobiophage* 16-3. *Acta Microbiol. Acad. Sci. Hung.* **17**: 185–194.

SIK, T., KONDOROSI, A., BARABÁS, I. and SVÁB, Z. 1976. Nitrate reductase and effectiveness in *Rhizobium*. Pages 374–382 in W. E. Newton and C. J. Nyman, eds. *Proc. 1st Int. Symp. on Nitrogen Fixation*. Washington State Univ. Press.

SIK, T. and BARABÁS, I. 1977. The correlation of nitrate reductase and nitrogenase in *Rhizobium* symbiosis. Pages 365–373 in W. Newton, J. R. Postgate and C. Rodriguez-Barrueco, eds. *Recent developments in nitrogen fixation*. Academic Press, London.