

Systemically suppressed isoflavonoids and their stimulating effects on nodulation and mycorrhization in alfalfa split-root systems

Jean Guy Catford · Christian Staehelin ·
Geneviève Larose · Yves Piché · Horst Vierheilig

Received: 1 March 2006 / Accepted: 11 April 2006 / Published online: 25 May 2006
© Springer Science+Business Media B.V. 2006

Abstract In split-root systems of alfalfa (*Medicago sativa* L.), already existing nodules or arbuscular mycorrhizal roots suppress further establishment of symbiosis in other root parts, a phenomenon named autoregulation. Roots treated with rhizobial nodulation signals (Nod factors) induce a similar systemic suppression of symbiosis. In order to test the hypothesis that flavonoids play a role in this systemic suppression, split-root systems of alfalfa plants were inoculated on one side of the split-root system with *Sinorhizobium meliloti* or *Glomus mosseae* or were treated with Nod factor. HPLC-analysis of alfalfa root extracts from both sides of the split-root system revealed a persistent local and systemic accumulation pattern of some flavonoids

associated with the different treatments. The two flavonoids, formononetin and ononin, could be identified to be similarly altered after rhizobial or mycorrhizal inoculation or when treated with Nod factor. Exogenous application of formononetin and ononin partially restored nodulation and mycorrhization pointing towards the involvement of these two secondary compounds in the autoregulation of both symbioses.

Keywords Arbuscular mycorrhiza · Autoregulation · Flavonoid · *Medicago sativa* · Nodulation · *Sinorhizobium meliloti*

Abbreviations

AMF Arbuscular mycorrhizal fungi
NFs Nod factors

J. G. Catford · Geneviève Larose · Y. Piché
CRBF, Faculté de Foresterie et de Géomatique,
Université Laval, Pavillon C.-E. Marchand, G1K 7P4
Ste-Foy, Québec, Canada

C. Staehelin
State Key Laboratory of Biocontrol, School of Life
Sciences, Sun Yat-Sen (Zhongshan) University,
Xingangxi Road 135, Guangzhou 510275, China

H. Vierheilig (✉)
Department für Angewandte Pflanzenwissenschaften
und Pflanzenbiotechnologie, Institut für
Pflanzenschutz, Universität für Bodenkultur Wien,
Peter-Jordan-Str. 82, A-1190 Wien, Austria
e-mail: nonhorst@boku.ac.at

Introduction

Apart from abiotic soil factors such as nutrient and water availability, the growth of land plants is positively affected by symbiotic microorganisms. Nitrogen-fixing rhizobia (*Rhizobiaceae* bacteria) infect roots of legumes and induce the formation of nodules. More than 80% of all land plants enter symbiosis with AMF that acquire nutrients, such as phosphorus, from the soil. In the nodule

symbiosis, the infection process depends on rhizobial nodulation signals, called NFs. During the early steps of infection, the two symbiotic associations seem to share a number of signalling steps. There is increasing evidence that components required for NF signalling also play a role in establishment of the AMF symbiosis (Guinel and Geil 2002; Hirsch and Kapulnik 1998; Riely et al. 2004; Staehelin et al. 2001; Xie et al. 1995). Similarities between the two symbioses also have been suggested during later symbiotic stages (Duc et al. 1989; Vierheilig 2004a; Vierheilig and Piché 2002).

Once a plant has formed nodules, further nodulation is suppressed in other parts of the root system by a long-distance signal exchange. This regulatory mechanism has been named autoregulation of nodulation (for details see review Caetones-Anollés and Gresshoff 1991). Interestingly, autoregulation of symbiosis has also been reported for the AMF root colonization. Working with split-root systems of barley, alfalfa and soybean, it could be shown that root colonization of one side of a split-root system strongly suppressed mycorrhization of “autoregulated roots” on the other side (Catford et al. 2003; Meixner et al. 2005; Vierheilig 2004b; Vierheilig et al. 2000a, b). Recently, Catford et al. (2003) reported that pre-inoculation with *Sinorhizobium meliloti* of one side of alfalfa split-roots negatively affected AMF root colonization on the other side. Conversely, pre-inoculation with AMF inhibited subsequent nodulation of autoregulated roots. Moreover, application of NFs to one side of the split-root system suppressed nodulation and mycorrhization on the other side. Meixner et al. (2005) reported that the soybean supernodulating mutant *nts1007* (mutated in the receptor kinase gene *GmNARK* (Searle et al. 2003), which lacks the autoregulatory mechanism to control nodulation, did not autoregulate AMF root colonization. All these data together indicated a similar mechanism of autoregulation in both symbioses.

Flavonoids—and isoflavonoids in particular—exuded by plant roots have been reported to activate expression of rhizobial *nod* genes required for synthesis of NFs. There is some information available on the possible limitation of

NF synthesis by flavonoids (Hungria and Phillips 1993; Schlaman et al. 1991; Zaat et al. 1988) and about the effect of additional flavonoid application on nodulation (Cunningham et al. 1991; Jain et al. 1990; Kapulnik et al. 1987; Kosslak et al. 1990; Zhang and Smith 1995). Most recently, Novák et al. (2002) showed that application of specific flavonoids to pea roots either increased or decreased nodulation. Flavonoids have also been suggested to be involved in the regulation of mycorrhization (Vierheilig and Piché 2002; Vierheilig 2004a). In a number of studies with various host plants, it has been shown that roots treated with certain flavonoids exhibited increased AMF root colonization (Nair et al. 1991; Scervino et al. 2005; Siqueira et al. 1991; Vierheilig et al. 1998a; Xie et al. 1995).

To our knowledge, no data are available yet whether flavonoids are involved in autoregulation of symbiosis. In this study, we report on alterations of isoflavonoid levels in split-root systems of alfalfa plants, which were inoculated on one side either with *S. meliloti* or with the AM fungus *Glomus mosseae*. In both symbioses, formononetin and its glycoside ononin were locally and systemically down-regulated. In a further series of experiments, we applied these isoflavonoids to split-roots and studied their effects on nodulation and symbiosis.

Materials and methods

Biological material, growing conditions and experimental design

Alfalfa (*Medicago sativa* L. cv. Sitel) plants were inoculated with *G. mosseae* (Nicolson & Gerdemann) Gerd. & Trappe (BEG12; European Bank for the Glomales) or for nodulation experiments with *S. meliloti* strain 1021. Corresponding control plants were grown under the same conditions. Alfalfa plants were grown in a growth chamber [day/night cycle: 16/8 h, 23/19°C; PAR (photosynthetically active radiation): 300 $\mu\text{mol}/\text{m}^{-2} \text{ s}^{-1}$; relative humidity 50%]. All experiments were repeated twice with five replicates per treatment.

Split-root systems

Alfalfa seeds were surface-sterilized by soaking in 0.75% sodium hypochlorite for 5 min, rinsed with tap water and germinated in pots in a steam-sterilized (30 min, 121°C) mixture of silicate sand, TurFace (Applied Industrial Materials Corp.; Buffalo Grove, Illinois, USA) and soil (1:1:1 by vol). To induce the development of lateral roots, tips of main roots from 10 days old plantlets were excised. Plantlets were then placed into the same substrate as described above. Twenty-five days later, split-root systems were established as described previously (Vierheilig et al. 2000a). The split-root system consists of two units, each containing one half of the alfalfa root system. The two compartments are separated by an impermeable PVC membrane in order to prevent any flow of molecules or root growth from one side to the other. Thus, symbiosis can be established in split-root systems on one side without any contact on the other side. Control plants were grown under the same growth conditions as the inoculated plants.

Isoflavonoid levels in a split-root system of alfalfa with one side inoculated with *S. meliloti*

Sinorhizobium meliloti strain 1021 was cultivated at 25°C in a TY–streptomycin (50 µg/ml) medium (Beringer 1974) on a rotary shaker (200 rpm) to reach an $OD_{600} = 0.25$. Bacteria were then harvested by centrifugation (6000g × 10 min) and resuspended in sterile H₂O. For inoculation with *S. meliloti*, one side of a split-root system was inoculated with 10 ml of this bacterial suspension (one day after the transfer of plants into the split-root system). After 15 days, the split-root system and non-inoculated control plants were harvested. The nodulated side of the split-root formed 168 ± 11 nodules per 100 mg root (FW), whereas no nodules were seen in the non-nodulated side. The harvested plant material was then frozen (–80°C) and lyophilized. Flavonoid analysis was performed with samples of all three different treatments (control roots; nodulated roots from the inoculated side of the split-root system; roots from the non-inoculated side of the split-root system).

Isoflavonoid levels in a split-root system of alfalfa with one side treated with NFs

The tetrameric Nod factor NodSm-IV(C16:2, S) was purified from *S. meliloti* strain 1021(pEK327) (Schultze et al. 1992; kindly provided by Eva Kondorosi, CNRS, Gif-sur-Yvette, France). Supernatants of bacterial cultures were extracted with *n*-butanol and fractionated by reverse-phase HPLC (Waters C18 column) with 35% acetonitrile/water, 40 mM ammonium acetate as the mobile phase. The fraction containing NodSm-IV(C16:2, S) was collected and desalted as described by Staehelin et al. (2000).

Eight days after establishment of the split-root system, 4 ml of a 10^{–8} M NF solution (dissolved in water) were daily applied to one side (volume 65 ml) of the system. After 8 days, the NF-treated side, the roots from the other side of the split-root system and non-treated control plants were harvested. Lyophilized root material was used for flavonoid analysis.

Isoflavonoid levels in a split-root system of alfalfa inoculated with *G. mosseae*

The outer side of each split-root compartment was equipped with a Nylon screen (Nitex monofilament, 100 µm mesh), which can be penetrated by hyphae but not by roots. To inoculate alfalfa with *G. mosseae*, one outer side of the split-root compartment was joined at the day of transfer of plants into the split-root system with a donor compartment, which on its sides was also equipped with the Nylon screen. This donor compartment contained beans (*Phaseolus vulgaris* L. cv. Sun Gold) colonized by *G. mosseae*. Hyphae of the AMF penetrated the Nylon screen and colonized one side of the split-root system, however, the second side of the split-root system remained without root colonization by the AMF. After 20 days the mycorrhizal and non-mycorrhizal roots of the split-root system and the roots of the control plants were harvested and frozen (–80°C). After lyophilization, roots were used for flavonoid analysis. In all samples, the degree of AMF root colonization was determined at the time of harvest.

Effects of isoflavonoid application on nodulation in split-root systems pre-inoculated with *S. meliloti*

One side of the split-root system of alfalfa plants was pre-inoculated with *S. meliloti* as described above. After 15 days, when nodules had formed on the first side of the split-root system (168 ± 11 nodules per 100 mg root FW), the second side was inoculated with *S. meliloti*. At the same time formononetin or ononin were applied to the second side of the split-root system. The application of isoflavonoids was repeated every second day during 15 days. Thereafter the root material was harvested and the number of nodules determined.

Effects of isoflavonoid application on mycorrhizal colonization in split-root systems pre-colonized by *G. mosseae*

One side of the split-root system was pre-inoculated with *G. mosseae* as described above. After 20 days, when the AMF symbiosis was well established on the first side of the split-root system ($54 \pm 3\%$ root colonization), the second side was inoculated with *G. mosseae*. At the same time formononetin or its glycoside ononin were applied to the second side of the split-root system and this treatment was repeated every second day during 20 days. Thereafter plants were harvested and the degree of AMF root colonization was determined.

Application of isoflavonoids to roots

Formononetin (1.34 mg) or ononin (2.15 mg) were dissolved in 4 ml ethanol and diluted with H₂O to a final volume of 1,000 ml. In a pre-experiment, it was found that 0.4% ethanol did not exhibit any effect on AMF root colonization. Volumes of 10 ml of the diluted isoflavonoid solutions (with a final concentration of 5 μ M) were applied every second day to the second side of the split-root system.

Determination of nodule number and AMF root colonization

At the time of harvest, roots were carefully rinsed with water and the root fresh weight and the

number of nodules were determined. To estimate AMF root colonization, several roots from each plant were cleared by boiling in 10% KOH and stained according to the method of Vierheilig et al. (1998b) by boiling in a 5% ink (Shaeffer; black)/household vinegar (=5% acetic acid) solution. Stained roots were observed with a light stereomicroscope to determine the percentage of root colonization according to a modified method of Newman (1966).

Root extraction and high-performance liquid chromatography

At harvest roots were rinsed with tap water and root fresh weight (FW) was determined. Thereafter roots were stored at -20°C and finally lyophilized. The freeze-dried roots (0.5–2.0 g) were crushed and flavonoids were extracted in 100% acetone overnight at 4°C as described by Edwards and Kessmann (1992). The mixture was filtered and the extract was evaporated. The residue was dissolved in 0.5 ml of methanol.

HPLC analysis was performed on a reverse-phase analytical column (Kingsorb C₁₈, 150 \times 4.6 mm id., 3 μ m) employing a diode array detector. The flavonoids were separated by a 45 min linear gradient from 20 to 75% solvent B (solvent A = H₂O + 0.1% TFA, solvent B = acetonitrile) with a flow rate of 0.5 ml/min. The chromatograms were recorded at 210, 250, 260, and 350 nm depending on the UV absorption maximum of the compounds. The compound identification was based on the peak retention times and the comparison with the UV spectra of the standards.

Retention times: Ononin (formononetin-7-*O*-glucoside) 14.74 min (Sequoia Research Products), 7,4'-dihydroxyflavone 15.50 min (Indofine), 7,4'-dihydroxyflavanone 18.30 min, 4,4'-dihydroxy-2'-methoxychalcone 20.30 min (kindly provided by D.A. Phillips, University of California, Davis, USA), daidzein (4',7-dihydroxyisoflavone) 17.20 min (Indofine), genistein (4',5,7-trihydroxyisoflavone) 23.11 min (Indofine), coumestrol 23.21 min (Fluka), formononetin (7-hydroxy-4'-methoxyisoflavone) 27.14 min (Indofine), medicarpin 31.44 min (Sequoia Research Products).

Results

Analysis of isoflavonoids in alfalfa roots indicated for each treatment a specific isoflavonoid pattern. Levels of certain isoflavonoids remained similar in all samples, whereas others varied in response to *S. meliloti* or *G. mosseae*. The data presented in this study are restricted to those isoflavonoids that exhibited similar altered levels during symbiosis and Nod factor treatment, namely formononetin and its glycoside ononin, medicarpin and daidzein. The levels of 7,4'-dihydroxyflavone, 7,4'-dihydroxyflavanone, 4,4'-dihydroxy-2'-methoxychalcone and genistein were not changed in any of the treatments (data not shown), whereas only in mycorrhizal plants coumestrol levels were locally and systemically enhanced (data not shown).

Compared to non-inoculated control plants, inoculation with *S. meliloti* on one side of a split-root system resulted in decreased levels of formononetin and ononin in the inoculated part of the split-root (nodulated roots). Interestingly, levels of formononetin and ononin were also low in the non-inoculated side of the split-root, indicating a systemic effect from inoculated to non-inoculated root parts. Medicarpin levels were low

in non-inoculated control plants and slightly reduced in both parts of the split-root system. Daidzein levels in nodulated roots were elevated, whereas the non-inoculated part of the split-root system exhibited levels comparable to those in non-inoculated control plants (Fig. 1). Application of NFs to one side of a split-root system strongly reduced the levels of formononetin, ononin and medicarpin. Lowest levels were found for formononetin, which were about 10-fold lower than in non-inoculated control plants. Levels of these isoflavonoids were also systemically down-regulated in the half of the split-root, which has been not treated with NFs. In contrast to the infection with *S. meliloti* resulting in nodulated roots, application of NFs to roots did not affect the daidzein levels in our split-root experiments (Fig. 2).

When one side of the split-root system was inoculated with the AMF *G. mosseae*, the levels of formononetin and ononin in mycorrhizal roots were not altered compared to non-inoculated control plants. The non-inoculated side of the split-root however, exhibited low levels of formononetin and ononin, indicating a systemic regulation of these isoflavonoids. Medicarpin and




Fig. 1 Flavonoid pattern in a split-root system of alfalfa with one side inoculated with *Sinorhizobium meliloti* (+R side) and the other side without inoculation (–R side) and

in roots of non-inoculated control plants (–R). Data represent mean \pm SE ($n = 5$)




Fig. 2 Flavonoid pattern in a split-root system of alfalfa with one side treated with nod factor (+NF side) and the other side without NF treatment (–NF side) and in roots of non-treated control plants (–NF). Data represent mean \pm SE ($n = 5$)

daidzein levels were stimulated in the side with mycorrhizal roots, when compared to non-mycorrhizal control plants. Daidzein was also strongly accumulated in the non-mycorrhizal part of the split-root system (Fig. 3).

Taken these data together, they indicate that formononetin and its glycoside ononin are systemically down-regulated in response to *S. meliloti* infection, treatment with NFs and root colonization by *G. mosseae*. Hence, low levels of formononetin and ononin in non-infected roots are metabolic markers for autoregulated roots. We therefore tested in a second series of experiments whether an application of formononetin or ononin to autoregulated roots can stimulate nodulation and mycorrhization. To induce autoregulation signalling, a split-root system was inoculated on one side. As reported previously (Catford et al. 2003), pre-inoculation

with *S. meliloti* on one side of the split-root system resulted in a drastic suppression of nodule formation on the later inoculated side of the split-root system. As shown in Fig. 4, only few nodules were counted on the second half of the split-root system, whereas plants, which have been not pre-inoculated, formed about 10-fold more nodules. When formononetin was applied to the second side of the split-root system of pre-inoculated plants, nodule formation on this autoregulated side was strongly increased. Nodule numbers reached values that were similar to those from not pre-inoculated plants. This finding indicates that formononetin counteracts autoregulation of nodulation. Similar to formononetin, its glycoside ononin also promoted nodulation on the autoregulated side of split-roots, albeit to a lower extent than the aglycone (Fig. 4).





Fig. 3 Flavonoid pattern in a split-root system of alfalfa with one side inoculated (+M side) with *Glomus mosseae* and the other side without inoculation (–M side) and in

roots of non-inoculated control plants (–M). Data represent mean \pm SE ($n = 5$)

Fig. 4 Effect of rhizobial pre-inoculation of one half of the split-root system of alfalfa on subsequent nodule formation on the second half of the split-root system with or without application of formononetin or ononin (10 ml of a 5 μ M solution every second day) to the second half. Data represent mean \pm SE ($n = 5$)



Pre-inoculation with *G. mosseae* on one side of the split-root system resulted in a drastic suppression of AMF root colonization on the second side of the split-root system, indicating autoregulation of mycorrhization (Catford et al. 2003; Fig. 5). When the glycoside ononin was applied to the autoregulated side of the split-root, AMF root colonization by *G. mosseae* was enhanced on this side. AMF root colonization, however, did not reach the degree of those plants that were not pre-inoculated with *G. mosseae*. In contrast to ononin, application of the aglycone formononetin did not affect mycorrhization in our split-root experiments (Fig. 5).


Discussion

Inoculation of one side of an alfalfa split-root system with *S. meliloti*, *G. mosseae* or the application of NF to one side of an alfalfa split-root system has been reported to suppress nodulation or mycorrhization on the other side (Catford et al. 2003). In this study, we have used split-root systems in order to test whether isoflavonoid levels in alfalfa roots are systemically altered in response to *S. meliloti* and *G. mosseae*. We could demonstrate that infection with either *S. meliloti* or *G. mosseae* influenced the levels of isoflavonoids in roots, which were not in contact with the invading symbionts. Hence, specific isoflavonoids in roots are systemically regulated upon infection.

Our study shows significant symbiosis-related alterations for four isoflavonoids: formononetin, its glycoside ononin, medicarpin and daidzein. In the pathway for biosynthesis of alfalfa isoflavonoids, formononetin is a precursor of the phytoalexine medicarpin. Daidzein however, is most likely not an intermediate in medicarpin synthesis. It has been proposed that 4'-*O*-methyltransferase is the key enzyme at the branching point for medicarpin synthesis (Liu and Dixon 2001). In this view, our data indicate that root colonization by *G. mosseae* activates the medicarpin synthesis pathway as reported previously (Larose et al. 2002), whereas infection with *S. meliloti* down-regulates this pathway. Daidzein levels accumulated in roots in response to both symbionts. Similar to the infection with *S. meliloti*, application of NFs resulted in reduced levels of formononetin, ononin and medicarpin, indicating a suppression of this synthetic pathway. A similar, albeit weaker, suppression of formononetin occurred in alfalfa roots that were challenged with Nod factors for 50 h (C.S., G.L. and H.V., unpublished results; Vierheilig et al. 2004b). The suppression effect of NFs on specific isoflavonoids involved in medicarpin synthesis is remarkable, as various genes involved in isoflavonoid biosynthesis have been reported to be up-regulated in response to NFs (i.e., in *M. truncatula*; Lohar et al. 2006).

In the split-root experiments of this study, inoculation with *S. meliloti* on one side also

Fig. 5 Effect of AM pre-inoculation of one half of the split-root system of alfalfa on subsequent AMF root colonization of the second half of the split-root system with or without application of formononetin or ononin (10 ml of a 5 μ M solution every second day) to the second half. Data represent mean \pm SE ($n = 5$)



altered isoflavonoids of the medicarpin synthesis pathway on the other side (Fig. 1). We conclude that levels of formononetin and its glycoside ononin are not only locally reduced in response to *S. meliloti*, but also systemically down-regulated in parts of the root system that were not in contact with *S. meliloti*. When NFs were used for the experiment, we observed a similar systemic reduction for formononetin, ononin as well as medicarpin (Fig. 2). As NFs are able to induce autoregulation of symbiosis in alfalfa split-roots (Catford et al. 2003), we suggest that systemic down-regulation of formononetin and ononin levels in non-treated root parts is a metabolic response to the putative autoregulation signals that mediate suppression of symbiosis. Root colonization by *G. mosseae* also can elicit suppression of symbiosis in other parts of the root system (Catford et al. 2003) and we therefore expected a systemic down-regulation of formononetin and ononin also during the mycorrhizal symbiosis. Indeed, our data indicate very low levels of these isoflavonoids in the non-inoculated side of the split-root (Fig. 3). Thus, all known treatments eliciting autoregulation signaling—i.e., inoculation with *S. meliloti*, application of NFs, and inoculation with *G. mosseae*—suppressed accumulation of formononetin and ononin in the whole root system.

Nodule formation and mycorrhization is suppressed in autoregulated parts of the root system. Our isoflavonoid analyses indicate low levels of formononetin and ononin in all autoregulated roots. We therefore wondered whether these isoflavonoids are limiting factors for establishment of symbiosis in autoregulated roots. Our data point to this direction, as application of formononetin or ononin to autoregulated roots promoted nodule formation. Ononin showed a similar stimulating effect on establishment of the AMF symbiosis. With other words, application of isoflavonoids to roots inactivated or compensated the symbiosis-suppressing effects induced by long-distance autoregulation signals.

In the rhizobial symbiosis, flavonoids have been proposed to play a limiting factor for NF synthesis and optimal establishment of symbiosis (Zaat et al. 1988; Schlaman et al. 1991; Hungria and Phillips 1993). Interestingly, exogenous

application of flavonoids to legumes has been reported to increase or decrease nodulation (Cunningham et al. 1991; Jain et al. 1990; Kapulnik et al. 1987; Kossak et al. 1990; Novák et al. 2002; Zhang and Smith 1995;). One possible explanation is that specific flavonoids induce rhizobial *nod* gene expression required for synthesis of NFs, whereas others do not or even repress NF synthesis. Formononetin however, did exhibit *nod* gene inducing activity on a *S. meliloti* strain (Zuanazzi et al. 1998). It is therefore unlikely that the promoting effect on nodulation induced by formononetin (Fig. 4) is related to NF synthesis of *S. meliloti* strain 1021.

Flavonoids have also been suggested to be involved in the regulation of mycorrhization (Vierheilig and Piché 2002; Vierheilig 2004a). A number of studies with various host plants showed that roots challenged with specific flavonoids increased AM root colonization (Nair et al. 1991; Scervino et al. 2005; Siqueira et al. 1991; Vierheilig et al. 1998a; Xie et al. 1995). Ononin promoted AMF root colonization in our experiments (Fig. 5), indicating that certain isoflavonoids have antagonistic effects on autoregulation of AMF root colonization. In contrast to the nodule symbiosis, formononetin did not stimulate mycorrhization in autoregulated parts of the root system. Hence, although autoregulation of nodulation and mycorrhization seem to share some common signalling events (Catford et al. 2003; Meixner et al. 2005), the effects of systemically regulated isoflavonoids on establishment of symbiosis are different.

Application of isoflavonoids to roots might induce multiple responses. One possibility is that flavonoids and NFs interfere with the polar auxin transport in roots (Boot et al. 1999; Jacobs and Rubery 1988; Mathesius et al. 1998). Treatment of roots with auxin transport inhibitors induced the formation of nodule-like structures in alfalfa (Hirsch et al. 1989) and promoted mycorrhizal colonization in various plants, including alfalfa (Xie et al. 1996, 1998). Moreover, analysis of phytohormones in soybean roots suggested that auxin levels might be involved in autoregulation of symbiosis (Caba et al. 2000; Meixner et al. 2005). Future split-root experiments are required to test whether formononetin and ononin alter

auxin transport in the autoregulated part of alfalfa roots and whether other isoflavonoids have similar effects on roots.

To summarize, our study shows that levels of specific isoflavonoids are systemically suppressed in response to *S. meliloti*, NFs and *G. mosseae*. Comparing the systemic isoflavonoid accumulation pattern in all three treatments, we propose that formononetin and ononin are possibly involved in the systemic suppression of nodulation and mycorrhization. Application of formononetin and ononin to autoregulated roots promoted nodulation, whereas ononin stimulated mycorrhization. Future studies with plant mutants lacking autoregulation are required to demonstrate that the long-distance signals controlling autoregulation of symbiosis are identical to those signals that systemically regulate isoflavonoid levels.

Acknowledgements This work was partially funded by a OECD grant to H.V.C.S. was supported by the Science and Technology Department of Guangdong and by the Guangdong Natural Foundation.

References

- Beringer JE (1974) R factor transfer in *Rhizobium leguminosarum*. J Gen Microbiol 84:188–198
- Boot KJM, van Brussel AA, Tak T, Spaik H, Kijne JW (1999) Lipochitin oligosaccharides from *Rhizobium leguminosarum* bv. *viciae* reduce auxin transport capacity in *Vicia sativa* subsp. *nigra* roots. Mol Plant-Microbe Interact 12:839–844
- Caba JM, Centeno ML, Fernandez B, Gresshoff PM, Ligeró F (2000) Inoculation and nitrate alter phytohormone levels in soybean roots: differences between a supernodulating mutant and the wild type. Planta 211:98–104
- Caetano-Anollés G, Gresshoff PM (1991) Plant genetic control of nodulation. Annu Rev Microbiol 45:345–382
- Catford JG, Staehelin C, Lerat S, Piché Y, Vierheilig H (2003) Suppression of arbuscular mycorrhizal colonization and nodulation in split-root systems of alfalfa after pre-inoculation and treatment with Nod factors. J Exp Bot 54:1481–1487
- Cunningham S, Kollmeyer WD, Stacey G (1991) Chemical control of interstrain competition for soybean nodulation by *Bradyrhizobium japonicum*. Appl Environ Microbiol 57:1886–1892
- Duc G, Trouvelot A, Gianinazzi-Pearson V, Gianinazzi S (1989) First report of non-mycorrhizal plant mutants (Myc-) obtained in pea (*Pisum sativum* L.) and fababea (*Vicia faba* L.). Plant Sci 60:215–222
- Edwards R, Kessmann H (1992) Isoflavonoid phytoalexins and their biosynthetic enzymes. In: Gurr SJ, McPherson MJ, Bowles DJ (eds) Molecular plant pathology: A practical approach. Oxford University Press, Oxford, pp 45–62
- Guinel FC, Geil RD (2002) A model for the development of the rhizobial and arbuscular mycorrhizal symbioses in legumes and its use to understand the roles of ethylene in the establishment of these two symbioses. Can J Bot 80:695–720
- Hirsch AM, Bhuvaneshwari TV, Torrey JG, Bisseling T (1989) Early nodulin genes are induced in alfalfa root outgrowths elicited by auxin transport inhibitors. Proc Natl Acad Sci USA 86:1244–1248
- Hirsch AM, Kapulnik Y (1998) Signal transduction pathways in mycorrhizal associations: comparisons with the *Rhizobium*-legume symbiosis. Fung Gen Biol 23:205–212
- Hungria M, Phillips DA (1993) Effects of a seed color mutation on rhizobial *nod*-gene-inducing flavonoids and nodulation in common bean. Mol Plant-Microbe Interact 6:418–422
- Jacobs M, Rubery PH (1988) Naturally occurring auxin transport regulators. Science 241:346–349
- Jain V, Garg N, Nainawatee HS (1990) Naringenin enhanced efficiency of *Rhizobium meliloti*-alfalfa symbiosis. World J Microbiol Biotechnol 6:434–436
- Kapulnik Y, Joseph CM, Phillips DA (1987) Flavone limitations to root nodulation and symbiotic nitrogen fixation in alfalfa. Plant Physiol 84:1193–1196
- Kosslak RM, Joshi RS, Bowen BA, Paaren HE, Appelbaum ER (1990) Strain-specific inhibition of *nod* gene induction in *Bradyrhizobium japonicum* by flavonoid compounds. Appl Environ Microbiol 56:1333–1341
- Larose G, Chenevert R, Moutoglis P, Gagne S, Piche Y, Vierheilig H (2002) Flavonoid levels in roots of *Medicago sativa* are modulated by the developmental stage of the symbiosis and the root colonizing arbuscular mycorrhizal fungus. J Plant Physiol 159:1329–39
- Liu CJ, Dixon RA (2001) Elicitor-induced association of isoflavone *O*-methyltransferase with endomembranes prevents the formation and 7-*O*-methylation of daidzein during isoflavonoid phytoalexin biosynthesis. Plant Cell 13:2643–2658
- Lohar DP, Sharopova N, Endre G, Penuela S, Samac D, Town C, Silverstein KAT, VandenBosch KA (2006) Transcript analysis of early nodulation events in *Medicago truncatula*. Plant Physiol 140:221–234
- Mathesius U, Schlaman HRM, Spaik HP, Sautter C, Rolfe BG, Djordjevic MA (1998) Auxin transport inhibition precedes root nodule formation in white clover roots and is regulated by flavonoids and derivatives of chitin oligosaccharides. Plant J 14:23–34
- Meixner C, Ludwig-Müller J, Miersch O, Gresshoff P, Staehelin C, Vierheilig H (2005) Lack of mycorrhizal autoregulation and phytohormonal changes in the supernodulating soybean mutant *nts1007*. Planta 222:709–715

- Nair MG, Safir GR, Siqueira JO (1991) Isolation and identification of vesicular-arbuscular mycorrhizal-stimulatory compounds from clover (*Trifolium repens*) roots. *Appl Environ Microbiol* 57:434–439
- Newman EI (1966) A method of estimating the total length of root in a sample. *J Appl Ecol* 3:139–145
- Novák K, Chovanec P, Škrdleta V, Kropáčová M, Lisá L, Němcová M (2002) Effect of exogenous flavonoids on nodulation of pea (*Pisum sativum* L.). *J Exp Bot* 53:1735–1745
- Riely BK, Ané JM, Varma Penmetsa R, Cook DG (2004) Genetic and genomic analysis in model legumes bring Nod-factor signaling to center stage. *Curr Opin Plant Biol* 7:408–413
- Scervino JM, Ponce MA, Erra-Bassels R, Vierheilig H, Ocampo JA, Godeas A (2005) Arbuscular mycorrhizal colonization of tomato by *Gigaspora* and *Glomus* species in presence of roots flavonoids. *J Plant Physiol* 162:625–633
- Schlaman HRM, Horvath B, Vijgenboom E, Okker RJH, Lugtenberg BJJ (1991) Suppression of nodulation gene expression in bacteroids of *Rhizobium leguminosarum* biovar *viciae*. *J Bacteriol* 173:4277–4287
- Schultze M, Quiclet-Sire B, Kondorosi E, Virelizier H, Glushka JN, Endre G, Géro SD, Kondorosi A (1992) *Rhizobium meliloti* produces a family of sulfated lipooligosaccharides exhibiting different degrees of plant host specificity. *Proc Natl Acad Sci USA* 89:192–196
- Searle IR, Men AE, Laniya TS, Buzas DM, Iturbe-Ormaetxe I, Carroll BJ, Gresshoff PM (2003) Long-distance signaling in nodulation directed by a CLAVATA1-like receptor kinase. *Science* 299:109–112
- Siqueira JO, Safir GR, Nair MG (1991) Stimulation of vesicular-arbuscular mycorrhiza formation and growth of white clover by flavonoid compounds. *New Phytol* 118:87–93
- Staelin C, Charon C, Crespi M, Boller T, Kondorosi A (2001) *Medicago truncatula* plants overexpressing the early nodulin gene *enod40* exhibit accelerated mycorrhizal colonization and enhanced formation of arbuscules. *Proc Natl Acad Sci USA* 98:15366–15371
- Staelin C, Schultze M, Tokuyasu K, Poinso V, Promé JC, Kondorosi E, Kondorosi A (2000) N-deacetylation of *Sinorhizobium meliloti* Nod factors increases their stability in the *Medicago sativa* rhizosphere and decreases their biological activity. *Mol Plant-Microbe Interact* 13:72–79
- Vierheilig H (2004a) Regulatory mechanisms during the plant-arbuscular mycorrhizal fungus interaction. *Can J Bot* 82:1166–1176
- Vierheilig H (2004b) Further root colonization by arbuscular mycorrhizal fungi in already mycorrhizal plants is suppressed after a critical level of root colonization. *J Plant Physiol* 161:339–341
- Vierheilig H, Bago B, Albrecht C, Poulin MJ, Piché Y (1998a) Flavonoids and arbuscular-mycorrhizal fungi. In: Manthey JA, Buslig BS (eds) Flavonoids in the living system. Plenum Press, New York, USA, pp 9–33
- Vierheilig H, Coughlan AP, Wyss U, Piché Y (1998b) Ink and vinegar, a simple staining technique for arbuscular-mycorrhizal fungi. *Appl Environ Microbiol* 64:5004–5007
- Vierheilig H, Garcia-Garrido JM, Wyss U, Piché Y (2000a) Systemic suppression of mycorrhizal colonization of barley roots already colonized by AM fungi. *Soil Biol Biochem* 32:589–595
- Vierheilig H, Maier W, Wyss U, Samson J, Strack D, Piché Y (2000b) Cyclohexenone derivative- and phosphate-levels in split-root systems and their role in the systemic suppression of mycorrhization in precolonized barley plants. *J Plant Physiol* 157:593–599
- Vierheilig H, Piché Y (2002) Signalling in arbuscular mycorrhiza, facts and hypotheses. In: Buslig B, Manthey J (eds) Flavonoids in cell functions. Kluwer Academic/Plenum Publishers, New York, pp 23–39
- Xie ZP (1996) Effects of Nod factors and phytohormones on the symbiotic interaction of legumes with rhizobia and arbuscular mycorrhizal fungi. PhD thesis. University of Basel, Switzerland
- Xie ZP, Müller J, Wiemken A, Broughton WJ, Boller T (1998) Nod factors and *tri*-iodobenzoic acid stimulate mycorrhizal colonization and affect carbohydrate partitioning in mycorrhizal roots of *Lablab purpureus*. *New Phytol* 139:361–366
- Xie ZP, Staehelin C, Vierheilig H, Wiemken A, Jabbouri S, Broughton WJ, Vögeli-Lange R, Boller T (1995) Rhizobial nodulation factors stimulate mycorrhizal colonization of nodulating and nonnodulating soybeans. *Plant Physiol* 108:1519–1525
- Zaat SAJ, Wijffelman CA, Mulders IHM, Van Brussel AAN, Lugtenberg BJJ (1988) Root exudates of various host plants of *Rhizobium leguminosarum* contain different sets of inducers of *Rhizobium* nodulation genes. *Plant Physiol* 86:1298–1303
- Zhang F, Smith DL (1995) Preincubation of *Bradyrhizobium japonicum* with genistein accelerates nodule development of soybean at suboptimal root zone temperatures. *Plant Physiol* 108:961–968
- Zuanazzi JAS, Clergeot PH, Quirion JC, Husson HP, Kondorosi A, Ratet P (1998) Production of *Sinorhizobium meliloti* nod gene activator and repressor flavonoids from *Medicago sativa* Roots. *Mol Plant-Microbe Interact* 11:784–794