

APPLIED RESEARCH

Alcohol beverages and biochemical changes in blood

SHELA GORINSTEIN¹ & SIMON TRAKHTENBERG²

¹Department of Medicinal Chemistry and Natural Products, The Hebrew University - Hadassah Medical School, Jerusalem, Israel & ²Kaplan University Medical Center, Rehovot, Israel

Abstract

Coronary atherosclerosis is the major cause of death in western civilization: one of every three in men as well as in women. It was shown that diets supplemented with moderate quantities of alcoholic beverages could lead to positive biochemical changes in blood of the consumers, which are regarded widely as indicators of improved prevention of atherosclerosis. This review summarizes the recent epidemiological, experimental and clinical investigations concerning mainly the plasma biochemical changes in lipid levels, anticoagulant and antioxidant activities.

Introduction

Atherosclerosis in general and coronary atherosclerosis in particular are still the most dangerous diseases in the industrial countries—the principal cause of death in western civilization.¹ Atherosclerosis is a multi-factorial process based on the action of various risk factors.² According to Ross,² the form and content of the advanced lesions of atherosclerosis demonstrate the results of the following fundamental biological processes: (a) accumulation of intimal smooth muscle cells, together with variable numbers of accumulated macrophages and T-lymphocytes; and (b) formation by the proliferated smooth muscle cells of large amounts of connective tissue matrix, principally in the form of cholesteryl esters and free cholesterol within the cells as well as in the surrounding connective tissues. In spite of the fact that some authors have found signs of inflammatory and immunological nature of

atherosclerosis,^{3–7} the oxidative theory is used widely as a basis for prevention and treatment of this disease.^{8–11} The oxidative theory states that only cholesterol-rich oxidized LDL-C particles are able to penetrate arterial walls causing their occlusions.^{8–11} Such pathological damage can be found practically in every artery.^{12,13} The most important damage is in the coronary arteries. Coronary artery disease (CAD) is a consequence of atherosclerotic process in these arteries. Occlusions of coronary arteries lead to myocardial infarctions, which in many cases can be fatal.¹⁴ Therefore, the international scientific community is searching for comprehensive schemes to fight this disease. Intensive multiple risk factor reduction has a beneficial effect on coronary atherosclerosis in men and women: the disease regression is twice as frequent in the risk reduction group as in the control group.¹⁵ An integral part of these preventive measures are

Correspondence to: Dr Shela Gorinstein, Department of Medicinal Chemistry and Natural Products, The Hebrew University of Jerusalem—Hadassah Medical School, POB 12065, Jerusalem 91120, Israel.
Tel: 972 6758690; Fax: 972 26757076; E-mail: gorin@cc.huji.ac.il

Received for publication 15th January 2003. Accepted 17th October 2003.

proper diets. The general principle of such diets is plenty of vegetables, fruit and cereal fibre.^{16,17} Alcoholic beverages have been food throughout the ages.^{18,19} In spite of the fact that even moderate consumption of alcoholic beverages leads to some structural changes in plasma circulating proteins,^{20,21} these beverages are an integral part of diets in most western countries and consist about 4–6% of the average energy intake.²² One of the reasons for widespread alcohol consumption is the claims of many investigators that alcoholic beverages have a cardioprotective effect.^{23–28} These authors have observed that consumption of alcoholic beverages influence positively the main risk factors of atherosclerosis. This review is based on recent epidemiological, experimental and clinical investigations concerning the influence of moderate consumption alcohol beverages on plasma lipid levels, plasma anticoagulant and plasma antioxidant activities.

Moderate alcohol consumption and plasma lipids

Since 1913 it has been an established fact that diet containing cholesterol creates atherosclerotic changes in the arteries of laboratory animals.²⁹ All authors currently agree that cholesterol is also the “building material” for atherosclerotic plaques in humans.^{30–32} Elevated levels of total cholesterol, low-density cholesterol (LDL-C), triglycerides, apolipoproteins B and C-III and reduced levels of high-density cholesterol (HDL-C) and apolipoprotein A-I are major risk factors for this disease.^{30–32}

Moderate consumption of alcoholic beverages leads first to an increase in HDL-cholesterol and has little effect on total and LDL-cholesterol.^{24–26} However, in experiments on laboratory animals it was demonstrated that diets supplemented with moderate quantities of alcoholic beverages led to decreased levels of plasma total cholesterol, LDL-C, triglycerides, total phospholipids and total cholesterol in liver and an increased level of HDL-cholesterol in animals fed added cholesterol.^{33–35} The same tendency was observed in clinical investigations of patients suffering from hypercholesterolemia: moderate consumption of alcoholic beverages led to a decrease in the level of total cholesterol, LDL-cholesterol, triglycerides and an increase of HDL-cholesterol.^{28,36}

Contrary to the above, increase of the levels of HDL-cholesterol after moderate alcoholic consumption is not disputable. An increase in the level of HDL-cholesterol was found in experiments on laboratory animals^{33–35} and in investigations of patients with and without hypercholesterolaemia.^{20,21,24,25,27,28,35–37} Recently, Sillanaukee *et al.*²⁴ have reported that HDL-cholesterol (the HDL₃ subfraction) was higher in those drinking 20–40 g alcohol daily. It was also found that the frequency of drinking was associated with a higher level of HDL-C.²⁵

It was shown that paraoxonase, an HDL-associated enzyme, is related to atherosclerosis.^{38–41} This organophosphate enzyme suppresses atherosclerosis in an unknown way.⁴¹ Fuhrman & Aviram³⁹ have shown that paraoxonase is associated physically with HDL, and its activity has been related inversely to the risk of cardiovascular diseases. These authors found that paraoxonase can hydrolyze specific lipid peroxides in oxidized lipoproteins and in atherosclerotic lesions. Fuhrman & Aviram³⁹ have demonstrated that consumption of wine flavonoids preserves paraoxonase activity by reducing the oxidative stress in apolipoprotein E-deficient mice, thereby contributing to paraoxonase hydrolytic activity on lipid peroxides in oxidized lipoproteins and atherosclerotic lesions. Paraoxonase inhibits macrophage cholesterol biosynthesis and atherogenesis probably through its phospholipase-A(2)-like activity.^{41,42} It was shown also in investigations of humans that paraoxonase activity is related inversely to the risk of cardiovascular diseases.³⁸ In this diet-controlled, randomized intervention study³⁸ the effects of daily moderate consumption of red wine, beer and spirits in comparison with mineral water on paraoxonase activity in serum was examined. For 3 weeks, eleven healthy middle-aged men each consumed three of the beverages (red wine, beer and spirits) with evening dinner. At the end of each 3-week period, blood samples were collected pre- and postprandially and after an overnight fast. Fasting paraoxonase activity was higher after intake of wine, beer and spirits than after water consumption ($p < 0.001$ in all three cases). The authors of this investigation suggest that increased serum paraoxonase may be one of the biological mechanisms underlying the reduced coronary heart disease risk in moderate alcohol consumers. Another study was performed to investigate the kinetics of alcohol-induced

increases in apo A-1, HDL cholesterol and paraoxonase activity.⁴³ This experiment was conducted with 30–40 g/day of beer. Serum apo A-I, HDL cholesterol and paraoxonase activity were increased significantly during 3 weeks of moderate alcohol consumption compared with no alcohol consumption. Moreover, the results suggest that there is a sequence in induction of these parameters. After an increase in apo A-1, HDL cholesterol is increased followed by an increase in paraoxonase activity. These authors⁴³ have concluded that the increased serum HDL cholesterol level and paraoxonase activity may be a mechanism of action not only in healthy middle-aged men but also in postmenopausal women, underlying the reduced coronary heart disease risk in moderate drinkers.

Moderate alcohol consumption and plasma antioxidant activity

As mentioned, some authors have found signs of the inflammatory and immunological nature of atherosclerosis.^{3–7,44} One of them even claims that atherosclerosis is a true inflammatory disease.⁶ However, after publication of the paper “Beyond cholesterol: modifications of low-density lipoprotein that increases its atherogenicity”,⁸ authors increasingly support the theory that atherosclerosis is an oxidative disease.^{8–11,40} Recently, Perez *et al.*⁴⁵ have proved that oxidative stress is a central mechanism for the pathogenesis of atherogenic heart disease and atherogenesis.

Some evidence suggests that one of the important mechanisms predisposing to development of atherosclerosis is the oxidation of cholesterol-rich LDL-C particles.^{9–11} Oxidation of LDL-C enhances its atherogenicity and facilitates penetration of lipids into the arterial wall.¹¹ Thus, according to some authors,^{8–11} prevention of atherosclerosis is a fight against LDL-C oxidation.

In most western industrial countries intake of food products containing high quantities of cholesterol is correlated positively with morbidity and mortality from coronary artery disease (CAD).²⁶ Total consumption of saturated fats in France is equal to that of other developed countries, while French mortality from CAD is only one-third of the average.^{26,45} This phenomenon is known as the French paradox for CAD.^{26,46,47} The distinguishing feature of the

French diet is regular consumption of red wine with meals. Therefore, it was supposed that regular consumption of red wine could explain why French mortality from CAD is only one-third of the average in other developed countries.^{26,46,47}

The antioxidant properties of wine were confirmed *in vitro*: Frankel *et al.*^{48,49} have shown that oxidation of human low-density lipoprotein is inhibited by the phenolic substances of red wine. The same results were received a year later by using not only wine but also beer phenolic substances.⁵⁰

In a very recent epidemiological study, a significant inverse relationship was found between oxidized LDL antibody titres and daily wine intake.⁵¹ In experiments on laboratory animals and in investigation of humans, it was also found that red wine, white wine and beer increase plasma antioxidant activity.^{28,33,34,52} However, the degree of this increase was different. The authors have explained the difference as follows. According to *in vitro* investigations, among the three widely used alcoholic beverages (red wine, white wine and beer) the highest content of total polyphenols is in red wine and the lowest in beer; this difference is significant ($p < 0.0025$).⁵³

However, the influence of diet, supplemented with beer on lipid metabolism and antioxidant activity is higher than of diets supplemented with white wine.^{33,34} It was found that these two beverages have different concentrations of some essential phenolics: the contents of procyanidins, epicatechin and ferulic acid are significantly higher in beer than in white wine ($p < 0.005$; $p < 0.05$ and $p < 0.0025$, respectively).⁵³ The high content of the above-mentioned phenolics is the most possible explanation of the higher biological activity of beer. This suggestion is supported by the findings of Paganga *et al.*⁵⁴ They have proved that the antioxidant activities of one glass of red wine (150 ml) is equivalent to 3.5 glasses of beer (500 ml) and equivalent to 12 glasses of white wine. Therefore, not only the level of total polyphenols play a role,⁵⁵ but also the content of some essential phenolics.⁵⁶

Among the three widely used alcoholic beverages (red wine, white wine and beer), red wine has the highest content of total polyphenols and the highest content of some essential phenolics. Therefore, some authors claim that red wine is the best choice for consumers.⁵⁷

Moderate alcohol consumption and plasma anticoagulant activity

Although not without exception, current evidence from epidemiological, experimental and clinical studies suggests a protective effect against the development of atherosclerosis with moderate consumption of alcoholic beverages. The exact nature of the protective effect remains to be established. However, mechanisms including alterations in haemostatic variables is being recognized increasingly as contributory.⁵⁸ Thrombosis of the coronary arteries is the main reason for fatal cases from myocardial infarctions due to coronary atherosclerosis.¹⁴ As shown, moderate consumption of alcoholic beverages lowers the risk and the death rate from cardiovascular diseases.^{59,60} Therefore, the decrease in the death rate could not be achieved without a beneficial influence on haemostatic risk factors.⁶¹

Many investigations have demonstrated that consumption of alcoholic beverages is correlated with haemostatic risk factors.^{27,62–68} It was found that platelet aggregation is inhibited by moderate alcohol consumption.^{26,63} According to Renaud & de Lorgeril,²⁶ this is the main reason for the low death rate from coronary atherosclerosis in France. It was also found in a clinical investigation that moderate beer consumption leads to a significant decrease in the prothrombin time.²⁷ Fibrinogen is one of the plasma circulating proteins. This protein is synthesized in liver and circulates in plasma at a concentration of 200–400 mg/dl. Fibrinogen plays an important role in blood clotting, fibrinolysis, cellular and matrix interactions. Fibrinogen is comprised of two sets of three polypeptide chains that are joined by disulfide bridging within the N-terminal E domain. These domains contain constitutive binding sites that participate in fibrinogen conversion to fibrin, fibrin assembly and platelet interactions.⁶⁹ Evidence links fibrinogen with coronary atherosclerosis and blood coagulation.⁶⁹ It was reported that fibrinogen levels predicted cardiovascular events independently of traditional risk factors.⁷⁰ Some authors have demonstrated that moderate drinking leads to a decrease in the plasma circulating fibrinogen concentration.^{71–73} Wang *et al.*⁷² have shown that daily consumption of moderate amounts of ethanol decreases circulating levels of fibrinogen by 18–20%. Also, some structural changes in plasma circulating fibrinogen after short-term moderate beer consumption were found using electrophoresis and

spectroscopy.³⁶ In order to evaluate the status of the plasma anticoagulant activity Factor VIIa, Factor VIIc and PAI were studied. A decrease was found in all these tests after moderate beer consumption.^{20,21,36} Another group of investigators have measured the levels of fibrinogen, plasma viscosity, von Willebrand factor, factor VII, plasminogen activator inhibitor antigen-1 and tissue plasminogen activator antigen in a cross-sectional analysis of 3223 enrolled in the Framingham Offspring Study.⁶⁷ They have found that light-to-moderate alcohol consumption is associated with lower levels of coagulatory activity, but higher intake is associated with impaired fibrinolytic potential.

All the above-mentioned epidemiological and clinical investigations have shown that moderate-only consumption of alcoholic beverages influences plasma anticoagulant activity favourably.

Moderate alcohol consumption and other biochemical changes

The positive influence of moderate alcohol consumption on plasma lipids, plasma antioxidant and plasma anticoagulant activity is well known. However, moderate consumption of alcoholic beverages leads to some other biochemical changes. It was found in a randomized, diet-controlled interventional study that moderate alcohol consumption reduces plasma C-reactive protein levels.⁷³ In another investigation it was demonstrated that red wine consumption influences positively preincubation of vascular smooth muscle cells, inhibits ligand binding and the subsequent tyrosine phosphorylation of the platelet-derived growth factor beta receptor, which plays a critical role in the pathogenesis of atherosclerosis.⁷⁴ There is evidence that consumption of alcoholic beverages influences positively synthase of nitric oxide, which plays a critical role in cardiovascular protection and is the responsible cardioprotective protein.⁷⁵ Also, Leikert *et al.*⁷⁶ have shown that red wine polyphenols increase endothelial nitric oxide synthase and subsequent endothelial nitric oxide release. Increased endothelial nitric oxide activity may antagonize the development of endothelial dysfunction and atherosclerosis, a hypothesis that supports the view that red wine may indeed have long-term protective cardiovascular properties mediated by its polyphenols.^{75,76}

The authors of the next investigation indicated that proliferation of vascular smooth muscle cells is critical to atherosclerosis formation.⁷⁷ They found that treatment with red wine polyphenols has a potent inhibitory effect on the proliferation and DNA syntheses in cultured rats. Not all authors have found a positive influence of moderate alcohol consumption on the iron stores in the body. Therefore, one group of investigators reported that wine consumption reduces iron stores and thereby the risk of coronary heart disease.⁷⁸ However, other authors have found that light drinking increases ferritin and, by inference, body iron stores.⁷⁹ Moderate beer consumption leads to an increase in plasma magnesium and as a consequence to a decrease in the heart arrhythmias.⁸⁰

Moderate alcohol consumption and the vascular wall functioning

In the well-known Rotterdam Study an inverse association between alcohol consumption and vascular wall functioning was found in non-smoking men and women.⁸¹ Failure to maintain an intact endothelium, as a result of episodic and persistent injury and perturbation of the vascular endothelium, promotes formation of fatty streaks, which are considered initiation events of atherosclerosis. Cellular constituents contributing to endothelial injury include endothelial cells, monocytes, platelets and smooth muscle cells.⁸² Evidence is emerging that resveratrol, a polyphenolic phytoalexin, which is present in dietary sources including red wine, may protect against atherosclerosis and cardiovascular disease by enhancing the integrity of the endothelium. The inhibitory effect of resveratrol on protein kinase C activity and therefore on the associated signalling networks may, in part, underline the mechanisms by which this agent exerts its beneficial effects on endothelial and cardiovascular function.⁸² It was shown that wine polyphenols induce vascular relaxation *in vitro* through the NO-cGMP pathway in rabbit blood vessels.⁸³ Zou *et al.*⁸³ claim that resveratrol of red wine may protect against atherosclerosis and cardiovascular disease by enhancing the integrity of the endothelium in experimental hypercholesterolemic rabbits. The authors found that administration of resveratrol, red wine or de-alcoholized red wine improved endothelial function, which in their opinion may be one of the mechanisms by which red wine

polyphenols exert their alcohol-independent cardioprotective effects. The effect of red wine and wine polyphenol resveratrol on endothelial function was also investigated in humans.^{84,85} It was demonstrated in a clinical study of healthy male volunteers that red wine counteracts endothelial dysfunction.⁸⁴ Cuevas *et al.*⁸⁴ have measured endothelial function as flow-mediated dilatation of the brachial artery, employing high-resolution ultrasound after an overnight fast. They have found that the deleterious effect on endothelial function by a high-fat diet was contracted by the protective role of wine. Other authors have also reported that wine in particular, but also beer, contains polyphenols, which act as antioxidants and could maintain the integrity of the endothelial function by reducing the formation of superoxide. Moreover, these antioxidants may protect against LDL oxidation and modulate the macrophage attack on the endothelium.^{86,87} The above-mentioned authors have concluded that red wine polyphenolic compounds can preserve a normal vascular reactivity by acting at different stages of the cascade that leads to lipid oxidation, endothelium dysfunction and vasospasm.

Discussion

It was shown that in most western industrial countries intake of food products containing high quantities of cholesterol is correlated positively with morbidity and mortality from CAD.² Total consumption of saturated fats in France is equal to that of other developed countries, while French mortality from CAD is only one-third of the average.^{26,46,47,74,77} The distinguishing feature of the French diet is regular consumption of red wine with meals. Therefore, it was supposed that regular consumption of red wine could explain the relatively low French mortality from CAD.^{26,47,74,77} Most of the authors cited in this review have shown that moderate consumption of alcoholic beverages is cardioprotective because of the positive influence on plasma lipids, plasma antioxidant and plasma anticoagulant activity and vascular wall function.^{18,88,89} The most comprehensive study to date summarizes the results of alcohol consumption and mortality among middle-aged and elderly American adults.¹⁹ Among 490 000 men and women who used alcohol beverages, 46 000 died during 9 years of follow-up. The authors have compared cause-specific death rates and death from all causes across

categories of baseline alcohol consumption. They found that the rates of death from all cardiovascular diseases between the ages of 35 and 69 were 30–40% lower among men and women reporting one drink daily than among non-drinkers.

Currently, all investigators agree that diets supplemented with moderate quantities of alcoholic beverages lead to biochemical changes, which are regarded widely as indicators of improved prevention of atherosclerosis.^{23–25} However, there is ongoing debate about the bioactive components of the alcoholic beverages which lead to this effect.

Some authors claim that ethanol itself is responsible for the cardioprotective effect of alcoholic beverages.^{61,72,89} In their case-control study, Brenner *et al.*⁶¹ have assessed and compared the effect of alcohol consumption from various sources on risk of coronary heart disease. Their sample included 312 patients with clinically stable, angiographically confirmed coronary heart disease and 479 healthy controls. The results of their investigation concluded that the protective effect of moderate alcohol consumption against coronary heart disease is mediated in part by beneficial effects of ethanol on lipids and haemostatic factors. Wang *et al.*⁷² have shown that daily consumption of moderate amounts of ethanol decreases circulating levels of fibrinogen by 18–20%. However, many other authors have proved that wine polyphenols are responsible for the cardioprotective effect of alcoholic beverages.^{48,50,90–98} Frankel *et al.*⁴⁸ have reported that phenolic substances of red wine inhibit oxidation of human low-density lipoprotein. Mosinger⁵⁰ observed that polyphenols but not alcohol in beer and wine protect serum low-density lipoprotein against atherogenic modifications. In subsequent years a number of experiments on laboratory animals were performed.^{33,34,91–93} Gorinstein *et al.* performed two experiments on rats: with dry matter of alcoholic beverages³³ and with alcohol-containing and alcohol-free different alcoholic beverages.³⁴ These authors reported that both original and lyophilized wine and beer increase antioxidant activity and improve lipid metabolism. There were no statistically significant differences in the results of the investigations performed on groups of rats fed basal diet supplemented with original wine and beer versus groups fed basal diet supplemented with

lyophilized wine and beer. In a well-performed experiment on hypercholesterolaemic golden Syrian hamsters it was found that red wine phenolic compounds reduce plasma lipids and apolipoprotein B and prevent early aortic atherosclerosis.⁹³ Similar results were obtained in investigations on humans.^{94–98} Fuhrman *et al.*⁹⁴ have demonstrated that consumption of red wine with meals reduces the susceptibility of human plasma and low density lipoprotein to lipid peroxidation. Serafini *et al.*⁹⁵ found that alcohol-free red wine enhances plasma antioxidant capacity. Carbonneau *et al.*⁹⁶ observed an improvement in plasma antioxidant status and in LDL-C level of subjects receiving a red wine phenolic mixture. Aviram & Fuhrman⁹⁸ proved that wine flavonoids protect against LDL oxidation and atherosclerosis. The above-mentioned experiments on laboratory animals and investigations of humans underline the role of phenolics in alcoholic beverages.

There is also a “compromise” opinion: both ethanol and phenolic substances of alcoholic beverages positively affect plasma lipid levels. In an already-mentioned experiment⁹³ the effects of a red wine phenolic extract and ethanol on plasma lipoproteins and early atherosclerosis were studied in hamsters. The aortic fatty streak area was reduced significantly in both groups receiving phenolic extract and ethanol (the later reduced the aortic fatty streak area by 60%). As far back as 1979, Leger *et al.*⁸⁸ reported a strong and specific negative association between CAD deaths and alcohol consumption of wines, beer and spirits. The report by Leger *et al.*⁸⁸ was supported by the results of the investigation by Goldberg *et al.*,⁹⁹ who found phenolic constituents even in distilled spirits. In 2000, van der Gaag *et al.*¹⁰⁰ claimed that the effects “are largely independent of the type of alcoholic beverage”. This review cited many epidemiological, experimental and clinical investigations which proved that moderate consumption of alcoholic beverages is cardioprotective.

However, it must be underlined that some investigators strongly oppose the use of alcohol beverages for general population.^{101,102} Llerena *et al.*¹⁰¹ claim that it has not been demonstrated that alcohol intake, even in moderate amounts, is beneficial for the general population, in particular men under the age of 40 and women under 50,

because it raises mortality due to other causes, especially injury, cirrhosis of the liver and some types of cancer, thereby outweighing the benefits for coronary artery disease. Thus, alcohol consumption should not be recommended as a prophylaxis for the general population. The same opinions are shared by Corrao *et al.*¹⁰² They have found that the estimated number of deaths attributed to alcohol use in Italy still exceeds by far the number of deaths prevented by alcohol use both for men and women.

In conclusion, (1) current evidence from the epidemiological, experimental and clinical studies shows that moderate consumption of alcoholic beverages leads to positive biochemical changes, which is believed widely to have a protective effect against the development of atherosclerosis. (2) Based on majority of the reviewed papers it can be concluded that the main biologically active compounds of alcoholic beverage are phenolics and ethanol. (3) Red wine has the highest content of total polyphenols and essential phenolics. Therefore, among alcoholic beverages red wine could be the best choice.

References

- Gaziano JM. Antioxidant vitamins and coronary artery disease. *Am J Med* 1994;97:18S–21, 22S–8.
- Ross R. The pathogenesis of atherosclerosis. In: Braunwald E, editor. *Heart Diseases*. Philadelphia: W. B. Saunders Co.; 1997, pp. 1105–21.
- Yla-Herttuala S, Palinsky W, Butler S, Picard S, Steinberg D, Witztum JL. Rabbit and human atherosclerotic lesions contain IgG that recognized epitopes of oxidized low density lipoproteins. *Atheroscler Thromb* 1994;14:32–40.
- Frostgard J, Wu R, Guscombe E. Induction of T cell activation by oxidized low density lipoprotein. *Atheroscler Thromb* 1992;12:461–7.
- Watanabe T, Haraoka S, Shimokama T. Inflammatory and immunological nature of atherosclerosis. *Int J Cardiol* 1996;54:S51–60.
- Ross R. Atherosclerosis is an inflammatory disease. *Am Heart J* 1999;138:S419–20.
- Wick G, Pershinka H, Qingbo Xu. Autoimmunity and atherosclerosis. *Am Heart J* 1999;138:S444–9.
- Steinberg D, Parthasarathy S, Carew T, Khoo J, Witztum J. Beyond cholesterol: modifications of low density lipoprotein that increases its atherogenicity. *N Engl J Med* 1989;320:915–24.
- Steinbrecher UP, Zhang H, Londheed M. Role of oxidativity modified LDL in atherosclerosis. *Free Radic Biol Med* 1990;9:155–68.
- Witztum JL, Steinberg D. Role of oxidized low density lipoprotein in atherogenesis. *J Clin Invest* 1991;88:1785–92.
- Aviram M. Modified forms of low density lipoprotein and atherosclerosis. *Atherosclerosis* 1993;98:1–9.
- Genest J, McNamara JR, Ordow MJM *et al.* Lipoprotein cholesterol, apolipoprotein A-I and B lipoprotein abnormalities in men with premature coronary artery disease. *J Am Coll Cardiol* 1992;19:792–802.
- Hodis HN, Mack WJ, Azen SP *et al.* Triglyceride- and cholesterol-rich lipoproteins have a different effect on mild/moderate and severe lesion progression as assessed by quantitative coronary angiography in a controlled trial of lovastatin. *Circulation* 1994;90:42–9.
- Brosins FC III, Roberts WS. Significance of coronary artery thrombosis in transmural acute myocardial infarction. A study of 54 necropsy patients. *Circulation* 1983;63:810–6.
- Haskell W, Alderman E, Fair J and the SCRIP Investigators. Intensive multiple-risk-factor reduction to modify coronary atherosclerosis in men and women with coronary artery disease. In: Woodford FP, Davignon J, Sniderman A, editors. *Atherosclerosis X*. Amsterdam: Elsevier, 1995; pp. 565–70.
- Rimm EB, Ascherio A, Giovannucci E, Spiegelman D, Stampfer M, Willett W. Vegetable, fruits, and cereal fiber intake and risk of coronary heart disease among men. *JAMA* 1996;275:447–51.
- Rimm EB, Katan MB, Ascherio A, Stampfer MJ, Willett W. Relation between intake of flavonoids and risk for coronary heart disease in male health professionals. *Ann Intern Med* 1996;125:384–9.
- Rimm EB, Giovannucci E, Willett WC *et al.* Prospective study of alcohol consumption and risk of coronary disease in men. *Lancet* 1991;338:464–8.
- Thun MJ, Peto R, Lopez AD *et al.* Alcohol consumption and mortality among middle-aged and elderly US adults. *N Engl J Med* 1997;337:1456–8.
- Gorinstein S, Caspi A, Libman I *et al.* Intrinsic tryptophan fluorescence of human serum proteins and related conformational changes. *Protein Chem* 2000;19:637–42.
- Gorinstein S, Caspi A, Goshev I *et al.* Beer consumption and changes in stability of human serum proteins. *J Agric Food Chem* 2001; 49:1441–5.
- Christiansen C, Thomsen C, Rasmussen D *et al.* Effect of alcohol on glucose, insulin free fatty acids and triacylglycerol responses to a light meal in non insulin dependent diabetic subjects. *Br J Nutr* 1994;71:449–54.
- Friedman LA, Kinball AW. Coronary artery disease and alcohol consumption in Framingham. *Am J Epidemiol* 1986;124:481–9.
- Sillanaukee P, Koivula T, Jokela H, Pitkääjärvi T, Seppä K. Alcohol consumption and its relation to lipid based cardiovascular risk factors among middle-aged women: the role of HDL3 cholesterol. *Atherosclerosis* 2000;152:503–10.

25. Hashimoto Y, Futamura A, Nakarai H, Nakahara K. Effect of the frequency of alcohol intake on risk factors for coronary heart disease. *Eur J Epidemiol* 2001;17:307–12.
26. Renaud S, de Lorgeril M. Wine, alcohol, platelets and the French paradox for coronary heart disease. *Lancet* 1992;339:1523–6.
27. Gorinstein S, Zemser M, Lichman I *et al.* Moderate beer consumption and the blood coagulation in patients with coronary atherosclerosis. *J Intern Med* 1997;241:47–51.
28. Gorinstein S, Zemser M, Berliner M *et al.* Moderate beer consumption and positive biochemical changes in patients with coronary artery disease. *J Intern Med* 1997;242:219–24.
29. Anitschkow N, Chalataw S. Uber experimentelle Cholesterinsteatose und ihre Bedeutung fur die Entstehung einiger pathologischer Prozesse. *Centrbl Allg Pathol Anat* 1913;24:1–9.
30. Castelli WB, Garrison RJ, Wilson PWF, Abbott RD, Kalousdian S, Kannel WB. Incidence of coronary heart disease and lipoprotein cholesterol levels. The Framingham Study. *JAMA* 1986;256:2835–8.
31. Genest J, McNamara JR, Ordow MJM *et al.* Lipoprotein cholesterol, apolipoprotein A-I and B lipoprotein abnormalities in men with premature coronary artery disease. *J Am Coll Cardiol* 1992;19:792–802.
32. Kwiterovich PO, Coresh J, Smith HH, Bachoric PS, Derby CA, Pearson TA. Comparison of plasma level of apolipoprotein B and A-1 and other risk factors in men and in women with premature coronary artery disease. *Am J Cardiol* 1992;69:1015–21.
33. Gorinstein S, Zemser M, Weisz M, Haruenkit R, Trakhtenberg S. The influence of dry matter of different alcoholic beverages on lipid and protein metabolism and antioxidant activity in serum of rats. *J Nutr Biochem* 1998;9:131–5.
34. Gorinstein S, Zemser M, Weisz M, Halevy SH, Martin-Bellosso O, Trakhtenberg S. The influence of alcohol-containing and alcohol-free beverages on lipid levels and lipid peroxides in serum of rats. *J Nutr Biochem* 1998;9:682–6.
35. Gorinstein S, Caspi A, Pawelzik E *et al.* Protein of beer affect lipid levels in rats. *Nutr Res* 2001;21:1159–69.
36. Gorinstein S, Caspi A, Goshev I *et al.* Structural changes in plasma circulating fibrinogen after moderate beer consumption as determined by electrophoresis and spectroscopy. *J Agric Food Chem* 2003;51:822–7.
37. Nakanishi N, Yoshida K, Kawashimo H, Tatara K. Influence of alcohol intake on risk for increased low-density lipoprotein cholesterol in middle-aged Japanese men. *Alcohol Clin Exp Res* 2001;25:1046–50.
38. Van der Gaag MS, van Tol A, Scheek LM *et al.* Daily moderate alcohol consumption increases serum paraoxonase activity; a diet-controlled, randomised intervention study in middle aged men. *Atherosclerosis* 1999;147:405–10.
39. Fuhrman B, Aviram M. Preservation of paraoxonase activity by wine flavonoids—possible role in protection of LDL from lipid peroxidation. *Alcohol and wine in health and disease. Ann NY Acad Sci* 2002;957:321–4.
40. Bin Ali A, Zhang Q, Lim YK, Fang D, Retnam L, Lim SK. Expression of major HDL-associated antioxidant PON-1 is gender dependent and regulated during inflammation. *Free Radic Biol Med* 2003;34:824–9.
41. Rozenberg O, Shih DM, Aviram M. Human serum paraoxonase 1 decreases macrophage cholesterol biosynthesis—possible role for its phospholipase-A(2)-like activity and lysophosphatidylcholine formation. *Ather Thromb Vasc Biol* 2003;23:461–7.
42. Marathe GK, Zimmerman GA, McIntyre TM. Platelet-activating factor acetylhydrolase, and not paraoxonase-1, is the oxidized phospholipid hydrolyase of high density lipoprotein particles. *J Biol Chem* 2003;278:3937–47.
43. Sierksma A, van der Gaag MS, van Tol A, James RW, Hendriks HFJ. Kinetics of HDL cholesterol and paraoxonase activity in moderate alcohol consumers. *Alcohol Clin Exp Res* 2002;26:1430–35.
44. Imhof A, Froelich M, Brenner H, Boeing H, Pepys MB, Koenig W. Effect of alcohol consumption on systemic markers of inflammation. *Lancet* 2001;357:763–7.
45. Perez DD, Strobel P, Foncea R *et al.* Wine, diet, antioxidant defenses, and oxidative damage. *Alcohol and wine in health and disease. Ann NY Acad Sci* 2002;957:136–45.
46. Stanley LL, Mazier MJP. Potential explanations for the French paradox. *Nutr Res* 1999;19:3–15.
47. Belleville J. The French paradox: possible involvement of ethanol in the protective effect against cardiovascular diseases. *J Nutr* 2002;18:173–7.
48. Frankel EN, Kanner J, German GB, Parks E, Kinsella JE. Inhibition of oxidation of human low-density lipoprotein by phenolic substances in red wine. *Lancet* 1993;341:454–7.
49. Frankel EN, Waterhouse AL, Kinsella JE. Inhibition of human LDL-C oxidation by resveratrol. *Lancet* 1993;341:1103–4.
50. Mosinger B. Polyphenolics but not alcohol in beer and wine protect serum low-density lipoprotein against atherogenic modification. *Cor Vasa* 1994;4:171–4.
51. Di Bari M, Zacchei S, Kritchevsky SB *et al.* Antioxidized LDL antibodies and wine consumption: a population-based epidemiological study in Dicomano, Italy. *Ann Epidemiol* 2003;13:189–95.
52. Gorinstein S, Caspi A, Libman I *et al.* Positive and negative effects of short-term moderate beer consumption. *Alcohol Res* 2001;6:233–9.
53. Gorinstein S, Caspi A, Zemser M, Trakhtenberg S. Comparative contents of some phenolics in beer, red and white wines. *Nutr Res* 2000;1:131–9.
54. Paganga G, Miller N, Rice-Evans CA. The polyphenolic content of fruit and vegetables and their antioxidant activities. What does a serving constitute? *Free Radic Res* 1999;30:153–62.

55. Abu-Amsha R, Croft KD, Puddey IB, Proudfoot JM, Beilin LJ. Phenolic content of various beverages determines the extent of inhibition of human serum and low-density lipoprotein oxidation *in vitro*: identification and mechanism of action of some cinnamic acid derivatives from red wine. *Clin Sci* 1996;91:449–58.
56. Rifici VA, Schneider SH, Khachadurian AK. Lipoprotein oxidation mediated by J774 murine macrophages is inhibited by individual red wine polyphenols but not by ethanol. *J Nutr* 2002;132:2532–7.
57. Klatsky AL, Armstrong MA. Alcohol beverages choice and risk of coronary artery disease. Mortality: do red wine drinkers fare best? *Am J Cardiol* 1993;71:467–9.
58. Wollin SD, Jones PJH. Alcohol, red wine and cardiovascular disease. *J Nutr* 2002;132:1401–4.
59. Sesso HD, Stamfer MJ, Rosner B, Hennekens CH, Manson JE, Gaziano JM. Seven-year changes in alcohol consumption and subsequent risk of cardiovascular disease in men. *Arch Intern Med* 2000;160:2605–12.
60. Wannamethee SG, Shaper AG. Taking up regular drinking in middle age: effect on major coronary heart disease event and mortality. *Heart* 2002;87:32–6.
61. Dimmitt SB, Rakic V, Puddey IB *et al.* The effects of alcohol on coagulation and fibrinolytic factors: a controlled trial. *Blood Coagul Fibrinol* 1998;9:39–45.
62. Lacoste L, Hung J, Lam JYT. Acute and delayed antithrombotic effects of alcohol in humans. *Am J Cardiol* 2001;87:82–5.
63. Mezzano D, Leighton F, Martinez C *et al.* Complementary effects of Mediterranean diet and moderate red wine intake on haemostatic cardiovascular risk factors. *Eur J Clin Nutr* 2001;55:444–51.
64. Brenner H, Rothenbacher D, Bode G, Marz W, Hoffmeister A, Koenig W. Coronary heart disease risk reduction in a predominantly beer-drinking population. *Epidemiology* 2001;12:390–5.
65. Sierksma A, van der Gaag MS, Schaafsma G, Kluff C, Bakker M, Hendriks FHJ. Moderate alcohol consumption and fibrinolytic factors of pre- and post-menopausal women. *Nutr Res* 2001;21:171–81.
66. Marquez-Vidal P, Montaye M, Haas B *et al.* Relationship between alcoholic beverages and cardiovascular risk factor levels in middle-aged men: the Prime study. *Atherosclerosis* 2001;157:431–40.
67. Mukamal KJ, Jadhav PP, D'Agostino RB *et al.* Alcohol consumption and hemostatic factors. Analysis of the Framingham Offspring cohort. *Circulation* 2001;104:1367–73.
68. Buemann B, Dyerberg J, Astrup A. Alcohol drinking and cardiac risk. *Nutr Res Rev* 2002;15:91–121.
69. Mosesson MW, Siebenlist KR, Meh DA. The structure and biological features of fibrinogen and fibrin. *Ann NY Acad Sci* 2001;936:11–30.
70. Palmieri V, Celentano A, Roman MJ de *et al.* Relation of fibrinogen to cardiovascular events is independent of preclinical cardiovascular disease: the Strong Heart Study. *Am Heart J* 2003;145:467–74.
71. Mennen LI, Balkau B, Vol S, Caces E, Eschwege E. Fibrinogen—a possible link between alcohol consumption and cardiovascular disease? *Arterioscler Thromb Vasc Biol* 1999;19:887–92.
72. Wang ZQ, Baker TH, Fuller GM. Alcohol at moderate levels decreases fibrinogen expression *in vivo* and *in vitro*. *Alcohol Clin Exp Res* 1999;23:1927–32.
73. Sierksma A, van der Gaag WS, Kluff S, Hendriks HFJ. Moderate alcohol consumption reduces plasma C-reactive protein and fibrinogen levels; a randomized, diet-controlled intervention study. *Eur J Clin Nutr* 2002;56:1130–6.
74. Rosenkranz S, Knirel D, Dietrich H, Flesch M, Erdmann E, Bohm M. Inhibition of the PDGF receptor by red wine flavonoids provides a molecular explanation for the “French paradox”. *FASEB J* 2002;16:U192–211.
75. Parks DA, Booyse FM. Cardiovascular protection by alcohol and polyphenols—role of nitric oxide. Alcohol and wine in health and disease. *Ann NY Acad Sci* 2002;957:115–21.
76. Leikert JF, Rathel TR, Wohlfart P, Cheynier V, Vollmar AM, Dirsch VM. Red wine polyphenols enhance endothelial nitric oxide synthase expression and subsequent nitric oxide release from endothelial cells. *Circulation* 2002;106:1614–17.
77. Iijima K, Yoshizumi M, Ouchi Y. Effect of red wine polyphenols on vascular smooth muscle cell function—molecular mechanism of the “French paradox”. *Mech Ageing Dev* 2002;123:1033–9.
78. De Lorgeril M, Salem P, Boucher F, de Leiris J, Paillard F. Effect of wine ethanol on serum iron and ferritin levels in patients with coronary heart disease. *Nutr Metab Cardiovasc Dis* 2001;11:176–80.
79. Whitfield JB, Zhu G, Heath AC, Powell LW, Martin NG. Effects of alcohol consumption on indices of iron stores and iron on alcohol intake markers. *Alcohol Clin Exp Res* 2001;25:1037–45.
80. Gorinstein S, Zenser M, Libman I, Trakhtenberg S, Caspi A. Effect of beer consumption on plasma magnesium: randomized comparison with mineral water. *J Roy Soc Med* 1998;91:631–3.
81. Vliegthart R, Geleijnse JM, Hofman A *et al.* Alcohol consumption and risk of peripheral arterial disease—the Rotterdam Study. *Am J Epidemiol* 2002;155:332–8.
82. Slater SJ, Seiz JL, Cook AC, Stagliano BA, Buzas CJ. Inhibition of protein kinase C by resveratrol. *Biochim Biophys Acta—Mol Basis Dis* 2003;1637:59–69.
83. Zou JG, Wang ZR, Huang YZ, Cao KJ, Wu JM. Effect of red wine and wine polyphenol resveratrol on endothelial function in hypercholesterolemic rabbits. *Int J Mol Med* 2003;11:317–20.

84. Cuevas AM, Guasch V, Castillo O *et al.* A high-fat diet induces and red wine counteracts endothelial dysfunction in human volunteers. *Lipids* 2000;35:143–8.
85. Wu JM, Wang ZR, Hsieh TC, Bruder JL, Zou JG, Huang YZ. Mechanism of cardioprotection by resveratrol, a phenolic antioxidant present in red wine. *Int J Mol Med* 2001;8:3–17.
86. Buemann B, Dyerberg J, Astrup A. Alcohol drinking and cardiac risk. *Nutr Res Rev* 2002;15:91–121.
87. Deckert V, Desrumaux C, Athias A *et al.* Prevention of LDL alpha-tocopherol consumption, cholesterol oxidation, and vascular endothelium dysfunction by polyphenolic compounds from red wine. *Atherosclerosis* 2002;165:41–50.
88. Leger AS, Cochrane AL, Moore I. Factors associated with cardiac mortality in developed countries with particular reference to the consumption of wine. *Lancet* 1979;1:1017–20.
89. Moor RD, Pearson TA. Moderate alcohol consumption and coronary disease. *Medicine* 1986;65:246–67.
90. Carando S, Teissedre PL, Leger C, Cabanis JC. Wine, beer and spirits consumption and cardiovascular diseases. *Sci Aliments* 1998;18:115–27.
91. Cestaro B, Simonetti P, Cervato G, Brusomolino A, Gatti P, Testolin G. Red wine effects on peroxidation indexes of rat plasma and erythrocytes. *Int J Food Sci Nutr* 1996;47:181–9.
92. Cascon E, Roig R, Ardevol A, Salvado MJ, Arola L, Blade C. Nonalcoholic components in wine reduce low density lipoprotein cholesterol in normocholesterolemic rats. *Lipids* 2001;36:383–8.
93. Auger C, Caporiccio B, Landrault N *et al.* Red wine phenolic compounds reduce plasma lipids and apolipoprotein B and prevent early aortic atherosclerosis in hypercholesterolemic golden Syrian hamsters (*Mesocricetus auratus*). *J Nutr* 2002;132:1207–13.
94. Fuhrman D, Lavy A, Aviram M. Consumption of red wine with meals reduces the susceptibility of human plasma and low density lipoprotein to lipid peroxidation. *Am J Clin Nutr* 1995;61:549–54.
95. Serafini M, Maiani G, Ferro-Luzzi A. Alcohol free red wine enhances plasma antioxidant capacity in humans. *J Nutr* 1998;128:1003–7.
96. Carbonneau MA, Leger CL, Descomps B, Michel F, Monnier L. Improvement in the antioxidant status of plasma and low-density lipoprotein in subjects receiving a red wine phenolics mixture. *J Am Oil Chem Soc* 1998;75:235–40.
97. Ghiselli A, Natella F, Guidi A, Montanari L, Fantozzi P, Scaccini C. Beer increases plasma antioxidant capacity in humans. *J Nutr Biochem* 2000;11:76–80.
98. Aviram M, Fuhrman B. Wine flavonoids protect against LDL oxidation and atherosclerosis. *Alcohol and wine in health and disease. Ann NY Acad Sci* 2002;957:146–61.
99. Goldberg DM, Hoffman B, Yang J, Soleas GJ. Phenolic constituents, furans, and total antioxidant status of distilled spirits. *J Agric Food Chem* 1999;47:3978–85.
100. Van der Gaag MS, van den Berg R, van den Berg H, Schaafsma G, Hendriks HFJ. Moderate consumption of beer, red wine and spirits has counteracting effects on plasma antioxidants in middle-aged men. *Eur J Clin Nutr* 2000;54:586–91.
101. Llerena IR, Huerta EM. Wine and the heart. *Rev Esp Cardiol* 1998;51:435–49.
102. Corrao G, Rubbiati L, Zambon A, Aricò S. Alcohol-attributable and alcohol-preventable mortality in Italy: a balance in 1983 and 1996. *Eur J Public Health* 2002;12:214–23.

Copyright of Addiction Biology is the property of Carfax Publishing Company and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.