

A Research Note

Amino Acid Composition and Its Change in Raw and Granulated Potatoes During Processing

S. GORINSTEIN, S. YAMAGATA, and D. HADZIYEV

ABSTRACT

Total amino acids and composition of raw and cooked tubers were determined by gas-liquid and ion-exchange column chromatography. Peeled raw potato, after reaching the end product state, lost 40% of its free amino acid pool. Processing into dehydrated granules by the freeze-thaw process caused only small losses in total amino acids and little change in their composition. There was only a 4.5% destruction of essential amino acids and a total loss of 2.1% of non-essential amino acids, excluding aspartic and glutamic acids which comprised 48% of peeled and sliced potatoes.

INTRODUCTION

THE EFFECTS of peeling and cooking on the amino acids of potatoes has been examined earlier (Augustin et al., 1979; Davies, 1977; Davies and Thomas, 1973; Eppendorfer et al., 1979; Fitzpatrick et al., 1964; Kozempel et al., 1982; Talley and Porter, 1968; Talley et al., 1970, 1983; Toma et al., 1978; Yamagata, 1983). The losses in amino acids of dehydrated granulated potatoes prepared by the freeze-thaw process were investigated in this study.

MATERIALS & METHODS

APPROXIMATELY 4.3 kg medium-size potato tubers (*Solanum tuberosum*, cv. Russet Burbank) grown in Alberta and obtained from I & S Produce Ltd. (Edmonton) were washed, peeled and sliced.

Granule preparation by the freeze-thaw (F-T) process developed by Ooraikul (1977) was used for the preparation of dehydrated potato granules. Samples were withdrawn from each stage indicated by an asterisk in Fig. 1 and then freeze-dried.

In order to follow the extent of destruction of labile amino acids (half-cystine, methionine, tryptophan and tyrosine), three different hydrolytic procedures (hydrolysis with HCL with a preliminary oxidation step (Blackburn, 1978); HCL without preliminary oxidation (Yamagata, 1983); and with methanesulfonic acid) were used for comparison studies for each sample (Liu and Chang, 1971; Simpson et al., 1976). Raw and processed potatoes were analyzed for amino acids by ion-exchange column chromatography (Blackburn, 1978; Moore et al., 1958; Moore and Stein, 1954; Rosen, 1957; Spackman et al., 1958; Syngde, 1977) and simultaneously gas-liquid chromatography (GLC), (Golan-Goldhirsh and Wolfe, 1979; Golan-Goldhirsch et al., 1982; Kirkman et al., 1980; Tajima et al., 1978).

In this study all statistical data were determined by Duncan's (1955) Multiple Range Test.

RESULTS & DISCUSSION

THE AMINO ACID COMPOSITION of dehydrated potato granules obtained by a semi-pilot scale freeze-thaw process is given in Table 1. The data were obtained by applying gas-liquid and ion-exchange chromatography methods and were arranged by the unit operations of the process.

As revealed by GLC, the peeled tuber contained 90 mg of apparent amino acids/g dry potato. However, when the amino




Fig. 1—Flow chart of the freeze-thaw process.

acids half-cystine, serine, isoleucine, leucine and histidine were included, the total amino acid content was raised to 104.5 mg/g dry potato. This decreased ($p < 0.05$) to 90.1 units after steam-cooking, 75.7 units after mashing, 74.2 units after freeze-thawing and 73.7 after granulation. Then end product content was 58.3 units and the discard, 71.6 units.

When ion-exchange chromatography data were analyzed, sliced samples provided 101.1 mg/g dry potato. This decreased ($p < 0.005$) to 92.6 units after steam-cooking, 91.6 units after mashing, 90.7 units after freeze-thawing and 85.6 units after granulation. The end product content was 84.5 units.

Total amino acids as found by GLC were either too low or the results obtained by ion-exchange chromatography were inflated (even though in the latter case the contents of γ -amino-butyric acid and ornithine were not included). The discrepancy between the two methods was tolerable only in slicing and steam-cooking steps but was too high in other steps (15.9 mg amino acid/g dry potato in mashing; 16.5 mg amino acid/g dry potato in freeze-thawing; 11.9 mg amino acid/g dry potato in granulation and 26.2 mg amino acid/g dry potato in end product.)

To resolve the discrepancy, the total crude protein-N content was followed during the granule process. The results (Table 1) revealed a steam-cook-induced loss of 4.6%, the loss ($p < 0.05$) increasing by 1.5% in mashing, 0.5% in freeze-thawing and 5.18% in granulation, giving a total loss of 11.78% total crude protein-N. The significant increase ($p < 0.05$) in the end product was only 1.1%. These results supported the ion-

Author Gorinstein is with the Dept. of Pharmaceutical Chemistry, The Hebrew Univ. of Jerusalem, Israel. Authors Yamagata and Hadziyev are with the Dept. of Food Science, Edmonton, The Univ. of Alberta, Canada.

Table 1—Total amino acid composition of peeled potato tubers cv Russet Burbank (Alberta) processed into dehydrated freeze-thaw granules, using ion-exchange and gas-liquid chromatography methods

Amino acid		Amino acid content mg/g dry potato ^a freeze-thaw processing step						
		Slicing	Steaming	Hot mashing	Thawing	Granulation	End product	Discard portion
Aspartic acid	a ^b	28.32 ± 0.28	26.14 ± 0.24	26.39 ± 0.31	27.31 ± 0.45	22.28 ± 0.33	21.79 ± 0.36	24.26 ± 0.40
	b	28.19 ± 0.34	20.07 ± 0.34	19.06 ± 0.37	18.87 ± 0.34	18.82 ± 0.42	17.99 ± 0.41	19.26 ± 0.39
Threonine ^c	a	3.49 ± 0.47	3.23 ± 0.57	3.18 ± 0.41	3.07 ± 0.39	3.05 ± 0.47	3.01 ± 0.48	2.79 ± 0.54
	b	2.98 ± 0.65	2.80 ± 0.71	2.56 ± 0.49	2.54 ± 0.37	0.90 ± 0.38	2.16 ± 0.55	2.21 ± 0.47
Serine	a	3.49 ± 0.81	3.42 ± 0.60	3.15 ± 0.74	3.31 ± 0.44	2.99 ± 0.42	2.95 ± 0.52	2.75 ± 0.51
	b	—	—	—	—	—	—	—
Glutamic acid	a	19.79 ± 0.39	16.42 ± 0.41	17.41 ± 0.39	16.47 ± 0.37	16.61 ± 0.34	16.46 ± 0.33	15.56 ± 0.39
	b	17.34 ± 0.36	18.18 ± 0.30	14.21 ± 0.39	14.07 ± 0.31	14.28 ± 0.36	14.00 ± 0.36	14.52 ± 0.40
Proline	a	3.03 ± 0.42	2.56 ± 0.39	2.72 ± 0.27	2.25 ± 0.29	2.61 ± 0.35	2.56 ± 0.33	2.06 ± 0.36
	b	3.13 ± 0.34	3.10 ± 0.34	1.84 ± 0.37	1.82 ± 0.24	2.38 ± 0.29	1.53 ± 0.27	1.70 ± 0.22
Glycine	a	3.07 ± 0.51	2.77 ± 0.47	2.94 ± 0.42	2.86 ± 0.42	2.47 ± 0.33	2.49 ± 0.39	2.19 ± 0.37
	b	3.21 ± 0.44	2.88 ± 0.46	2.80 ± 0.41	2.77 ± 0.39	1.66 ± 0.37	1.66 ± 0.41	2.13 ± 0.42
Alanine	a	3.13 ± 0.36	2.90 ± 0.40	2.65 ± 0.37	2.69 ± 0.38	2.62 ± 0.39	2.55 ± 0.40	2.41 ± 0.31
	b	3.23 ± 0.39	2.74 ± 0.31	2.56 ± 0.36	2.55 ± 0.33	1.98 ± 0.30	1.97 ± 0.31	2.23 ± 0.31
1/2 Cystine ^c	a	1.08 ± 0.24	0.86 ± 0.26	0.92 ± 0.19	0.43 ± 0.21	1.03 ± 0.24	1.08 ± 0.29	0.81 ± 0.21
	b	—	—	—	—	—	—	—
Valine ^c	a	5.18 ± 0.69	4.91 ± 0.44	4.79 ± 0.46	4.66 ± 0.31	4.58 ± 0.47	4.52 ± 0.50	4.32 ± 0.34
	b	3.79 ± 0.55	4.49 ± 0.50	4.29 ± 0.43	4.25 ± 0.42	1.86 ± 0.42	2.84 ± 0.37	4.12 ± 0.36
Methionine ^c	a	1.06 ± 0.29	1.29 ± 0.24	0.96 ± 0.20	1.30 ± 0.24	1.19 ± 0.22	1.21 ± 0.21	1.27 ± 0.25
	b	4.69 ± 0.39	0.66 ± 0.22	0.73 ± 0.21	0.72 ± 0.19	1.36 ± 0.26	0.88 ± 0.24	0.91 ± 0.20
Isoleucine ^c	a	3.32 ± 0.34	3.08 ± 0.30	2.98 ± 0.31	2.99 ± 0.33	2.82 ± 0.29	2.79 ± 0.34	2.69 ± 0.31
	b	—	—	—	—	—	—	—
Leucine ^c	a	5.16 ± 0.77	4.78 ± 0.69	4.53 ± 0.62	4.42 ± 0.57	4.36 ± 0.49	4.31 ± 0.48	4.02 ± 0.47
	b	—	—	—	—	—	—	—
Tyrosine ^c	a	4.12 ± 0.44	4.13 ± 0.42	3.87 ± 0.40	4.01 ± 0.42	3.94 ± 0.41	3.88 ± 0.39	3.62 ± 0.38
	b	4.88 ± 0.39	4.99 ± 0.42	3.44 ± 0.43	2.97 ± 0.37	4.67 ± 0.47	3.25 ± 0.33	3.17 ± 0.37
Phenylalanine ^c	a	4.02 ± 0.34	3.64 ± 0.36	3.51 ± 0.33	3.59 ± 0.34	3.40 ± 0.34	3.33 ± 0.33	3.16 ± 0.31
	b	4.89 ± 0.40	2.75 ± 0.37	2.06 ± 0.29	2.04 ± 0.30	4.44 ± 0.39	2.22 ± 0.27	2.11 ± 0.27
Lysine ^c	a	5.57 ± 0.31	5.51 ± 0.31	5.08 ± 0.33	4.78 ± 0.40	4.92 ± 0.29	4.99 ± 0.31	4.40 ± 0.36
	b	4.36 ± 0.29	2.82 ± 0.28	3.09 ± 0.27	3.06 ± 0.23	2.01 ± 0.27	2.49 ± 0.28	2.79 ± 0.30
Histidine ^c	a	1.93 ± 0.30	1.92 ± 0.28	1.77 ± 0.29	1.68 ± 0.29	1.71 ± 0.23	1.68 ± 0.30	1.49 ± 0.26
	b	—	—	—	—	—	—	—
Arginine ^c	a	5.30 ± 0.54	5.02 ± 0.54	4.75 ± 0.47	4.86 ± 0.44	5.05 ± 0.30	4.89 ± 0.41	4.55 ± 0.38
	b	4.06 ± 0.48	7.05 ± 0.46	3.30 ± 0.37	3.27 ± 0.34	2.35 ± 0.31	1.29 ± 0.29	1.84 ± 0.31
γ-aminobutyric acid	b	3.88 ± 0.33	2.41 ± 0.24	1.65 ± 0.23	1.63 ± 0.23	2.13 ± 0.20	1.63 ± 0.24	1.59 ± 0.30
Ornithine	b	1.37 ± 0.35	0.98 ± 0.22	0.81 ± 0.26	0.80 ± 0.24	1.58 ± 0.19	1.58 ± 0.21	1.18 ± 0.26
% Crude nitrogen		1.580 ± 0.32	1.507 ± 0.27	1.484 ± 0.31	1.476 ± 0.35	1.394 ± 0.35	1.376 ± 0.32	1.340 ± 0.33

^a Determination for nonoxidized sample. Each value is the mean of eight determinations ± SD.

^b (a) ion-exchange chromatography; (b) gas-liquid chromatography

^c Essential amino acids

exchange chromatography data but strongly disagreed with those obtained by GLC. This is not surprising since potatoes contain a rather reactive and unstable mix of compounds: free amino acids, proteins and carbohydrates among others. Both processing and analysis must have mild conditions if destruction is to be prevented.

As shown by ion-exchange chromatography, potato tuber processing into dehydrated granules by a freeze-thaw process caused only small losses in total amino acids and little change in their composition. There was only a 4.5% loss ($p < 0.05$) of total essential amino acids and a 2.1% loss ($p < 0.05$) in non-essential amino acids, excluding aspartic and glutamic acids, which comprised 48% peeled and sliced potatoes. In the granule process their content significantly decreased to 37.9% ($p < 0.05$), which was the only significant loss ($p < 0.05$) recorded by the granule process.

When the granule results are compared to potato flakes made by laboratory drum drying (Jaswal, 1973), the granule process appears much superior. In the granule process, essential amino acid destruction was less than half that for flakes, while the nonessential amino acid losses were similar and involved primarily significant losses ($p < 0.05$) of aspartic and glutamic acids. A previous study (Golan-Goldhirsh et al, 1982) revealed only the change of the free amino acid pool during processing. Peeled raw potato, after reaching the end product state, decreased ($p < 0.05$) its free amino acid pool by 40%.

The major loss of 27.3% ($p < 0.05$) occurred in the pre-cooking step. Slightly more than half of the losses were assigned to essential amino acids. These data confirm the recent works of Talley et al. (1983, 1984) and Sullivan et al. (1985) where it was found that some losses of arginine, aspartic, glutamic and aminobutyric acids occurred during hot water blanching and a significant loss of methionine occurred during drum drying. However, amino acids can be considered quite stable during potato flake production. No other data are available in the existing literature on a granule process.

CONCLUSIONS

THE GRANULE PROCESS had little effect on the total amino acids and their composition. Only 4.5% destruction of essential amino acids and a total loss of 2.1% of nonessential acids classified the process as least detrimental for commercial potato processing.

REFERENCES

- Augustin, J., Swanson, B.G., Pometto, S.F., Teitzel, C., Artz, W.E., and Huang, C.-P. 1979. Changes in nutrient composition of dehydrated potato products during commercial processing. *J. Food Sci.* 44: 216.
Blackburn, S. 1978. Amino acid analysis: an important technique. In: "Amino Acid Determination—Methods and Techniques," 2nd ed., S. Blackburn (Ed.). Marcel Dekker, Inc., New York.

PROCESSING EFFECT ON POTATOE AMINO ACIDS...

- Davies, A.M.C. 1977. The free amino acids of tubers of potato varieties grown in England and Ireland. *Potato Res.* 20: 9.
- Davies, M.G. and Thomas, A.J. 1973. An investigation of hydrolytic techniques for the amino acid analysis of food stuffs. *J. Sci. Food Agric.* 24: 1525.
- Duncan, D.B. 1955. Multiple range and multiple F tests. *Biometrics* 11: 1.
- Eppendorfer, W.H., Eggum, B.D., and Bille, S.W. 1979. Nutritive value of potato crude protein as influenced by manuring and amino acid composition. *J. Sci. Food Agric.* 30: 361.
- Fitzpatrick, T.J., Talley, E.A., Porter, W.L., and Murphy, H.J. 1964. Chemical composition of potatoes. III. Relationships between specific gravity and the nitrogenous constituents. *Am. Potato J.* 41(3): 75.
- Golan-Goldhirsh, A., Hogg, A.M., and Wolfe, F.H. 1982. Gas chromatographic analysis of the free amino acid pool of the potato and gas chromatography—mass spectrometry identification of γ -aminobutyric acid and ornithine. *J. Agric. Food Chem.* 30: 320.
- Golan-Goldhirsh, A. and Wolfe, F.H. 1979. Gas chromatographic analysis of protein amino acid N-heptafluorobutyl isopropyl esters. *Can. Inst. Food Sci. Technol. J.* 12: 123.
- Jaswal, A.S. 1973. Effects of various processing methods on free and bound amino acid contents of potatoes. *Am. Potato J.* 50: 86.
- Kirkman, M.A., Burrell, M.M., Lea, P.J., and Mills, W.R. 1980. Identification and measurement of homoserine by gas-liquid chromatography. *Anal. Biochem.* 101: 364.
- Kozempel, M.F., Sullivan, J.F., Della Monica, E.S., Egoville, M.J., Talley, E.A., Jones, W.J., and Craig, J.R., J.C. 1982. Application of leaching model to describe potato nutrient losses in hot water blanching. *J. Food Sci.* 47: 1519.
- Liu, T.Y. and Chang, Y.H. 1971. Hydrolysis of proteins with p-toluene-sulfonic acid-determination of tryptophan. *J. Biol. Chem.* 246: 2842.
- Moore, S. and Stein, W.H. 1954. Procedure for the chromatographic determination of amino acids on four per cent cross-linked sulfonated polystyrene resins. *J. Biol. Chem.* 211: 893.
- Moore, S., Spackman, D.H., and Stein, W.H. 1958. Chromatography of amino acids on sulfonated polystyrene resins—an improved system. *Anal. Chem.* 30: 1185.
- Ooraikul, B. 1977. Production of potato granules. U.S. patent 4,007,286.
- Rosen, H. 1957. A modified ninhydrin colorimetric analysis for amino acids. *Arch. Biochem. Biophys.* 67: 10.
- Simpson, R.J., Neuberger, M.R., and Liu, T.Y. 1976. Complete amino acid analysis of protein from a single hydrolysate. *J. Biol. Chem.* 251: 1936.
- Spackman, D.H., Stein, W.H., and Moore, S. 1958. Automatic recording apparatus for use in the chromatography of amino acids. *Anal. Chem.* 30: 1190.
- Sullivan, J.F., Kozempel, M.F., Egoville, M.J., and Talley, E.A. 1985. Loss of amino acids and water soluble vitamins during potato processing. *J. Food Sci.* 50: 1249.
- Synge, R.L. 1977. Free amino acids of potato tubers: a survey of published results set out according to potato variety. *Potato Res.* 20: 1.
- Tajima, M., Tadokoro-Yasui, S., Suzuki, T., Shinoda-Kenmochi, K., Kitano, T., Tsuchiya, K., and Fulenshima, H. 1978. Comparative study between gas chromatography and ion-exchange chromatography of amino acid analysis of foods. *Agric. Biol. Chem.* 42: 1949.
- Talley, E.A., Fitzpatrick, T.J., and Porter, W.L. 1964. Chemical composition of potatoes. IV. Relationship of the free amino acid concentrations to specific gravity and storage time. *Am. Potato J.* 41: 357.
- Talley, E.A., Fitzpatrick, T.J. and Porter, W.L. 1970. Chemical composition of potatoes. VIII. Effect of variety, location and year of growth on the content of nitrogen compounds. *Am. Potato J.* 47: 231.
- Talley, E.A., and Porter, W.L. 1968. New quantitative approach to the study of nonenzymatic browning. *J. Agric. Food Chem.* 16: 262.
- Talley, E.A., Toma, R.B., and Orr, P.H. 1983. Composition of raw and cooked potato peel and flesh: amino acid content. *J. Food Sci.* 48: 1360.
- Talley, E.A., Toma, R.B., and Orr, P.H. 1984. Amino acid composition of freshly harvested and stored potatoes. *Am. Potato J.* 61(5): 267.
- Toma, R.B., Augustin, J., Orr, P.H., True, R.H., Hogan, J.M., and Shaw, R.L. 1978. Changes in the nutrient composition of potatoes during home preparation: I. Proximate composition. *Am. Potato J.* 55: 639.
- Yamagata, S. 1983. Amino acids and proteins of raw and granulated potatoes. M. Sc. Thesis, Univ. of Alberta, Edmonton, Canada.
- Ms received 2/17/87; revised 6/29/87; accepted 8/7/87.

The authors greatly acknowledge the Alberta potato dehydration industry for financial assistance (Alberta, Edmonton, Canada) and Prof. Trakhtenberg and Mr. A. Gorinstein (Israel) for their help with the statistical studies.

QUALITY OPTIMIZATION IN ASEPTIC PROCESSING... From page 307

- of heat induced changes in food contents with the aid of reaction kinetics. *Zeit. Lebensmitteltechnol. und Verfahrenstechnik* 5: 372.
- Leonard, S.J., Luh, B.S., and Simone, M. 1964. Aseptic canning of foods. *Food Technol.* 18(1): 81.
- Leonard, S.J., Merson, R.L., Marsh, G.L., and Heil, J.R. 1986. Estimating thermal degradation in processing foods. *J. Agric. Food Chem.* 34(3): 392.
- Lund, D.B. 1982. Quantifying reactions influencing quality of foods: texture, flavor, and appearance. *J. Food Proc. Preserv.* 6: 133.
- Mansfield, T. 1962. High temperature-short time sterilization. *Proc. 1st Int'l Cong. Food Sci. and Technol.* London, 4: 311.
- Mansfield, T. 1974. A brief study of cooking. Quoted in Lund, D.B. (1975). In "Principles of Food Science. Part II. Physical Principles of Food Preservation." Fennema, O.W. (Ed.), Ch. 2. Marcel Dekker, New York, NY.
- Nelson, P.R. 1983. Stability prediction using Arrhenius model. *Comm. Programs in Biomedicine* 16: 55.
- Ohlsson, T. 1980a. Temperature dependence of sensory quality changes during thermal processing. *J. Food Sci.* 45: 836.
- Ohlsson, T. 1980b. Optimal sterilization temperatures for flat containers. *J. Food Sci.* 45: 848.
- Ohlsson, T. 1980c. Optimal sterilization temperatures for sensory quality in cylindrical containers. *J. Food Sci.* 45: 1517.
- Rao, M.A., Lee, C.Y., Katz, J., and Cooley, H.J. 1981. A kinetic study of the loss of vitamin C color and firmness during thermal processing of canned peas. *J. Food Sci.* 46: 636.
- Reichert, J.E. 1974. Optimale sterilizationstemperaturen für Fertiggerichte. *Die Fleischwirtschaft* 54: 1305.
- Reichert, J.E. 1977. Der C-wet als Hilfsmittel zur Prozessoptimierung. *Zeit für Lebensm. — tech und Vergahr* 28: 1.
- Reuter, H. 1980. Evaluation of thermic effectiveness of ultrahigh temperature plants of milk. *Milchwissenschaft* 35(9): 536. [In *Biol Abstr.* 71(7):44872].
- Saguy, I. 1983. Optimization of dynamic systems. In "Computer-Aided Techniques in Food Technology," Saguy, I. (Ed.). Marcel Dekker, New York, NY.
- Saguy, I. and Karel, M. 1979. Optimal retort temperature profile in optimizing thiamine retention in conduction-type heating of canned foods. *J. Food Sci.* 44: 1485.
- Saguy, I. and Karel, M. 1986. Index of deterioration and prediction of quality losses. In "Objective Methods in Food Quality Assessment," Kapsalis, J.G. (Ed.), C.R.C. Press, Boca Raton, FL.
- Saguy, I., Mishkin, M., and Karel, M. 1983. A review: Optimization methods and available software. *CRC Crit. Rev. Food Technol.* 20(4): 275.
- Scott, V.N. 1985. Aseptic processing and packaging of particulates commercial sterility and enzymatic considerations. *Proc. of National Food Processors Association Conf.*, Washington, DC. April 11–12, 1985. p. 47.
- Swartzel, K.R. 1982. Arrhenius kinetics as applied to product constituent losses in ultra high temperature processing. *J. Food Sci.* 47: 1886.
- Swartzel, K.R. 1986. Equivalent-point method for thermal evaluation of continuous flow systems. *J. Agric. Food Chem.* 34(3): 396.
- Toledo, R.T. 1986. Postprocessing changes in aseptically packed beverages. *J. Agric. Food Chem.* 34(3): 405.
- Villota, R. and Hawkes, J.G. 1983. Effect of processing on kinetics of nutrients and organoleptic changes in foods. 1983 Winter Meeting, American Society of Agricultural Engineers, Dec. 13–16, Chicago, IL.
- von Bokelmann, B.A.H. and von Bokelmann, I.L.I. 1986. Aseptic packaging of liquid food products: A review. *J. Agric. Food Chem.* 34(3): 384.
- Ms received 4/8/87; revised 7/10/87; accepted 8/3/87.